

Measles Information Sheets



Our colleagues at the Ohio Department of Health Vaccine Preventable Disease program have requested that we share this information with you about measles. There are two fact sheets for parents, with each being two pages long. One is for pre-K settings, while the other is for the K-12 setting. We have copied the language on the forms below, as well as the images of the pages. Please feel free to share which ever is appropriate for your setting.

For ECE Settings:

Measles: What Parents of Children Attending Early Childcare in Ohio Should Know

Measles is not just a little rash.

It can spread very quickly person-to-person **through the air** when an infected person breathes, coughs, or sneezes. Even brief contact or being in a shared space with someone who has measles can put you at high risk for getting the disease if you are not protected.

One person with measles can spread it to nine (9) out of 10 unprotected people around them.

Symptoms Include:

- High Fever.
- Cough.
- Runny nose.
- Red, watery eyes.
- Rash—usually starts on the head and spreads down to the rest of the body.

Measles is contagious from four days before the rash appears through four days after the rash appears.

Measles can result in a medical complication for approximately three (3) out of 10 people who are infected.

Complications can include:

- Ear infections.
- Pneumonia (infection of the lungs).
- Encephalitis (swelling of the brain, which can lead to seizures).
- Diarrhea (which can lead to dehydration).
- Death.

DID YOU KNOW?

Measles virus can remain in the air for up to two (2) hours after an infected person leaves an area.

What is the treatment for measles?

- There is no specific antiviral therapy for measles.
- Medical care is supportive and helps to relieve symptoms or complications, such as bacterial infections.

Two doses of the measles, mumps, and rubella (MMR) vaccine is the best way to protect your child against measles.

- First dose at 12 to 15 months.
- Second dose at 4 to 6 years.

MMR can be given to children as young as 6 months of age who are at high risk of exposure, such as during a **community outbreak** or **international travel**.

If MMR vaccine is given between 6–11 months, two additional doses are still needed after 12 months of age.

Two doses are about 97% effective at preventing measles. One dose is about 93% effective at preventing measles.

Early childcare vaccine requirements in Ohio.

- Ohio Revised Code 5104.014 requires children to be fully protected against 14 vaccine preventable diseases, including measles.

Measles vaccines are available at:

- Many doctor's offices.
- Local Health Departments.
- Pharmacies (check with location based on your child's age).

How to protect babies who are too young for the MMR vaccine:

- Ensure (to the extent possible) that those around your baby are vaccinated.
- Limit your baby's exposure to crowds, other children, and anyone with cold symptoms.
- Keep your child home when they are sick.
- Prevent germs at home. Regularly disinfect objects and surfaces in your home.
- Practice good hand hygiene. Wash hands using soap and water and scrub for at least 20 seconds.

Seek care with your child's doctor, urgent care, or emergency room if you think your child has measles. Make sure to call the healthcare facility before you arrive (if possible) to inform them that you believe your child has measles so they can provide care to your child while protecting others.

Measles:

What Parents of Children Attending Early Childcare in Ohio Should Know



Measles is not just a little rash.

It can spread very quickly person-to-person **through the air** when an infected person breathes, coughs, or sneezes. Even brief contact or being in a shared space with someone who has measles can put you at high risk for getting the disease if you are not protected.

One person with measles can spread it to nine (9) out of 10 unprotected people around them.

Symptoms Include:

- High Fever.
- Cough.
- Runny nose.
- Red, watery eyes.
- Rash—usually starts on the head and spreads down to the rest of the body.

Measles is contagious from four days before the rash appears through four days after the rash appears.

Measles can result in a medical complication for approximately three (3) out of 10 people who are infected.

Complications can include:

- Ear infections.
- Pneumonia (infection of the lungs).
- Encephalitis (swelling of the brain, which can lead to seizures).
- Diarrhea (which can lead to dehydration).
- Death.



DID YOU KNOW?

Measles virus can remain in the air for up to two (2) hours after an infected person leaves an area.

What is the treatment for measles?

- There is no specific antiviral therapy for measles.
- Medical care is supportive and helps to relieve symptoms or complications, such as bacterial infections.

Two doses of the measles, mumps, and rubella (MMR) vaccine is the best way to protect your child against measles.

- First dose at 12 to 15 months.
- Second dose at 4 to 6 years.

MMR can be given to children as young as 6 months of age who are at high risk of exposure, such as during a **community outbreak** or **international travel**.

If MMR vaccine is given between 6–11 months, two additional doses are still needed after 12 months of age.

Two doses are about 97% effective at preventing measles. One dose is about 93% effective at preventing measles.

Early childcare vaccine requirements in Ohio.

- **Ohio Revised Code 5104.014** requires children to be fully protected against 14 vaccine-preventable diseases, including measles.

Measles vaccines are available at:

- Many doctor's offices.
- [Local Health Departments](#).
- Pharmacies (check with location based on your child's age).

How to protect babies who are too young for the MMR vaccine:

- Ensure (to the extent possible) that those around your baby are vaccinated.
- Limit your baby's exposure to crowds, other children, and anyone with cold symptoms.
- Keep your child home when they are sick.
- Prevent germs at home. Regularly disinfect objects and surfaces in your home.
- Practice good hand hygiene. Wash hands using soap and water and scrub for at least 20 seconds.

Seek care with your child's doctor, urgent care, or emergency room if you think your child has measles. Make sure to call the healthcare facility before you arrive (if possible) to inform them that you believe your child has measles so they can provide care to your child while protecting others.

May 2025

For K-12 settings:

Measles: What Parents of Students Attending Kindergarten through 12th Grade in Ohio Should Know

Measles is not just a little rash.

It can spread very quickly person-to-person through the air when an infected person breathes, coughs, or sneezes. Even brief contact or being in a shared space with someone who has measles can put you at high risk for getting the disease if you are not protected.

One person with measles can spread it to nine (9) out of 10 unprotected people around them.

Symptoms Include:

- High Fever.
- Cough.
- Runny nose.
- Red, watery eyes.
- Rash—usually starts on the head and spreads down to the rest of the body.

Measles is contagious from four days before the rash appears through four days after the rash appears.

Measles can result in a medical complication for approximately three (3) out of 10 people who are infected.

Complications can include:

- Ear infections.
- Pneumonia (infection of the lungs).
- Encephalitis (swelling of the brain, which can lead to seizures).
- Diarrhea (which can lead to dehydration).
- Death.

DID YOU KNOW?

Measles virus can remain in the air for up to two (2) hours after an infected person leaves an area.

What is the treatment for measles?

- There is no specific antiviral therapy for measles.
- Medical care is supportive and helps to relieve symptoms or complications, such as bacterial infections.

Two doses of the measles, mumps, and rubella (MMR) vaccine is the best way to protect your child against measles.

- First dose at 12 to 15 months.
- Second dose at 4 to 6 years.

It's not too late to vaccinate.

If your child is over 6 years old and hasn't received all their recommended vaccines, don't worry—there's still time. Talk to your healthcare provider or local health department to find out what your child needs to stay protected, especially against measles.

Two doses are about 97% effective at preventing measles. One dose is about 93% effective at preventing measles.

A healthy school year starts with protection!

Measles vaccines are available at:

- Many doctor's offices.
- Local Health Departments.
- Pharmacies (check with location based on your child's age).

K-12 school vaccine requirements in Ohio:

- Ohio Revised Code 3313.671 requires students to be fully protected against 10 vaccine preventable diseases, including measles.
- During an outbreak at school, children without proof of two (2) MMR vaccine doses or previous measles disease will be excluded from school until documentation is provided. In an outbreak situation, a written statement of previous measles disease is acceptable in lieu of vaccination only if it is signed by a physician and gives the month and year of infection. Children with written waivers or exemptions shall be excluded for 21 days after the onset of the last known case in the affected school. (Director's Journal-School Requirements).

Seek care with your child's doctor, urgent care, or emergency room if you think your child has measles. Make sure to call the healthcare facility before you arrive (if possible) to inform them that you believe your child has measles so they can provide care to your child while protecting others.

Measles:

What Parents of Students Attending Kindergarten through 12th Grade in Ohio Should Know

Measles is not just a little rash.

It can spread very quickly person-to-person **through the air** when an infected person breathes, coughs, or sneezes. Even brief contact or being in a shared space with someone who has measles can put you at high risk for getting the disease if you are not protected.

One person with measles can spread it to nine (9) out of 10 unprotected people around them.

Symptoms Include:

- High Fever.
- Cough.
- Runny nose.
- Red, watery eyes.
- Rash—usually starts on the head and spreads down to the rest of the body.

Measles is contagious from four days before the rash appears through four days after the rash appears.

Measles can result in a medical complication for approximately three (3) out of 10 people who are infected.

Complications can include:

- Ear infections.
- Pneumonia (infection of the lungs).
- Encephalitis (swelling of the brain, which can lead to seizures).
- Diarrhea (which can lead to dehydration).
- Death.



DID YOU KNOW?

Measles virus can remain in the air for up to two (2) hours after an infected person leaves an area.

What is the treatment for measles?

- There is no specific antiviral therapy for measles.
- Medical care is supportive and helps to relieve symptoms or complications, such as bacterial infections.

Two doses of the measles, mumps, and rubella (MMR) vaccine is the best way to protect your child against measles.

- First dose at 12 to 15 months.
- Second dose at 4 to 6 years.

It's not too late to vaccinate.

If your child is over 6 years old and hasn't received all their recommended vaccines, don't worry—there's still time. Talk to your healthcare provider or local health department to find out what your child needs to stay protected, especially against measles.

Two doses are about 97% effective at preventing measles. One dose is about 93% effective at preventing measles.

A healthy school year starts with protection!

Measles vaccines are available at:

- Many doctor's offices.
- [Local Health Departments](#).
- Pharmacies (check with location based on your child's age).

K-12 school vaccine requirements in Ohio:

- **Ohio Revised Code 3313.671** requires students to be fully protected against 10 vaccine-preventable diseases, including measles.
- During an outbreak at school, children without proof of two (2) MMR vaccine doses or previous measles disease will be excluded from school until documentation is provided. In an outbreak situation, a written statement of previous measles disease is acceptable in lieu of vaccination only if it is signed by a physician and gives the month and year of infection. Children with written waivers or exemptions shall be excluded for 21 days after the onset of the last known case in the affected school. ([Director's Journal-School Requirements](#)).

Seek care with your child's doctor, urgent care, or emergency room if you think your child has measles. Make sure to call the healthcare facility before you arrive (if possible) to inform them that you believe your child has measles so they can provide care to your child while protecting others.