

# Ohio Trauma Registry 2026

## Trauma Acute Care Registry Data Dictionary

Version 2026.0 (01/01/2026)

*This edition is effective for all trauma patients  
presenting for treatment on or after January 1, 2026.*



Department of  
Public Safety



## ACKNOWLEDGEMENTS

The Ohio State Board of Emergency Medical, Fire and Transportation Services and the EMS Division of the Ohio Department of Public Safety would like to thank the myriad of people – too numerous to list here – who have worked tirelessly to create, expand and transform the Ohio Trauma Registry from its inception and embryonic beginnings in the late 1990s into the powerful research and policymaking tool it is today. This growth and development would not have been possible without the strength of their combined knowledge, wisdom and hard work.

*Trauma Acute Care Registry (TACR) is a component of the Ohio Trauma Registry (OTR) and is maintained by the Ohio Department of Public Safety, 1970 W. Broad St., Columbus, Ohio 43223. For more information about the TACR, OTR and/or the State of Ohio's Trauma System, contact the Ohio Department of Public Safety, Division of EMS, Research and Analysis Section, at (800)233-0785, EMSdata@dps.ohio.gov or visit [www.ems.ohio.gov](http://www.ems.ohio.gov).*

## **Table of Contents**

ACKNOWLEDGEMENTS.....	2
STATEMENT ABOUT ITDX / TECHNICAL STANDARDS VS. CLINICAL STANDARDS.....	9
NATIONAL ELEMENTS THAT WILL NOT BE COLLECTED IN THE OHIO TRAUMA ACUTE CARE REGISTRY.....	10
OHIO SPECIFIC ELEMENTS .....	11
DIFFERENCES BETWEEN OHIO AND NATIONAL TRAUMA DATA STANDARD (NTDS).....	12
TACR INCLUSION/EXCLUSION CRITERIA – ICD-10 .....	15
OTR TACR INCLUSION/EXCLUSION DECISION TREE – ICD-10 .....	17
COMMON NULL VALUES .....	18
HOSPITAL CODE.....	19
UNIQUE ADMISSION NUMBER .....	20
TRAUMA TRACKING NUMBER .....	21
FACILITY TYPE.....	22
PATIENT’S HOME CITY.....	23
PATIENT’S HOME STATE.....	24
PATIENT’S HOME COUNTY .....	25
PATIENT’S HOME ZIP CODE .....	26
PATIENT’S HOME COUNTRY .....	27
ALTERNATE HOME RESIDENCE .....	28
DATE OF BIRTH.....	29
AGE .....	30
AGE UNITS .....	31
SEX ASSIGNED AT BIRTH .....	32
GENDER .....	33
GENDER-AFFIRMING HORMONE THERAPY.....	34
RACE.....	35
ETHNICITY .....	36
PRIMARY ICD-10 EXTERNAL CAUSE CODE.....	37
ADDITIONAL ICD-10 EXTERNAL CAUSE CODE .....	38
ICD-10 PLACE OF OCCURRENCE EXTERNAL CAUSE CODE.....	39
WORK-RELATED.....	40
PATIENT’S OCCUPATIONAL INDUSTRY .....	41

PATIENT'S OCCUPATION .....	42
INJURY INCIDENT DATE .....	43
INJURY INCIDENT TIME .....	44
INCIDENT CITY .....	45
INCIDENT STATE .....	46
INCIDENT COUNTY .....	47
INCIDENT LOCATION ZIP CODE .....	48
INCIDENT COUNTRY .....	49
PROTECTIVE DEVICES .....	50
CHILD SPECIFIC RESTRAINT .....	51
AIRBAG DEPLOYMENT .....	52
TRANSPORT MODE FOR ARRIVAL AT YOUR HOSPITAL .....	53
TRANSPORT AGENCY .....	54
OTHER TRANSPORT MODES .....	55
EMS PATIENT CARE REPORT UNIQUE IDENTIFIER (UUID) .....	56
EMS DISPATCH DATE TO SCENE OR TRANSFERRING FACILITY .....	57
EMS DISPATCH TIME TO SCENE OR TRANSFERRING FACILITY .....	58
EMS UNIT ARRIVAL DATE AT SCENE OR TRANSFERRING FACILITY .....	59
EMS UNIT ARRIVAL TIME AT SCENE OR TRANSFERRING FACILITY .....	60
EMS UNIT DEPARTURE DATE FROM SCENE OR TRANSFERRING FACILITY .....	61
EMS UNIT DEPARTURE TIME FROM SCENE OR TRANSFERRING FACILITY .....	62
INITIAL FIELD SYSTOLIC BLOOD PRESSURE .....	63
INITIAL FIELD PULSE RATE .....	64
INITIAL FIELD RESPIRATORY RATE .....	65
INITIAL FIELD OXYGEN SATURATION .....	66
INITIAL FIELD GCS - EYE .....	67
INITIAL FIELD GCS - VERBAL .....	68
INITIAL FIELD GCS - MOTOR .....	69
INITIAL FIELD GCS - TOTAL .....	70
INITIAL FIELD GCS QUALIFIER .....	71
SCENE INTERVENTIONS .....	72
PREHOSPITAL CARDIAC ARREST .....	73

INTUBATION PRIOR TO ARRIVAL .....	74
INTUBATION LOCATION.....	75
INTER-FACILITY TRANSFER .....	76
TRANSFERRING HOSPITAL CODE.....	77
ED/HOSPITAL ARRIVAL DATE.....	78
ED/HOSPITAL ARRIVAL TIME .....	79
HIGHEST ACTIVATION .....	80
TRAUMA SURGEON ARRIVAL DATE.....	81
TRAUMA SURGEON ARRIVAL TIME .....	82
INITIAL ED/HOSPITAL SYSTOLIC BLOOD PRESSURE .....	83
INITIAL ED/HOSPITAL PULSE RATE .....	84
INITIAL ED/HOSPITAL RESPIRATORY RATE.....	85
INITIAL ED/HOSPITAL RESPIRATORY ASSISTANCE .....	86
INITIAL ED/HOSPITAL OXYGEN SATURATION.....	87
INITIAL ED/HOSPITAL SUPPLEMENTAL OXYGEN .....	88
INITIAL ED/HOSPITAL TEMPERATURE .....	89
INITIAL ED/HOSPITAL GCS - EYE .....	90
INITIAL ED/HOSPITAL GCS - VERBAL.....	91
INITIAL ED/HOSPITAL GCS - MOTOR .....	92
INITIAL ED/HOSPITAL GCS - TOTAL.....	93
INITIAL ED/HOSPITAL GCS ASSESSMENT QUALIFIERS .....	94
HEIGHT .....	95
WEIGHT .....	96
ED DISCHARGE ORDER WRITTEN DATE.....	97
ED DISCHARGE ORDER WRITTEN TIME .....	98
ED DISCHARGE DATE.....	99
ED DISCHARGE TIME .....	100
ED DISCHARGE DISPOSITION .....	101
ED TRANSFER TO HOSPITAL.....	102
PRIMARY TRAUMA SERVICE TYPE.....	103
PRIMARY MEDICAL EVENT.....	104
ALCOHOL SCREEN .....	105

ALCOHOL SCREEN RESULTS .....	106
DRUG SCREEN .....	107
ICD-10 HOSPITAL PROCEDURES .....	108
PROCEDURE LIST FOR HOSPITAL PROCEDURE ELEMENTS .....	109
PROCEDURE EPISODE.....	110
HOSPITAL PROCEDURE START DATE.....	111
HOSPITAL PROCEDURE START TIME.....	112
ADVANCE DIRECTIVE LIMITING CARE.....	113
ALCOHOL USE DISORDER .....	115
ANTICOAGULANT THERAPY.....	117
ATTENTION DEFICIT DISORDER/ATTENTION DEFICIT HYPERACTIVITY DISORDER (ADD/ADHD) .....	119
AUTISM SPECTRUM DISORDER (ASD) (PRE-EXISTING CONDITIONS).....	121
BIPOLAR I/II DISORDER .....	123
BLEEDING DISORDER .....	125
BRONCHOPULMONARY DYSPLASIA/CHRONIC LUNG DISEASE.....	127
CEREBRAL VASCULAR ACCIDENT (CVA) .....	129
CHRONIC OBSTRUCTIVE PULMONARY DISEASE (COPD).....	131
CHRONIC RENAL FAILURE.....	133
CIRRHOSIS.....	135
CONGENITAL ANOMALIES.....	137
CONGESTIVE HEART FAILURE (CHF).....	139
CURRENT SMOKER .....	141
CURRENTLY RECEIVING CHEMOTHERAPY FOR CANCER .....	143
DEMENTIA .....	145
DIABETES MELLITUS.....	147
DISSEMINATED CANCER.....	149
FUNCTIONALLY DEPENDENT HEALTH STATUS.....	151
HYPERTENSION .....	153
MAJOR DEPRESSIVE DISORDER .....	155
MYOCARDIAL INFARCTION (MI).....	157
OTHER MENTAL/PERSONALITY DISORDERS.....	159
PERIPHERAL ARTERIAL DISEASE (PAD).....	161

POST-TRAUMATIC STRESS DISORDER.....	163
PREGNANCY .....	165
PREMATURITY .....	167
SCHIZOAFFECTIVE DISORDER.....	169
SCHIZOPHRENIA .....	171
STEROID USE .....	173
SUBSTANCE USE DISORDER.....	175
VENTILATOR DEPENDENCE .....	178
DNR STATUS.....	180
ICD-10 INJURY DIAGNOSES.....	181
AIS CODE.....	182
AIS VERSION .....	183
INJURY SEVERITY SCORE.....	184
TOTAL ICU LENGTH OF STAY .....	185
TOTAL VENTILATOR DAYS .....	186
HOSPITAL DISCHARGE ORDER WRITTEN DATE.....	187
HOSPITAL DISCHARGE ORDER WRITTEN TIME .....	188
HOSPITAL DISCHARGE DATE .....	189
HOSPITAL DISCHARGE TIME .....	190
HOSPITAL DISCHARGE DISPOSITION .....	191
INPATIENT TRANSFER TO HOSPITAL.....	192
DISCHARGE STATUS .....	193
DATE OF DEATH.....	194
PRIMARY METHOD OF PAYMENT .....	195
AUTOPSY PERFORMED .....	196
ACUTE KIDNEY INJURY (AKI).....	197
ACUTE RESPIRATORY DISTRESS SYNDROME (ARDS).....	199
ALCOHOL WITHDRAWAL SYNDROME .....	202
CARDIAC ARREST WITH CPR .....	204
CATHETER-ASSOCIATED URINARY TRACT INFECTION (CAUTI).....	206
CENTRAL LINE-ASSOCIATED BLOODSTREAM INFECTION (CLABSI) .....	210
DEEP SURGICAL SITE INFECTION .....	215

DEEP VEIN THROMBOSIS (DVT) .....	219
DELIRIUM.....	221
MYOCARDIAL INFARCTION (MI).....	223
ORGAN/SPACE SURGICAL SITE INFECTION .....	225
OSTEOMYELITIS.....	228
PULMONARY EMBOLISM (PE).....	230
PRESSURE ULCER .....	232
SEVERE SEPSIS .....	234
STROKE/CVA .....	236
SUPERFICIAL INCISIONAL SURGICAL SITE INFECTION.....	240
UNPLANNED ADMISSION TO ICU .....	243
UNPLANNED INTUBATION .....	245
UNPLANNED RETURN TO THE OPERATING ROOM .....	247
VENTILATOR-ASSOCIATED PNEUMONIA (VAP) .....	249
Appendix A - Discharge Disposition Definitions .....	263
Appendix B - Calculating ICU Length of Stay and Ventilator Days.....	264
Appendix C - Glossary of Abbreviations .....	265
Appendix D – Acronyms.....	266
Appendix E – Ohio Regional Trauma System Data Dictionary .....	267
Appendix F – CHANGE LOG for the Ohio Trauma Acute Care Data (TACR) .....	268



## STATEMENT ABOUT ITDX / TECHNICAL STANDARDS VS. CLINICAL STANDARDS

---

The State of Ohio recognizes the ITDX as the transmission standard. The Ohio Trauma Acute Care Registry Data Dictionary reflects the American College of Surgeons (ACS) reporting requirements adopted by the State of Ohio for 2025. The manner of end-point collection is left to the trauma vendor(s) provided that these vendors are able to meet both State and ACS reporting requirements.

## NATIONAL ELEMENTS THAT WILL NOT BE COLLECTED IN THE OHIO TRAUMA ACUTE CARE REGISTRY

---

**The following elements will not be collected and should be defaulted to Not Applicable.**

- Initial ED/Hospital GCS 40 – Eye
- Initial ED/Hospital GCS 40 – Verbal
- Initial ED/Hospital GCS 40 – Motor

## OHIO SPECIFIC ELEMENTS

---

- Hospital Code
- Unique Admission Number
- Trauma Tracking Number
- Facility Type
- Transport Agency
- EMS Dispatch Date
- EMS Dispatch Time
- EMS Unit Arrival Date at Scene or Transferring Facility
- EMS Unit Arrival Time at Scene or Transferring Facility
- EMS Unit Departure Date From Scene or Transferring Facility
- EMS Unit Departure Time From Scene or Transferring Facility
- Initial Field Systolic Blood Pressure
- Initial Field Pulse Rate
- Initial Field Respiratory Rate
- Initial Field Oxygen Saturation
- Initial Field GCS Eye
- Initial Field GCS Verbal
- Initial Field GCS Motor
- Initial Field GCS Total
- Initial Field GCS Qualifier
- Scene Interventions
- Transferring Hospital Code
- ED Discharge Order Written Date
- ED Discharge Order Written Time
- ED Transfer to Hospital
- Procedure Episode
- DNR Status
- Injury Severity Score
- Hospital Discharge Order Written Date
- Hospital Discharge Order Written Time
- Inpatient Transfer To Hospital
- Discharge Status
- Date of Death
- Autopsy Performed

## DIFFERENCES BETWEEN OHIO AND NATIONAL TRAUMA DATA STANDARD (NTDS)

---

### Inclusion Criteria Differences

- Ohio follows NTDS Inclusion Criteria with exceptions:
  - Ohio INCLUDES: S00, S10, S20, S30, S40, S50, S60, S70, S80, S90 (Patients with these isolated injuries that were transferred in/out or died.) (Patients transferred/discharged to hospice (e.g., hospice facility, hospice unit, home hospice)
    - Note that these codes are excluded when patients with these isolated injuries were **NOT** transferred in/out or died, per NTDS.
  - Ohio INCLUDES:
    - J70.5
    - T20-28
    - T30-32
    - T33
    - T34
    - T59.81
    - T67
    - T68
    - T69
    - T70.4
    - T70.8
    - T70.9
    - T71
    - T74.1
    - T74.4
    - T75
    - T75.1
    - T75.4

#### Inclusion criteria –

- **NON-TRAUMA CENTERS:** S00, S10, S20, S30, S40, S50, S60, S70, S80, S90 (Patients with these isolated injuries that were transferred in/ out (FOR TRAUMA EVALUATION) or died) –Centers only capture patients transferred for trauma care. This would exclude patients being transferred to another non-trauma center/ acute care facility for medical purposes.

**TRAUMA CENTERS:** S00, S10, S20, S30, S40, S50, S60, S70, S80, S90 (Patients with these isolated injuries that were transferred in or died THAT WERE EVALUATED BY TRAUMA SERVICE) –Centers only capture patients transferred for trauma care

#### Exclusion criteria –

- **NON-TRAUMA CENTERS:** S00, S10, S20, S30, S40, S50, S60, S70, S80, S90 (Patients with these isolated injuries that were transferred in/ out (**NOT** FOR TRAUMA EVALUATION) or died) –Centers only capture patients transferred for trauma care. This would exclude patients being transferred to another non-trauma center/ acute care facility for medical purposes.
- **TRAUMA CENTERS:** S00, S10, S20, S30, S40, S50, S60, S70, S80, S90 (Patients with these isolated injuries that were transferred in or died THAT WERE **NOT** EVALUATED BY TRAUMA SERVICE) –Centers only capture patients transferred for trauma care.

### OH Definition Is Different Than NTDS

- Height
  - OH does not include “within 24 hours or less of ED/Hospital arrival” in definition
- Weight

- OH does not include “within 24 hours or less of ED/Hospital arrival” in definition

#### **OH Additional Information Is Different Than NTDS**

- Patient Home City
  - OH does not include “Only reported when patients home zip postal code is not known not recorded and country is US.”
  - OH does not include “Null value NA is reported if patients home zip / postal code is reported.”
- Patient Home State
  - OH does not include “Only reported when patient home zip / postal code is not known / not recorded and country is US.”
  - OH does not include “Null value NA is reported if patients home zip / postal code is reported.”
- Patient Home County
  - OH does not include “Only reported when patient home zip / postal code is not known not recorded and country is US.”
- Highest Activation
  - Non-Trauma Center Facilities report the null value “Not Applicable” (NA)
  - OH does not include “Null value NA is reported if patients home zip / postal code is reported.”
- Patient Home Zip Code
  - OH does not include “May require adherence to HIPAA regulations.”
- Age
  - OH does not include the “Null value not applicable is reported if date of birth is reported.”
- Age Units
  - OH does not include the “Null value not applicable is reported if date of birth is reported.”
- Incident City
  - OH does not include “Only recorded when incident location zip / postal code is not known / not recorded and country is US.”
  - OH does not include the “Null value not applicable is reported if incident location zip / postal code is reported.”
- Incident State
  - OH does not include “Only recorded when incident location zip / postal code is not known / not recorded and country is US.”
  - OH does not include the “Null value not applicable is reported if incident location zip / postal code is reported.”
- Incident County
  - OH does not include “Only recorded when incident location zip / postal code is not known / not recorded and country is US.”
  - OH does not include the “Null value not applicable is reported if incident location zip postal code is reported.”
- Incident Zip Code
  - NTDS says “Can be stored as a 5 or 9-digit code (XXXXX-XXXX) for US or CA and can be stored in the postal code format of the applicable country.”
  - Ohio says “Stored as a five-digit code (XXXXX)”
- Transport mode for arrival at your hospital
  - OH added examples
- Other Transport Modes
  - OH added examples
- Height
  - OH does not include “...within 24 hours or less of ED/Hospital arrival” in the 4<sup>th</sup> bullet point
- Weight
  - OH does not include “...within 24 hours or less of ED/Hospital arrival” in the 4<sup>th</sup> bullet point
- Hospital Procedure State Date
  - OH added “Linked to hospital procedures element”

- Hospital Procedure Start Time
  - OH added “Linked to hospital procedures element”
  - OH added “If distinct procedures with the same procedure code are performed, their start time must be different”

#### **Other Element Name and Definition Differences**

- ED Discharge Order Written Date
- ED Discharge Order Written Time
  - These are Ohio specific elements. However, they match in definition to NTDS ED Discharge Date and ED Discharge Time
- ED Discharge Date
- ED Discharge Time
  - These are NTDS elements, however the Ohio definition is different
- Hospital Discharge Order Written Date
- Hospital Discharge Order Written Time
  - These are Ohio specific elements. However, they match in definition to NTDS Hospital Discharge Date and Hospital Discharge Time
- Hospital Discharge Date
- Hospital Discharge Time
  - These are NTDS elements, however the Ohio definition is different

#### **Element Value Differences**

- ED Discharge Disposition
  - Ohio added “12 Interventional Radiology (IR)”
- Primary Method of Payment
  - Ohio added “8 Workers Compensation”

#### **Edit Check Differences**

- For element Hospital Procedure Start Date, the following edit check should not be present:
  - 6607 Hospital Procedure Start Date is later than Hospital Discharge Order Written Date. (Note: NTDS refers to this field as Hospital Discharge Date – Ohio has a different definition for this field.)
- For element Hospital Procedure Start Time, the following edit check should not be present:
  - 6707 Hospital Procedure Start Time is later than Hospital Discharge Order Written Time. (Note: NTDS refers to this field as Hospital Discharge Time – Ohio has a different definition for this field.)

**NOTE:** Reference to this section is included on each individual element page that is affected by the differences listed.

## TACR INCLUSION/EXCLUSION CRITERIA – ICD-10

---

### TRAUMA PATIENT DEFINITION

To ensure consistent data collection across the State of Ohio and to follow the National Trauma Data Standard, a trauma patient is defined as a patient sustaining a traumatic injury within 14 days of initial hospital encounter\*\*\* and meeting the following criteria:

### PATIENT INCLUSION CRITERIA

To be included in the Trauma Acute Care Registry (TACR):

The patient must have incurred at least one of the injury diagnostic codes defined in the International Classification of Diseases, Tenth Revision, Clinical Modification (ICD-10-CM):

- **J70.5 with character modifier of A ONLY** (Respiratory conditions due to smoke inhalation – initial encounter)
- **S00-S99 with 7<sup>th</sup> character modifier of A, B or C ONLY** (Injuries to specific body parts – initial encounter):
  - **T07** (Unspecified multiple injuries);
  - **T14** (Injury of unspecified body region);
  - **T20-T28 with 7<sup>th</sup> character modifier of A ONLY** (Burns by specified body parts – initial encounter);
  - **T30-T32** (Burn by TBSA percentage);
  - **T33 with character modifier of A ONLY** (Superficial frostbite – initial encounter)
  - **T34 with character modifier of A ONLY** (Frostbite with tissue necrosis – initial encounter)
  - **T59.81 with 7<sup>th</sup> character modifier of A ONLY** (Toxic effect of smoke - initial encounter)
  - **T67 with character modifier of A ONLY** (Effects of heat and light – initial encounter)
  - **T68 with character modifier of A ONLY** (Hypothermia – initial encounter)
  - **T69 with character modifier of A ONLY** (Other effects of reduced temperature – initial encounter)
  - **T70.4 with character modifier of A ONLY** (Effects of high-pressure fluids – initial encounter)
  - **T70.8 with character modifier of A ONLY** (Other effects of air pressure and water pressure – initial encounter)
  - **T70.9 with character modifier of A ONLY** (Effect of air pressure and water pressure, unspecified – initial encounter)
  - **T71 with character modifier of A ONLY** (Asphyxiation – initial encounter)
  - **T74.1 with character modifier of A ONLY** (Physical abuse, confirmed – initial encounter)
  - **T74.4 with character modifier of A ONLY** (Shaken infant syndrome – initial encounter)
  - **T75.0 with character modifier of A ONLY** (Effects of lightning – initial encounter)
  - **T75.1 with character modifier of A ONLY** (Unspecified effects of drowning and nonfatal submersion – initial encounter)
  - **T75.4 with character modifier of A ONLY** (Electrocution – initial encounter)
  - **T79.A1-T79.A9 with 7<sup>th</sup> character modifier of A ONLY** (Traumatic compartment syndrome – initial encounter)
- **NON-TRAUMA CENTERS: S00, S10, S20, S30, S40, S50, S60, S70, S80, S90** (Patients with these isolated injuries that were transferred in/ out (FOR TRAUMA EVALUATION) or died) –Centers only capture patients transferred for trauma care. This would exclude patients being transferred to another non-trauma center/ acute care facility for medical purposes.

- **TRAUMA CENTERS:** S00, S10, S20, S30, S40, S50, S60, S70, S80, S90 (Patients with these isolated injuries that were transferred in or died THAT WERE EVALUATED BY TRAUMA SERVICE) –Centers only capture patients transferred for trauma care.

#### **PATIENT EXCLUSION CRITERIA**

Patients with the following isolated ICD-10-CM codes are **EXCLUDED** from the TACR:

- **NON-TRAUMA CENTERS:** S00, S10, S20, S30, S40, S50, S60, S70, S80, S90 (Patients with these isolated injuries that were transferred in/ out (**NOT** FOR TRAUMA EVALUATION) or died) –Centers only capture patients transferred for trauma care. This would exclude patients being transferred to another non-trauma center/ acute care facility for medical purposes.
- **TRAUMA CENTERS:** S00, S10, S20, S30, S40, S50, S60, S70, S80, S90 (Patients with these isolated injuries that were transferred in or died THAT WERE **NOT** EVALUATED BY TRAUMA SERVICE) –Centers only capture patients transferred for trauma care.
- **7<sup>th</sup> character modifiers of D through S** (Late effects)

#### **THE PATIENT MUST ALSO IN ADDITION TO THE ABOVE INCLUSION CRITERIA**

- Death resulting from the traumatic injury (independent of hospital admission or hospital transfer status);  
OR
- Patient transfer from one acute care hospital\*\* to another acute care hospital; OR
- Patients transferred/discharged to hospice (e.g., hospice facility, hospice unit, home hospice)  
OR
- Patients directly admitted to your hospital (exclude patients with isolated injuries admitted for elective and/or planned surgical intervention);  
OR
- Patients who were an in-patient admission and/or observed.

\*In-house traumatic injuries sustained after initial ED/Hospital arrival and before hospital discharge at the index hospital (the hospital reporting data), and all data associated with that injury event, are excluded.


\*\*Exclude patient injuries sustained at your facility after initial ED/Hospital arrival and before hospital discharge, and all data associated with that injury event.

\*\*Acute Care Hospital is defined as a hospital that provides inpatient medical care and other related services for surgery, acute medical conditions or injuries (usually for short-term illness or condition). “CMS Data Navigator Glossary of Terms”, [https://www.cms.gov/Research-Statistics-Data-and-Systems/Research/ResearchGenInfo/Downloads/DataNav\\_Glossary\\_Alpha.pdf](https://www.cms.gov/Research-Statistics-Data-and-Systems/Research/ResearchGenInfo/Downloads/DataNav_Glossary_Alpha.pdf).

\*\*\* Initial hospital encounter is referring to the patient’s first encounter for this injury event at your hospital.

**NOTE:** INCLUSION / EXCLUSION CRITERIA differs from NTDS. Please refer to the section “DIFFERENCES BETWEEN OHIO AND NATIONAL TRAUMA DATA STANDARD (NTDS)” starting on Page 12 for more information.





## COMMON NULL VALUES

---

### Definition

*Common Null Values* are terms to be used with OTR TACR Data Elements as described in this document for specifically-defined data fields when an answer cannot be provided.

### Element Values

NA= Not Applicable

ND= Not Known/Not Recorded/Not Documented

### Additional Information

- Although not written out on the following pages, these Common Null Values are included in the TACR dataset for every allowable data element. To ascertain their allowability by data field, see the “Accepts Null Value” notation on every data element descriptor page.
- *Not Applicable (Element Value NA)*: This null value code applies if, at any time of patient care documentation, the information requested was “Not Applicable” (NA) to the patient, the hospitalization or the patient care event. For example, variables documenting EMS care would be NA if a patient self-transportes to the hospital.
- *Not Known/Not Recorded/Not Documented (Element Value ND)*: This null value applies if, at the time of patient care documentation, information was “Not Known” (to the patient, family, healthcare provider) or no value for the element was recorded for the patient. This documents that there was an attempt to obtain information, but it was unknown by all parties or the information was missing at the time of documentation. For example, injury date and time may be documented in the hospital patient care report as “Unknown”. Another example, Not Known/Not Recorded/Not Documented should also be coded when documentation was expected, but none was provided (i.e., no EMS run sheet in the hospital record for patient transported by EMS).
- For any collection of data to be of value and reliably represent what was intended, a strong commitment must be made to ensure the correct documentation of incomplete data. When data elements associated with the TACR are to be electronically stored in a database or moved from one database to another, the indicated null values should be applied.

### References to Other Databases

- NTDS 2026

## HOSPITAL CODE

---

### Element Intent

This identifies the facility where the patient was seen, which is helpful for data linkage purposes and for further clarifying the resources that were available to treat the patient.

### Description

*Hospital Code* is a four-digit (4) hospital code assigned by the Ohio Department of Public Safety.

### Element Values

- Relevant value for data element

### Common Null Values

- Not Accepted

### Additional Information

- Stored as a four-digit code (xxxx)

### Data Source Hierarchy Guide

- 1 Ohio Department of Public Safety Hospital (Facility) Code List

### References to Other Databases

- Not an NTDS element

## UNIQUE ADMISSION NUMBER

---

### Element Intent

This identifies the specific trauma injury incident, which can help with record identification and data linkage purposes.

### Description

*Unique Admission Number* is a number assigned to the trauma patient at your facility. A patient encounter number or account number can be used.

### Element Values

- Relevant value for data element

### Common Null Values

- Not Accepted

### Additional Information

- Use an identifiable number specific to your facility, e.g. patient encounter or account number

### References to Other Databases

- Not an NTDS Element

## TRAUMA TRACKING NUMBER

---

### Element Intent

This identifies the specific trauma injury incident, which can help with record identification and data linkage purposes.

### Description

*Trauma Tracking Number* is a number automatically generated by the trauma registry system.

### Element Values

- Relevant value for data element

### Common Null Values

- Not Accepted

### References to Other Databases

- Not an NTDS Element

## FACILITY TYPE

---

### Element Intent

This is helpful for data linkage purposes and for further clarifying the resources that were available to treat the patient.

### Description

*Facility Type* is the type of facility at time of admission, transfer in or transfer out for each patient.

### Element Values

- 1 Free Standing Emergency Department
- 2 Acute Care Hospital
- 3 Adult Trauma 1
- 4 Adult Trauma 2
- 5 Adult Trauma 3
- 6 Pediatric Trauma 1
- 7 Pediatric Trauma 2

### Common Null Values

- Not Accepted

### References to Other Databases

- Not an NTDS Element

## PATIENT'S HOME CITY

---

### Element Intent

To conduct geospatial analyses or for linkage with geographic data sources.

### Description

*Patient's Home City* is the patient's city, township, or village of residence.

### Element Values

- Relevant value for data element

### Common Null Values

- Accepted

### Additional Information

- Used to calculate FIPS code
- The null value "Not Applicable" is reported for non-US hospitals.

### Data Source Hierarchy Guide

- 1 Face Sheet
- 2 Billing Sheet
- 3 Admission Form

### References to Other Databases

- NTDS 2026

**NOTE:** PATIENT HOME CITY differs from NTDS. Please refer to the section "DIFFERENCES BETWEEN OHIO AND NATIONAL TRAUMA DATA STANDARD (NTDS)" starting on Page 12 for more information.

## PATIENT'S HOME STATE

---

### Element Intent

To conduct geospatial analyses or for linkage with geographic data sources.

### Description

*Patient's Home State* is the state, territory, or province (or the District of Columbia) of the patient's residence.

### Element Values

- Relevant value for data element (two-digit FIPS code)

### Common Null Values

- Accepted

### Additional Information

- Used to calculate FIPS code
- The null value "Not Applicable" is reported for non-US hospitals.

### Data Source Hierarchy Guide

- 1 Face Sheet
- 2 Billing Sheet
- 3 Admission Form

### References to Other Databases

- NTDS 2026

**NOTE:** PATIENT HOME STATE differs from NTDS. Please refer to the section "DIFFERENCES BETWEEN OHIO AND NATIONAL TRAUMA DATA STANDARD (NTDS)" starting on Page 12 for more information.



## PATIENT'S HOME COUNTY

---

### Element Intent

To conduct geospatial analyses or for linkage with geographic data sources.

### Description

*Patient's Home County* is the patient's county (or parish) of residence.

### Element Values

- Relevant value for data element

### Common Null Values

- Accepted

### Additional Information

- Used to calculate FIPS code
- The null value "Not Applicable" is reported for non-US hospitals.

### Data Source Hierarchy Guide

- 1 Face Sheet
- 2 Billing Sheet
- 3 Admission Form

### References to Other Databases

- NTDS 2026

**NOTE:** PATIENT HOME COUNTY differs from NTDS. Please refer to the section "DIFFERENCES BETWEEN OHIO AND NATIONAL TRAUMA DATA STANDARD (NTDS)" starting on Page 12 for more information.

## PATIENT'S HOME ZIP CODE

---

### Element Intent

To conduct geospatial analyses or for linkage with geographic data sources.

### Description

*Patient's Home Zip Code* is the zip code of the patient's primary residence.

### Element Values

- Relevant value for data element

### Common Null Values

- Accepted

### Additional Information

- Can be stored as a 5 or 9-digit code (XXXXX-XXXX) for US and CA, or can be stored in the postal code format of the applicable country.
- If ZIP/Postal code is "Not Applicable," report variable: Alternate Home Residence.
- If ZIP/Postal code is "Not Known/Not Recorded," report variables: Patient's Home Country, Patient's Home State (US only), Patient's Home County (US only) and Patient's Home City (US only).
- If ZIP/Postal code is documented, must also report Patient's Home Country.

### Data Source Hierarchy Guide

- 1 Face Sheet
- 2 Billing Sheet
- 3 Admission Form

### References to Other Databases

- NTDS 2026

**NOTE:** PATIENT HOME ZIP CODE differs from NTDS. Please refer to the section "DIFFERENCES BETWEEN OHIO AND NATIONAL TRAUMA DATA STANDARD (NTDS)" starting on Page 12 for more information.

## PATIENT'S HOME COUNTRY

---

### Element Intent

To conduct geospatial analyses or for linkage with geographic data sources.

### Description

*Patient's Home Country* is the country where the patient resides.

### Element Values

- Relevant value for data element (two-digit alpha country code)

### Common Null Values

- Accepted

### Additional Information

- Values are two character FIPS codes representing the country (e.g. U.S.)
- If Patient's Home Country is not US, then the null value "Not Applicable" is reported for: Patient's Home State, Patient's Home County, and Patient's Home City.

### Data Source Hierarchy Guide

- 1 Face Sheet
- 2 Billing Sheet
- 3 Admission Form

### References to Other Databases

- NTDS 2026

## ALTERNATE HOME RESIDENCE

---

### Element Intent

To conduct geospatial analyses or for linkage with geographic data sources.

### Description

*Alternate Home Residence* is documentation of the residential status of a patient who has no home zip code.

### Element Values

- 1 Homeless
- 2 Undocumented Resident
- 3 Migrant Worker

### Common Null Values

- Accepted

### Additional Information

- Only used when Patient's Home ZIP/Postal Code is "Not Applicable"
- Report all that apply
- *Homeless* is defined as a person who lacks housing. The definition also includes a person living in transitional housing or a supervised public or private facility providing temporary living quarters
- *Undocumented* Citizen is defined as a national of another country who has entered or stayed in another country without permission
- *Migrant Worker* is defined as a person who temporarily leaves his/her principal place of residence within a country in order to accept seasonal employment in the same or different country.
- The null value "Not Applicable" is reported if Patient's Home ZIP/Postal Code is documented

### Data Source Hierarchy Guide

- 1 Face Sheet
- 2 Billing Sheet
- 3 Admission Form

### References to Other Databases

- NTDS 2026

## DATE OF BIRTH

---

### Element Intent

To calculate the patient's age at the time of the injury event, which is used for reporting and as a predictor of adverse outcomes.

### Description

*Date of Birth* is the patient's date of birth at time of injury.

### Element Values

- Relevant value for data element

### Common Null Values

- Accepted

### Additional Information

- Collected as YYYY-MM-DD
- If Date of Birth is "Not Known/Not Recorded," report variables: Age and Age Units.
- If Date of Birth equals Injury Date, then the Age and Age Units variables must be reported.

### Data Source Hierarchy Guide

- 1 Face Sheet
- 2 Billing Sheet
- 3 Admission Form
- 4 Triage / Trauma Flow Sheet
- 5 EMS Run Report

### References to Other Databases

- NTDS 2026

## AGE

---

### Element Intent

In the absence of the patient's date of birth, to calculate the patient's age at the time of the injury event, which is used for reporting and as a predictor of adverse outcomes.

### Description

*Age* is the patient's age (or best approximation) at the time of injury.

### Element Values

- Relevant value for data element

### Common Null Values

- Accepted

### Additional Information

- If Date of Birth is "Not Known/Not Recorded," report variables: Age and Age Units.
- If Date of Birth equals ED/Hospital Arrival Date, then the Age and Age Units variables must be reported.
- Must also report variable: Age Units.

### Data Source Hierarchy Guide

- 1 Face Sheet
- 2 Billing Sheet
- 3 Admission Form
- 4 Triage / Trauma Flow Sheet
- 5 EMS Run Report

### References to Other Databases

- NTDS 2026

**NOTE:** AGE differs from NTDS. Please refer to the section "DIFFERENCES BETWEEN OHIO AND NATIONAL TRAUMA DATA STANDARD (NTDS)" starting on Page 12 for more information.

## AGE UNITS

---

### Element Intent

In the absence of the patient's date of birth, to calculate the patient's age at the time of the injury event, which is used for reporting and as a predictor of adverse outcomes.

### Description

*Age Units* are the units used to document the patient's age (hours, days, months, years, minutes, weeks).

### Element Values

- 1 Hours
- 2 Days
- 3 Months
- 4 Years
- 5 Minutes
- 6 Weeks

### Common Null Values

- Accepted

### Additional Information

- If Date of Birth is "Not Known/Not Recorded," report variables: Age and Age Units.
- If Date of Birth equals ED/Hospital Arrival Date, then the Age and Age Units variables must be reported.
- Must also report variable: Age.

### Data Source Hierarchy Guide

- 1 Face Sheet
- 2 Billing Sheet
- 3 Admission Form
- 4 Triage / Trauma Flow Sheet
- 5 EMS Run Report

### References to Other Databases

- NTDS 2026

**NOTE:** AGE UNITS differs from NTDS. Please refer to the section "DIFFERENCES BETWEEN OHIO AND NATIONAL TRAUMA DATA STANDARD (NTDS)" starting on Page 12 for more information.

## SEX ASSIGNED AT BIRTH

---

### Element Intent

To analyze variations in injury patterns and outcomes.

### Description

The patient's sex assigned at birth

### Element Values

- 1 Male
- 2 Female
- 3 intersex

### Additional Information

- Also referred to as birth sex, natal sex, biological sex.

### Data Source Hierarchy Guide

- 1 Face Sheet
- 2 Billing Sheet
- 3 Admission Form
- 4 Triage/Trauma Flow Sheet
- 5 EMS Run report
- 6 History & Physical

### References to Other Databases

- NTDS 2026



## GENDER

---

### Element Intent

To analyze variations in injury patterns and outcomes.

### Description

The patient's gender identity

### Element Values

1. Man
2. Woman
3. Non-binary, genderqueer, gender nonconforming
4. Non-disclosed

### Additional Information

- Patient gender should be based upon self-report or identified by a family member.

### Data Source Hierarchy Guide

1. Face Sheet
2. Billing Sheet
3. Admission Form
4. Triage/Trauma Flow Sheet
5. EMS Run report
6. History & Physical

### References to Other Databases

- NTDS 2026

## GENDER-AFFIRMING HORMONE THERAPY

---

### Element Intent

To analyze variations in injury patterns and outcomes.

### Description

Is the patient currently (i.e., within the past 30 days) taking gender-affirming hormone therapy?

### Element Values

1. Yes
2. No
3. Non-disclosed

### Additional Information

- Gender-affirming hormone therapy includes but is not limited to estrogen, antiandrogens, and testosterone.
- If unclear if medication was for gender-affirming hormone therapy, then consult TMD or relevant physician/physician extender.

### Data Source Hierarchy Guide

1. Face Sheet
2. Billing Sheet
3. Admission Form
4. Triage/Trauma Flow Sheet
5. EMS Run report
6. History & Physical

### References to Other Databases

- NTDS 2026

## RACE

---

### Element Intent

To analyze variations in injury patterns and outcomes.

### Description

*Race* is the patient's race.

### Element Values

- 1 Asian
- 2 Native Hawaiian or Other Pacific Islander
- 3 Other Race
- 4 American Indian
- 5 Black or African American
- 6 White

### Common Null Values

- Accepted

### Additional Information

- Patient race should be based upon self-report or identified by a family member
- Based on the 2010 US Census Bureau
- Select all that apply

### Data Source Hierarchy Guide

- 1 Face Sheet
- 2 Billing Sheet
- 3 Admission Form
- 4 Triage/Trauma Flow Sheet
- 5 EMS Run report
- 6 History & Physical

### References to Other Databases

- NTDS 2026

## ETHNICITY

---

### Element Intent

To analyze variations in injury patterns and outcomes.

### Description

*Ethnicity* is the patient's ethnicity in terms of Hispanic heritage.

### Element Values

- 1 Hispanic or Latino
- 2 Not Hispanic or Latino

### Common Null Values

- Accepted

### Additional Information

- Patient ethnicity should be based upon self-report or identified by a family member
- The maximum number of ethnicities that may be reported for an individual patient is 1
- Based on the 2010 US Census Bureau

### Data Source Hierarchy Guide

- 1 Face Sheet
- 2 Billing Sheet
- 3 Admission Form
- 4 Triage/Trauma Flow Sheet
- 5 History & Physical
- 6 EMS Run Report

### References to Other Databases

- NTDS 2026

## PRIMARY ICD-10 EXTERNAL CAUSE CODE

---

### Element Intent

To identify potential injuries and are used as predictors of adverse outcomes.

### Description

*Primary External Cause Code* is a designation used to describe the mechanism (or external factor) that caused the injury event.

### Element Values

- Relevant ICD-10-CM code value for injury event

### Common Null Values

- Not Accepted

### Additional Information

- The Primary External Cause Code should describe the main reason a patient is admitted to the hospital.
- ICD-10-CM codes are accepted for this data element. Activity codes should not be reported for this data element.
- Activity codes should not be reported for this data element.
- Multiple Cause Coding Hierarchy: If two or more events cause separate injuries, an external cause code should be assigned for each cause. The first-listed external cause code will be selected in the following order:
  - External cause codes for child and adult abuse take priority over all other external cause codes.
  - External cause codes for terrorism events take priority over all other external cause codes except child and adult abuse.
  - External cause codes for cataclysmic events take priority over all other external cause codes except child and adult abuse, and terrorism.
  - External cause codes for transport accidents take priority over all other external cause codes except cataclysmic events, and child and adult abuse, and terrorism.
  - The first listed external cause code should correspond to the cause of the most serious diagnosis due to an assault, accident or self-harm, following the order of hierarchy listed above.

### Data Source Hierarchy Guide

- 1 EMS Run Sheet
- 2 Triage Form/Trauma Flow Sheet
- 3 Nursing Notes/Flow Sheet
- 4 History & Physical
- 5 Progress Notes

### References to Other Databases

- NTDS 2026

## ADDITIONAL ICD-10 EXTERNAL CAUSE CODE

---

### Element Intent

To identify potential injuries and are used as predictors of adverse outcomes.

### Description

*Additional External Cause Code* is used in conjunction with the Primary External Cause Code if multiple external cause codes are required to describe the injury event.

### Element Values

- Relevant ICD-10-CM code value for injury event

### Common Null Values

- Accepted

### Additional Information

- The null value “Not Applicable” is used if no additional external cause codes are used
- Activity codes should not be reported for this data element
- Report all that apply (maximum 2)
- Multiple Cause Coding Hierarchy: If two or more events cause separate injuries, an external cause code should be assigned for each cause. The first-listed external code will be selected in the following order:
  - External cause codes for child and adult abuse take priority over all other external cause codes
  - External cause codes for terrorism events take priority over all other external cause codes except child and adult abuse.
  - External cause codes for cataclysmic events take priority over all other external cause codes except child and adult abuse, and terrorism.
  - External cause codes for transport accident take priority over all other external cause codes except cataclysmic events, and child and adult abuse, and terrorism.
  - The first listed external cause code should correspond to the cause of the most serious diagnosis due to an assault, accident or self-harm, following the order of hierarchy listed above.

### Data Source Hierarchy Guide

- 1 EMS Run Sheet
- 2 Triage Form/Trauma Flow Sheet
- 3 Nursing Notes/ Flow Sheet
- 4 History & Physical
- 5 Progress Notes

### References to Other Databases

- NTDS 2026

## ICD-10 PLACE OF OCCURRENCE EXTERNAL CAUSE CODE

---

### Element Intent

To provide geographic context to the injury and describes the nature, activity, and cause.

### Description

*ICD-10 Place of Occurrence external cause code* is a Y92.x code used to describe the place, site or location of the injury event.

### Element Values

- Relevant ICD-10-CM or ICD-10-CA code value for injury event

### Common Null Values

- Not Accepted

### Additional Information

- Only ICD-10-CM or ICD-10-CA codes will be accepted for ICD-10 Place of Occurrence External Cause Code.

### Data Source Hierarchy Guide

- 1 EMS Run Sheet
- 2 Triage Form/Trauma Flow Sheet
- 3 Nursing Notes/ Flow Sheet
- 4 History & Physical
- 5 Progress Notes

### References to Other Databases

- NTDS 2026

## WORK-RELATED

---

### Element Intent

To analyze variations in injury patterns and outcomes.

### Description

*Work-related* is whether the injury occurred during paid employment.

### Element Values

- 1 Yes
- 2 No

### Common Null Values

- Accepted

### Additional Information

- If work-related, two additional data elements must be completed, *Patient's Occupational Industry* and *Patient's Occupation*

### Data Source Hierarchy Guide

- 1 EMS Run Report
- 2 Triage/Trauma Flow Sheet
- 3 History & Physical
- 4 Face Sheet
- 5 Billing Sheet

### References to Other Databases

- NTDS 2026



## PATIENT'S OCCUPATIONAL INDUSTRY

---

### Element Intent

To analyze variations in injury patterns and outcomes.

### Description

*Patient's Occupational Industry* is the occupational industry associated with the patient's work environment.

### Element Values

- |                                    |                              |
|------------------------------------|------------------------------|
| 1 Finance, Insurance, Real Estate  | 8 Construction               |
| 2 Manufacturing                    | 9 Government                 |
| 3 Retail Trade                     | 10 Natural Resources, Mining |
| 4 Transportation, Public Utilities | 11 Information Services      |
| 5 Agriculture, Forestry, Fishing   | 12 Wholesale Trade           |
| 6 Professional, Business Services  | 13 Leisure, Hospitality      |
| 7 Education, Health Services       | 14 Other Services            |

### Common Null Values

- Accepted

### Additional Information

- If work related, also report *Patient's Occupation*
- Based upon US Bureau of Labor Statistics Industry Classification
- Code as *NA* if injury is not work-related AND Work-Related value is coded is given a value of "2. No".

### Data Source Hierarchy Guide

- 1 Billing Sheet
- 2 Face Sheet
- 3 Case Management/Social Services Notes
- 4 EMS Run Report
- 5 Nursing Notes/Flow Sheet

### References to Other Databases

- NTDS 2026

## PATIENT'S OCCUPATION

---

### Element Intent

To analyze variations in injury patterns and outcomes.

### Description

*Patient's Occupation* is the occupation of the patient.

### Element Values

- |   |   |
|---|---|
| 1 Business, Financial Operations Occupations      | 13 Computer, Mathematical Occupations         |
| 2 Architecture, Engineering Occupations           | 14 Life, Physical, Social Science Occupations |
| 3 Community, Social Services Occupations          | 15 Legal Occupations                          |
| 4 Education, Training, Library Occupations        | 16 Arts, Design, Entertainment, Sports, Media |
| 5 Healthcare Practitioners, Technical Occupations | 17 Healthcare Support Occupations             |
| 6 Protective Service Occupations                  | 18 Food Preparation, Serving Related          |
| 7 Building, Grounds Cleaning & Maintenance        | 19 Personal Care, Service Occupations         |
| 8 Sales & Related Occupations                     | 20 Office, Administrative Support Occupations |
| 9 Farming, Fishing, Forestry Occupations          | 21 Construction, Extraction Occupations       |
| 10 Installation, Maintenance, Repair Occupations  | 22 Production Occupations                     |
| 11 Transportation, Material Moving Occupations    | 23 Military Specific Occupations              |
| 12 Management Occupations                         |   |

### Common Null Values

- Accepted

### Additional Information

- Only report if injury is work related.
- If work related, also report *Patient's Occupational Industry*.
- Based upon 1999 US Bureau of Labor Statistics Standard Occupational Classification (SOC).
- Code as *NA* if injury is not work-related AND Work-Related value is coded is given a value of "2. No".

### Data Source Hierarchy Guide

- 1 Billing Sheet
- 2 Face Sheet
- 3 Case Management/Social Services Notes
- 4 EMS Run Report
- 5 Nursing Notes/Flow Sheet

### References to Other Databases

- NTDS 2026

## INJURY INCIDENT DATE

---

### Element Intent

To analyze the timeline of the care event and the timeliness of interventions.

### Description

*Injury Incident Date* is the date that the injury occurred.

### Element Values

- Relevant value for data element

### Common Null Values

- Accepted

### Additional Information

- Collected as YYYY-MM-DD
- Estimates of the date of injury should be based upon report by patient, witness, family or health care provider (onset date). Other proxy measures (e.g. 911 call-time) should NOT be used.

### Data Source Hierarchy Guide

- 1 EMS Run report
- 2 Triage/Trauma Flow Sheet
- 3 History & Physical
- 4 Face Sheet

### References to Other Databases

- NTDS 2026

## INJURY INCIDENT TIME

---

### Element Intent

To analyze the timeline of the care event and the timeliness of interventions.

### Description

*Injury Incident Time* is the time of day that the injury occurred.

### Element Values

- Relevant value for data element

### Common Null Values

- Accepted

### Additional Information

- Collected as HHMM military time
- Estimates of time of injury should be based upon report by patient, witness, family, or health care provider (onset time). Other proxy measures (e.g. 911 call-time) should NOT be used.

### Data Source Hierarchy Guide

- 1 EMS Run report
- 2 Triage/Trauma Flow Sheet
- 3 History & Physical
- 4 Face Sheet

### References to Other Databases

- NTDS 2026

## INCIDENT CITY

---

### Element Intent

To conduct geospatial analyses or for linkage with geographic data sources.

### Description

*Incident City* is the city or township where the incident occurred.

### Element Values

- Relevant value for data element (five-digit FIPS code)

### Common Null Values

- Accepted

### Additional Information

- Used to calculate FIPS code
- If incident location resides outside of formal city boundaries, report nearest city/town.
- If Incident Country is not US, report the null value "Not Applicable."

### Data Source Hierarchy Guide

- 1 EMS Run Report
- 2 Triage/Trauma Flow Sheet

### References to Other Databases

- NTDS 2026

**NOTE:** INCIDENT CITY differs from NTDS. Please refer to the section "DIFFERENCES BETWEEN OHIO AND NATIONAL TRAUMA DATA STANDARD (NTDS)" starting on Page 12 for more information.

## INCIDENT STATE

---

### Element Intent

To conduct geospatial analyses or for linkage with geographic data sources.

### Description

*Incident State* is the state, territory or province where the incident occurred.

### Element Values

- Relevant value for data element (two-digit numeric FIPS code)

### Common Null Values

- Accepted

### Additional Information

- Used to calculate FIPS code
- If Incident Country is not US, report the null value "Not Applicable."

### Data Source Hierarchy Guide

- 1 EMS Run Report
- 2 Triage/Trauma Flow Sheet

### References to Other Databases

- NTDS 2026

**NOTE:** INCIDENT STATE differs from NTDS. Please refer to the section “DIFFERENCES BETWEEN OHIO AND NATIONAL TRAUMA DATA STANDARD (NTDS)” starting on Page 12 for more information.

## INCIDENT COUNTY

---

### Element Intent

To conduct geospatial analyses or for linkage with geographic data sources.

### Description

*Incident County* is the county or parish where the incident occurred.

### Element Values

- Relevant value for data element (three-digit FIPS code)

### Common Null Values

- Accepted

### Additional Information

- Used to calculate FIPS code
- If Incident Country is not US, report the null value "Not Applicable."

### Data Source Hierarchy Guide

- 1 EMS Run Report
- 2 Triage/Trauma Flow Sheet

### References to Other Databases

- NTDS 2026

**NOTE:** INCIDENT COUNTY differs from NTDS. Please refer to the section “DIFFERENCES BETWEEN OHIO AND NATIONAL TRAUMA DATA STANDARD (NTDS)” starting on Page 12 for more information.

## INCIDENT LOCATION ZIP CODE

---

### Element Intent

To conduct geospatial analyses or for linkage with geographic data sources.

### Description

*Incident Location Zip Code* is the zip code of the location where the patient was injured.

### Element Values

- Relevant value for data element

### Common Null Values

- Accepted

### Additional Information

- Stored as a five-digit code (XXXXX).
- If "Not Known/Not Recorded," report variables: Incident Country, Incident State (US Only), Incident County (US Only) and Incident City (US Only).
- May require adherence to HIPAA regulations.
- If ZIP/Postal code is documented, then must report Incident Country.

### Data Source Hierarchy Guide

- 1 EMS Run Report
- 2 Triage/Trauma Flow Sheet

### References to Other Databases

- NTDS 2026

**NOTE:** INCIDENT LOCATION ZIP CODE differs from NTDS. Please refer to the section “DIFFERENCES BETWEEN OHIO AND NATIONAL TRAUMA DATA STANDARD (NTDS)” starting on Page 12 for more information.



## INCIDENT COUNTRY

---

### Element Intent

To conduct geospatial analyses or for linkage with geographic data sources.

### Description

*Incident Country* is the country where the incident occurred.

### Element Values

- Relevant value for data element (two-digit alpha country code)

### Common Null Values

- Accepted

### Additional Information

- Values are two character FIPS codes representing a country (e.g. US)
- If Incident Country is not US, then the null value "Not Applicable" is reported for: Incident State, Incident County, and Incident Home City

### Data Source Hierarchy Guide

- 1 EMS Run report
- 2 Triage/Trauma Flow Sheet

### References to Other Databases

- NTDS 2026

## PROTECTIVE DEVICES

---

### Element Intent

To analyze the prevalence and effects of safety equipment.

### Description

*Protective Devices* is the safety equipment in use or worn by the patient at the time of the injury.

### Element Values

- |  |   |
|--|---|
| 1 None Used  | 7 Helmet (e.g., bicycle, skiing, motorcycle)      |
| 2 Lap Belt   | 8 Airbag Present                                  |
| 3 Personal Floatation Device   | 9 Protective Clothing (e.g. padded leather pants) |
| 4 Protective Non-Clothing Gear (e.g. shin guard)                           | 10 Shoulder Belt                                  |
| 5 Eye Protection   | 11 Other  |
| 6 Child Restraint (child car seat, infant car seat, or child booster seat) |   |

### Common Null Values

- Accepted
- Element cannot be “Not Applicable”

### Additional Information

- Report all that apply
- If “Child Restraint” is present, report variable *Child Specific Restraint*
- If “Airbag” is present, report variable *Airbag Deployment*
- Evidence of the use of safety equipment may be reported or observed
- “Lap belt” should be reported to include those patients that are restrained, but not further specified
- If chart indicates “3-point-restraint,” report element value “2. Lap Belt” and “10. Shoulder Belt.”
- If documented that a “Child Restraint (booster seat or child care seat)” was used or worn, but not properly fastened, either on the child or in the car, report Element Value “1. None.”

### Data Source Hierarchy Guide

- 1 EMS Run Sheet
- 2 Triage/Trauma Flow Sheet
- 3 Nursing Notes / Flow Sheet
- 4 History & Physical

### References to Other Databases

- NTDS 2026

## CHILD SPECIFIC RESTRAINT

---

### Element Intent

To analyze the prevalence and effects of safety equipment.

### Description

*Child Specific Restraint* indicates protective child restraint devices used by the pediatric patient at the time of injury.

### Element Values

- 1 Child Car Seat
- 2 Infant Car Seat
- 3 Child Booster Seat

### Common Null Values

- Accepted

### Additional Information

- Evidence of the use of child restraint may be reported or observed
- Only reported when *Protective Devices* include "6. Child Restraint (booster seat or child car seat)."
- The null value "Not Applicable" is reported if Element Value 6. "Child Restraint" is NOT reported for Protective Devices.
- Report Element Value "1. Child Car Seat" for forward-facing child seats.
- Report Element Value "2. Infant Car Seat" for rear-facing child seats.

### Data Source Hierarchy Guide

- 1 EMS Run Sheet
- 2 Triage/Trauma Flow Sheet
- 3 Nursing Notes / Flow Sheet
- 4 History & Physical

### References to Other Databases

- NTDS 2026

## AIRBAG DEPLOYMENT

---

### Element Intent

To analyze the prevalence and effects of safety equipment.

### Description

*Airbag Deployment* indicates whether an airbag deployed during a motor vehicle crash.

### Element Values

- 1 Airbag Not Deployed
- 2 Airbag Deployed Front
- 3 Airbag Deployed Side
- 4 Airbag Deployed Other (knee, airbelt, curtain, etc.)

### Common Null Values

- Accepted

### Additional Information

- Report all that apply.
- Evidence of the use of airbag deployment may be reported or observed.
- Only report when Protective Devices include "8. Airbag Present."
- Airbag Deployed Front should be reported for patients with documented airbag deployments, but are not further specified.
- The null value "Not Applicable" is reported if Element Value 8. "Airbag Present" is NOT reported for Protective Devices.

### Data Source Hierarchy Guide

- 1 EMS Run Sheet
- 2 Triage/Trauma Flow Sheet
- 3 Nursing Notes / Flow Sheet
- 4 History & Physical

### References to Other Databases

- NTDS 2026

## TRANSPORT MODE FOR ARRIVAL AT YOUR HOSPITAL

---

### Element Intent

To analyze patterns between different transportation modes.

### Description

*Transport Mode for Arrival at Your Hospital* is the manner of transport delivering the patient to your hospital.

### Element Values

- 1 Ground Ambulance
- 2 Helicopter Ambulance
- 3 Fixed-wing Ambulance
- 4 Private or Public Vehicle or Walk-in
- 5 Police Transport
- 6 Other Transport Mode

### Common Null Values

- Accepted

### Additional Information

- Example of “Other Transport Mode” include boat
- Examples of “Public or Private or Walk-in” include: bus, bicycle or personal vehicle
- If a patient was a visitor/in-house patient at your facility and experienced an event to require admission to the ED select patient’s mode of arrival as “4/Private or Public Vehicle or Walk-In”.

### Data Source Hierarchy Guide

- 1 EMS Run Report

### References to Other Databases

- NTDS 2026

**NOTE:** TRANSPORT MODE FOR ARRIVAL AT YOUR HOSPITAL differs from NTDS. Please refer to the section “DIFFERENCES BETWEEN OHIO AND NATIONAL TRAUMA DATA STANDARD (NTDS)” starting on Page 12 for more information.

## TRANSPORT AGENCY

---

### Element Intent

This identifies the EMS agency that treated the patient prior to arrival at the destination facility and provides vital information regarding the patient's prehospital care.

### Description

*Transport Agency* is the EMS agency or air ambulance that delivered the patient to your hospital.

### Element Values

- Relevant value for data element (ODPS-assigned EMS Agency ID)

### Common Null Values

- Accepted

### Additional Information

- "Non-applicable" (NA) is used to indicate that a patient arrived via "Private or Public Vehicle or Walk-in," "Police Transport," or "Other Transport Mode"

### Data Source Hierarchy Guide

- 1 EMS Run Report
- 2 ED Record

### References to Other Databases

- Not an NTDS element

## OTHER TRANSPORT MODES

---

### Element Intent

To analyze patterns between different transportation modes.

### Description

*Other Transport Modes* documents all other types of transport used during patient care prior to the patient arriving at your hospital, except the transport mode delivering the patient to your hospital.

### Element Values

- 1 Ground Ambulance
- 2 Helicopter Ambulance
- 3 Fixed-wing Ambulance
- 4 Private or Public Vehicle or Walk-in
- 5 Police Transport
- 6 Other Transport Mode

### Common Null Values

- Accepted

### Additional Information

- For patients with an unspecified mode of transport, select 6, *Other*
- The null value “Not Applicable” is reported to indicate that a patient had a single mode of transport.
- Report all that apply with a maximum of 5.
- An example is an ambulance transporting the patient to the helicopter landing zone.

### Data Source Hierarchy Guide

1. EMS Run Report
2. Transfer Facility Records

### References to Other Databases

- NTDS 2026

**NOTE:** OTHER TRANSPORT MODES differs from NTDS. Please refer to the section “DIFFERENCES BETWEEN OHIO AND NATIONAL TRAUMA DATA STANDARD (NTDS)” starting on Page 12 for more information.

## EMS PATIENT CARE REPORT UNIQUE IDENTIFIER (UUID)

---

### Element Intent

To link hospital and EMS data systems.

### Description

The universally unique identifier (UUID) of the patient care report (PCR) of each emergency service (EMS) unit treating the patient from the time of injury to arrival at your ED/hospital.

### Element Values

- Relevant value for data element
- Must be represented in canonical form, matching the following regular expression:  
[a-fA-F0-9]{8}-[a-fA-F0-9]{4}-[1-5][a-fA-F0-9]{3}-[89abAB][a-fA-F0-9]{3}-[afA-F0-9]{12}

### Additional Information

- Report all that apply (maximum 20).
- A sample UUID is: e48cd734-01cc-4da4-ae6a-915b0b1290f6
- Automated abstraction technology provided by registry product providers/vendors must be used for this data element. In the absence of automated technology, report the null value "Not Known/Not Recorded."
- Consistent with NEMSIS v3.5.0.
- The null value "Not Known/Not Recorded" must be reported if the UUID is not documented on the EMS Run Report. The UUID will not be documented on EMS Run Reports in NEMSIS versions lower than 3.5.0. In collaboration with NEMSIS, the ACS will communicate when NEMSIS 3.5.0 is widely implemented.
- The null value "Not Applicable" must be reported if the patient was never transported via EMS prior to arrival at your hospital.
- Assigned by any applicable transporting EMS agency in accordance with the IETF RFC 4122 standard.

### Data Source Hierarchy Guide

- 1 EMS Run Report

### References to Other Databases

- NTDS 2026



## EMS DISPATCH DATE TO SCENE OR TRANSFERRING FACILITY

---

### Element Intent

This provides information regarding the patient's prehospital care in relation to how quickly the EMS agency was able to arrive at the patient, spend time at the scene, and transport the patient to a destination facility.

### Description

The date the unit *transporting to your hospital* was notified by dispatch.

### Element Values

- Relevant value for data element

### Common Null Values

- Accepted

### Additional Information

- Collected as YYYY-MM-DD
- For inter-facility transfer patients, this is the date on which the unit transporting the patient to your facility from the transferring facility was notified by dispatch or assigned to this transport.
- For patients transported from the scene of injury to your hospital, this is the date on which the unit transporting the patient to your facility from the scene was dispatched.
- The null value "Not Applicable" is reported for patients who were not transported by EMS

### Data Source Hierarchy Guide

- 1 EMS Run Report

### References to Other Databases

- Not an NTDS element

## EMS DISPATCH TIME TO SCENE OR TRANSFERRING FACILITY

---

### Element Intent

This provides information regarding the patient's prehospital care in relation to how quickly the EMS agency was able to arrive at the patient, spend time at the scene, and transport the patient to a destination facility.

### Description

The time the unit *transporting to your hospital* was notified by dispatch.

### Element Values

- Relevant value for data element

### Common Null Values

- Accepted

### Additional Information

- Collected as HHMM military time
- For inter-facility transfer patients, this is the time at which the unit transporting the patient to your facility from the transferring facility was notified by dispatch.
- For patients transported from the scene of injury to your hospital, this is the time at which the unit transporting the patient to your facility from the scene was dispatched.
- The null value "Not Applicable" is used for patients who were not transported by EMS

### Data Source Hierarchy Guide

- 1 EMS Run Report

### References to Other Databases

- Not an NTDS element

## EMS UNIT ARRIVAL DATE AT SCENE OR TRANSFERRING FACILITY

---

### Element Intent

This provides information regarding the patient's prehospital care in relation to how quickly the EMS agency was able to arrive at the patient, spend time at the scene, and transport the patient to a destination facility.

### Description

The date the unit *transporting to your hospital* arrived on the scene/transferring facility (the time the vehicle stopped moving).

### Element Values

- Relevant value for data element

### Common Null Values

- Accepted

### Additional Information

- Collected as YYYY-MM-DD
- For inter-facility transfer patients, this is the date on which the unit transporting the patient to your facility from the transferring facility arrived at the transferring facility (arrival is defined at date/time when the vehicle stopped moving).
- For patients transported from the scene of injury to your hospital, this is the date on which the unit transporting the patient to your facility from the scene arrived at the scene (arrival is defined at date/time when the vehicle stopped moving).
- The null value "Not Applicable" is used for patients who were not transported by EMS

### Data Source Hierarchy Guide

- 1 EMS Run Report

### References to Other Databases

- Not an NTDS element

## EMS UNIT ARRIVAL TIME AT SCENE OR TRANSFERRING FACILITY

---

### Element Intent

This provides information regarding the patient's prehospital care in relation to how quickly the EMS agency was able to arrive at the patient, spend time at the scene, and transport the patient to a destination facility.

### Description

The time the unit *transporting to your hospital* arrived on the scene (the time the vehicle stopped moving).

### Element Values

- Relevant value for data element

### Common Null Values

- Accepted

### Additional Information

- Collected as HHMM military time
- For inter-facility transfer patients, this is the time at which the unit transporting the patient to your facility from the transferring facility arrived at the transferring facility (arrival is defined at date/time when the vehicle stopped moving).
- For patients transported from the scene of injury to your hospital, this is the time at which the unit transporting the patient to your facility from the scene arrived at the scene (arrival is defined at date/time when the vehicle stopped moving).
- The null value "Not Applicable" is used for patients who were not transported by EMS

### Data Source Hierarchy Guide

- 1 EMS Run Report

### References to Other Databases

- Not an NTDS element

## EMS UNIT DEPARTURE DATE FROM SCENE OR TRANSFERRING FACILITY

---

### Element Intent

This provides information regarding the patient's prehospital care in relation to how quickly the EMS agency was able to arrive at the patient, spend time at the scene, and transport the patient to a destination facility.

### Description

The date the unit *transporting to your hospital* left the scene (the time the vehicle started moving).

### Element Values

- Relevant value for data element

### Common Null Values

- Accepted

### Additional Information

- Collected as YYYY-MM-DD
- For inter-facility transfer patients, this is the date on which the unit transporting the patient to your facility from the transferring facility departed from the transferring facility (departure is defined at date/time when the vehicle started moving).
- For patients transported from the scene of injury to your hospital, this is the date on which the unit transporting the patient to your facility from the scene departed from the scene (arrival is defined at date/time when the vehicle started moving).
- The null value "Not Applicable" is used for patients who were not transported by EMS

### Data Source Hierarchy Guide

- 1 EMS Run Report

### References to Other Databases

- Not an NTDS element

## EMS UNIT DEPARTURE TIME FROM SCENE OR TRANSFERRING FACILITY

---

### Element Intent

This provides information regarding the patient's prehospital care in relation to how quickly the EMS agency was able to arrive at the patient, spend time at the scene, and transport the patient to a destination facility.

### Description

The time the unit *transporting to your hospital* left the scene (the time the vehicle started moving).

### Element Values

- Relevant value for data element

### Common Null Values

- Accepted

### Additional Information

- Collected as HHMM military time
- For inter-facility transfer patients, this is the time at which the unit transporting the patient to your facility from the transferring facility departed from the transferring facility (departure is defined at date/time when the vehicle started moving).
- For patients transported from the scene of injury to your hospital, this is the time at which the unit transporting the patient to your facility from the scene departed from the scene (arrival is defined at date/time when the vehicle started moving).
- The null value "Not Applicable" is used for patients who were not transported by EMS

### Data Source Hierarchy Guide

- 1 EMS Run Report

### References to Other Databases

- Not an NTDS element

## INITIAL FIELD SYSTOLIC BLOOD PRESSURE

---

### Element Intent

This provides information regarding the patient's prehospital care and health status.

### Description

Initial Field *Systolic Blood Pressure* is the first recorded systolic blood pressure measured.

### Element Values

- Relevant value for data element

### Common Null Values

- Accepted

### Additional Information

- If patient is transferred to your facility with no EMS run sheet from the scene of injury, record as *Not Known/Not Recorded/Not Documented*
- Measurement recorded must be without the assistance of CPR or any type of mechanical chest compression device. For those patients who are receiving CPR or any type of mechanical chest compressions, report the value obtained while compressions are paused.
- The null value "Not Applicable" is reported for patients who arrive by "4. Private/Public Vehicle/Walk-in."
- The null value "Not Known/Not Recorded" is reported if the patient's first recorded initial field systolic blood pressure was NOT measured

### Data Source Hierarchy Guide

- 1 EMS Run Report

### References to Other Databases

- Not an NTDS element

## INITIAL FIELD PULSE RATE

---

### Element Intent

This provides information regarding the patient's prehospital care and health status.

### Description

*Initial Field Pulse Rate* is the first recorded pulse measured (palpated or auscultated), expressed as a number per minute.

### Element Values

- Relevant value for data element

### Common Null Values

- Accepted

### Additional Information

- If patient is transferred to your facility with no EMS run sheet from the scene of injury, record as *Not Known/Not Recorded/Not Documented*
- Measurement recorded must be without the assistance of CPR or any type of mechanical chest compression device. For those patients who are receiving CPR or any type of mechanical chest compressions, report the value obtained while compressions are paused.
- The null value "Not Applicable" is reported for patients who arrive by "4. Private/Public Vehicle/Walk-in."
- The null value "Not Known/Not Recorded" is reported if the patient's first recorded initial field pulse rate was NOT measured

### Data Source Hierarchy Guide

- 1 EMS Run Report

### References to Other Databases

- Not an NTDS element



## INITIAL FIELD RESPIRATORY RATE

---

### Element Intent

This provides information regarding the patient's prehospital care and health status.

### Description

*Initial Field Respiratory Rate* is the first recorded respiratory rate measured (expressed as a number per minute).

### Element Values

- Relevant value for data element

### Common Null Values

- Accepted

### Additional Information

- If patient is transferred to your facility with no EMS run sheet from the scene of injury, record as *Not Known/Not Recorded/Not Documented*
- The null value "Not Applicable" is reported for patients who arrive by "4. Private/Public Vehicle/Walk-in."
- The null value "Not Known/Not Recorded" is reported if the patient's first recorded initial field respiratory rate was NOT measured

### Data Source Hierarchy Guide

- 1 EMS Run Report

### References to Other Databases

- Not an NTDS element

## INITIAL FIELD OXYGEN SATURATION

---

### Element Intent

This provides information regarding the patient's prehospital care and health status.

### Description

*Initial Field Oxygen Saturation* is the first recorded oxygen saturation measured (expressed as a percentage).

### Element Values

- Relevant value for data element

### Common Null Values

- Accepted

### Additional Information

- If patient is transferred to your facility with no EMS run sheet from the scene of injury, record as *Not Known/Not Recorded/Not Documented*
- The null value "Not Applicable" is reported for patients who arrive by 4. Private/Public Vehicle/Walk-in.
- The null value "Not Known/Not Recorded" is reported if the patient's first recorded initial field oxygen saturation was NOT measured

### Data Source Hierarchy Guide

- 1 EMS Run Report

### References to Other Databases

- Not an NTDS element

## INITIAL FIELD GCS - EYE

---

### Element Intent

This provides information regarding the patient's prehospital care and health status.

### Description

*Initial Field GCS Eye Opening* is the first recorded Glasgow Coma Score eye assessment done.

### Element Values

- 1 No eye movement when assessed
- 2 Opens eyes in response to painful stimulation
- 3 Opens eyes in response to verbal stimulation
- 4 Opens eyes spontaneously

### Common Null Values

- Accepted

### Additional Information

- If patient is transferred to your facility with no EMS run sheet from the scene of injury, record as *Not Known/Not Recorded/Not Documented*
- If a patient does not have a numeric GCS score recorded, but written documentation closely (or directly) relates to verbiage describing a specific level of functioning within the GCS scale, the appropriate numeric score may be listed. E.g. the chart indicates: "patient's pupils are PERRL," an Eye GCS of 4 may be recorded, IF there is no other contradicting documentation
- The null value "Not Applicable" is used for patients who arrive by 4. Private/Public Vehicle/ Walk-in
- The null value "Not Known/Not Recorded" is reported if the patient's first recorded initial field GCS-Eye was NOT measured

### Data Source Hierarchy Guide

- 1 EMS Run Report

### References to Other Databases

- Not an NTDS element

## INITIAL FIELD GCS - VERBAL

---

### Element Intent

This provides information regarding the patient's prehospital care and health status.

### Description

*Initial Field GCS Verbal Response* is the first recorded Glasgow Coma Score verbal assessment done.

### Element Values

- Pediatric(<= 2 years of age)
  - 1 No vocal response
  - 2 Inconsolable, agitated
  - 3 Inconsistently consolable, moaning
  - 4 Cries but is consolable, inappropriate interactions
  - 5 Smiles, oriented to sounds, follows objects, interacts
- Adult
  - 1 No verbal response
  - 2 Incomprehensible sounds
  - 3 Inappropriate words
  - 4 Confused
  - 5 Oriented

### Common Null Values

- Accepted

### Additional Information

- If patient is transferred to your facility with no EMS run sheet from the scene of injury, record as *Not Known/Not Recorded/Not Documented*
- If patient is intubated, then the GCS Verbal score is equal to 1
- If a patient does not have a numeric GCS score recorded, but written documentation closely (or directly) relates to verbiage describing a specific level of functioning within the GCS scale, the appropriate numeric score may be listed. E.g. the chart indicates: "patient is oriented to person place and time," a Verbal GCS of 5 may be recorded, IF there is no other contradicting documentation
- The null value "Not Applicable" is used for patients who arrive by 4. Private/Public Vehicle/ Walk-in
- The null value "Not Known/Not Recorded" is reported if the patient's first recorded initial field GCS - Verbal was NOT measured

### Data Source Hierarchy Guide

- 1 EMS Run Report

### References to Other Databases

- Not an NTDS element

## INITIAL FIELD GCS - MOTOR

---

### Element Intent

This provides information regarding the patient's prehospital care and health status.

### Description

*Initial Field GCS Motor Response* is the first recorded Glasgow Coma Score motor assessment done.

### Element Values

- |   |   |
|---|---|
| <ul style="list-style-type: none"><li>• <u>Pediatric (&lt;= 2 years of age)</u><ul style="list-style-type: none"><li>1 No motor response</li><li>2 Extension to pain</li><li>3 Flexion to pain</li><li>4 Withdrawal from pain</li><li>5 Localizing pain</li><li>6 Appropriate response to stimulation</li></ul></li></ul> | <ul style="list-style-type: none"><li>• <u>Adult</u><ul style="list-style-type: none"><li>1 No motor response</li><li>2 Extension to pain</li><li>3 Flexion to pain</li><li>4 Withdrawal from pain</li><li>5 Localizing pain</li><li>6 Obeys commands</li></ul></li></ul> |
|---|---|

### Common Null Values

- Accepted

### Additional Information

- If patient is transferred to your facility with no EMS run sheet from the scene of injury, record as *Not Known/Not Recorded/Not Documented*
- If a patient does not have a numeric GCS score recorded, but written documentation closely (or directly) relates to verbiage describing a specific level of functioning within the GCS scale, the appropriate numeric score may be listed. E.g. the chart indicates: "patient withdraws from a painful stimulus," a Motor GCS of 4 may be recorded, IF there is no other contradicting documentation.
- The null value "Not Applicable" is reported for patients who arrive by "4. Private/Public Vehicle/Walk-in"
- The null value "Not Known/Not Recorded" is reported if the patient's first recorded initial field GCS - Motor was NOT measured

### Data Source Hierarchy Guide

- 1 EMS Run Report

### References to Other Databases

- Not an NTDS element

## INITIAL FIELD GCS - TOTAL

---

### Element Intent

This provides information regarding the patient's prehospital care and health status.

### Description

*Initial Field GCS Total* is the first recorded total Glasgow Coma Score done.

### Element Values

- Relevant value for data element

### Common Null Values

- Accepted

### Additional Information

- If the patient is transferred to your facility with no EMS run sheet from the scene of injury, record as *Not Known/Not Recorded/Not Documented*
- If a patient does not have a numeric GCS recorded, but there is documentation related to their level of consciousness such as "AAOx3," "awake alert and oriented," or "patient with normal mental status," interpret this as GCS of 15 IF there is no other contradicting documentation
- The null value "Not Applicable" is reported for patients who arrive by "4. Private/Public Vehicle/Walk-in"
- The null value "Not Known/Not Recorded" is reported if the patient's first recorded initial field GCS - Total was NOT measured

### Data Source Hierarchy Guide

- 1 EMS Run Report

### References to Other Databases

- Not an NTDS element

## INITIAL FIELD GCS QUALIFIER

---

### Element Intent

This provides information regarding the patient's prehospital care and health status.

### Description

*Initial Field GCS Qualifier* documents circumstances related to the patient when or near the time that the *Initial Field GCS Total* was obtained.

### Element Values

- 1 Patient is chemically sedated or paralyzed
- 2 Obstruction to the patient's eye(s) prevents accurate eye assessment
- 3 Patient is intubated
- 4 GCS is valid meaning that the patient is not sedated, not intubated and without eye obstruction

### Common Null Values

- Accepted

### Additional Information

- Identifies treatments given to the patient that may affect the first assessment of GCS. This element does not apply to self-medications the patient may administer (i.e., ETOH, prescriptions, etc.)
- Element Value "1. Patient Chemically Sedated or Paralyzed" is reported if an intubated patient has recently received an agent that results in neuromuscular blockade such that a motor or eye response is not possible.
- Neuromuscular blockade is typically induced following the administration of agent like succinylcholine, mivacurium, rocuronium, (cis)atracurium, vecuronium, or pancuronium. While these are the most common agents, please review what might be typically used in your center so it can be identified in the medical record.
- Each of these agents has a slightly different duration of action, so their effect on the GCS depends on when they were given. For example, succinylcholine's effects last for only 5-10 minutes.
- Please note that the first recorded vitals do not need to be from the same assessment.
- Select NA if the patient was not transported to your hospital by EMS

### Data Source Hierarchy Guide

- 1 EMS Run Report

### References to Other Databases

- Not an NTDS element

## SCENE INTERVENTIONS

---

### Element Intent

This provides information regarding the patient's prehospital care and health status.

### Description

*Scene Interventions* indicates whether a critical procedure was performed by EMS at the scene or en route to your hospital, and if so, the procedure that was performed.

### Element Values

- 1 CPR
- 2 Needle Thoracostomy or Chest Tube
- 3 Nasal Endotracheal Tube
- 4 Oral Endotracheal Tube
- 5 Surgical Airway (i.e. surgical, needle or percutaneous cricothyrotomy, tracheostomy)
- 6 Other Non-Surgical Airway (Supraglottic Airway (e.g., Laryngeal Mask Airway, King, Combitube))
- 7 Tourniquet

### Common Null Values

- Accepted

### Additional Information

- Select *NA* If the patient was not treated at the scene by EMS

### Data Source Hierarchy Guide

- 1 EMS Run Report

### References to Other Databases

- Not an NTDS element



## PREHOSPITAL CARDIAC ARREST

---

### Element Intent

Pre-hospital cardiac arrest is associated with increased risk of mortality which could impact care decisions and increase the risk of adverse outcomes.

### Description

*Prehospital Cardiac Arrest* is indication of whether patient experienced cardiac arrest prior to ED/Hospital arrival.

### Element Values

- 1 Yes
- 2 No

### Common Null Values

- Accepted

### Additional Information

- A patient who experienced a sudden cessation of cardiac activity. The patient was unresponsive with no normal breathing and no signs of circulation
- The event must have occurred outside of the reporting hospital, prior to admission at the center in which the registry is maintained.
- Pre-hospital cardiac arrest could occur at a transferring institution
- Any component of basic and/or advanced cardiac life support must have been initiated

### Data Source Hierarchy Guide

- 1 EMS Run Report
- 2 Nursing Notes/Flow Sheet
- 3 History & Physical
- 4 Transfer Facility Records

### References to Other Databases

- NTDS 2026

## INTUBATION PRIOR TO ARRIVAL

---

### Element Intent

To indicate respiratory compromise, which could increase the risk of adverse outcomes.

### Description:

The patient is intubated with a definitive airway due to this injury prior to arrival at your hospital.

### INCLUDE:

- Definitive airways placed below the vocal cords (e.g., endotracheal tube (ET), tracheostomy, cricothyroidotomy).

### EXCLUDE:

- Airways not placed below the vocal cords (e.g., combitube, KING, laryngeal mask airway (LMA), I-Gel).

### Element Values

1. Yes
2. No

### Additional Information :

- If Element Value “1. Yes” is reported, report Intubation Location.
- The Null value “Not Applicable” is reported for patients who had an established airway prior to this injury event (e.g., Chronic Ventilator Dependence).

### Data Source Hierarchy Guide

1. Triage/Trauma Flow Sheet
2. ED Record
3. Face Sheet
4. Billing Sheet
5. Discharge Summary
6. Transfer Facility Records

### References to Other Databases

- NTDS 2026

## INTUBATION LOCATION

---

### Element Intent

To determine if the patient was intubated inside or outside of a hospital setting, which can inform outreach and improve prehospital care strategies.

### Description

The location the patient was intubated at prior to hospital arrival.

### Element Values

1. Out of the Hospital Intubation
2. Transferring Facility

### Additional Information

- Only reported if Intubation Prior to Arrival is Element Value “1. Yes”.
- The null value “Not Applicable” is reported if Intubation Prior to Arrival is reported as Element Value “2. No”
- The null value “Not Applicable” is reported if Intubation Prior to Arrival is reported as “Not Applicable.”
- The null value “Not Known/Not Recorded” is reported if Intubation Prior to Arrival is reported as “Not Known/Not Recorded.”
- Element Value “1. Out of Hospital Intubation” included intubations performed in the field, during transport occurred, report the location of the first intubation.

### Data Source Hierarchy Guide

1. Triage/Trauma Flow Sheet
2. ED Record
3. Face Sheet
4. Billing Sheet
5. Discharge Summary
6. Transfer Facility Records

### References to Other Databases

- NTDS 2026

## INTER-FACILITY TRANSFER

---

### Element Intent

To analyze system utilization and outcomes.

### Description

Was the patient transferred to your facility from another acute care facility?

### Element Values

- 1 Yes
- 2 No

### Common Null Values

- Accepted

### Additional Information

- Include, patients who require physical transfer from a free-standing emergency department (ED) to an affiliated trauma center.
- Patients transferred from a private doctor's office or stand-alone ambulatory surgery centers are NOT considered inter-facility transfers.
- Outlying facilities purporting to provide emergency care services or utilized to stabilize a patient are considered acute care facilities.
- Acute Care Hospital is defined as a hospital that provides inpatient medical care and other related services for surgery, acute medical conditions, or injuries (usually for a short-term illness or condition). "CMS Data Navigator Glossary of Terms", [https://www.cms.gov/Research-Statistics-Data-and-Systems/Research/ResearchGenInfo/Downloads/DataNav\\_Glossary\\_Alpha.pdf](https://www.cms.gov/Research-Statistics-Data-and-Systems/Research/ResearchGenInfo/Downloads/DataNav_Glossary_Alpha.pdf).

### Data Source Hierarchy Guide

1. EMS Run Report
2. Triage/Trauma Flow sheet
3. History & Physical
4. Transfer Facility Records

### References to Other Databases

- NTDS 2026

## TRANSFERRING HOSPITAL CODE

---

### Element Intent

If the patient was transferred from another hospital, this identifies that facility. This helps for data linkage purposes and to also track the level of care the patient received.

### Description

*Transferring Hospital Code* documents the Ohio Department of Public Safety (ODPS) assigned-number for the acute care facility which transferred a trauma patient to your hospital.

### Element Values

1. Four-digit hospital code assigned by the Ohio Department of Public Safety.

### Common Null Values

- Accepted

### Data Source Hierarchy Guide

- 1 ED Record
- 2 History & Physical

### References to Other Databases

- Not an NTDS element

## ED/HOSPITAL ARRIVAL DATE

---

### Element Intent

To calculate metrics such as hospital length of stay, provider response times, and medical intervention start times.

### Description

*ED/Hospital Arrival Date* is the date that the patient arrived at your ED/hospital.

### Element Values

1. Relevant value for data element

### Common Null Values

- Accepted

### Additional Information

- If the patient was brought to the ED, enter date patient arrived at ED. If the patient was directly admitted to the hospital, enter date patient was admitted to the hospital
- Collected as YYYY-MM-DD

### Data Source Hierarchy Guide

- 1 Triage/Trauma Flow Sheet
- 2 ED Record
- 3 Face Sheet
- 4 Billing Sheet
- 5 Discharge Summary

### References to Other Databases

- NTDS 2026

## ED/HOSPITAL ARRIVAL TIME

---

### Element Intent

To calculate metrics such as hospital length of stay, provider response times, and medical intervention start times.

### Description

*ED/Hospital Arrival Time* is the time of day that the patient arrived to your ED/hospital.

### Element Values

1. Relevant value for data element

### Common Null Values

- Accepted

### Additional Information

1. If the patient was brought to your hospital ED, enter the time patient arrived at the ED. If the patient was a directly admit to your hospital and bypassed the ED, enter that time that the patient was admitted to your hospital.
2. Collected as HHMM military time

### Data Source Hierarchy Guide

- 1 Triage/Trauma Flow Sheet
- 2 ED Record
- 3 Face Sheet
- 4 Billing Sheet
- 5 Discharge Summary

### References to Other Databases

1. NTDS 2026

## HIGHEST ACTIVATION

---

### Element Intent

To analyze response times, under/over triage, and resource utilization.

### Description

Patient received the highest level of trauma activation at your hospital.

### INCLUDE:

1. Patients who received the highest level of trauma activation initiated by emergency medical services (EMS) or by emergency department (ED) personnel at your hospital.
2. Patients who received the highest level of trauma activation initiated by EMS or by ED personnel at your hospital and were downgraded after arrival to your center.
3. Patients who received a lower level of trauma activation initiated by EMS or by ED personnel at your hospital and were upgraded to the highest level of trauma activation.

### EXCLUDE:

1. Patients who received the highest level of trauma activation after ED discharge.

### Element Values

- 1 Yes
- 2 No

### Additional Information

- Highest level of activation is defined by your hospital's criteria.
- Non-Trauma Center Facilities report the null value "Not Applicable" (NA)

### Data Source Hierarchy Guide

- 1 Triage/Trauma Flow Sheet
- 2 ED Record
- 3 History & Physical
- 4 Physician Notes
- 5 Discharge Summary

### References to Other Databases

- NTDS 2026



## TRAUMA SURGEON ARRIVAL DATE

---

### Element Intent

To analyze provider response times.

### Description

The date the first trauma surgeon arrived at the patient's bedside.

### Element Values

1. Relevant value for data element

### Additional Information

- Reported as YYYY-MM-DD.
- Limit reporting to the 24 hours after ED/Hospital arrival.
- The trauma surgeon leads the trauma team and is responsible for the overall care of trauma patient, including coordinating care with other specialties and maintaining continuity of care.
- The null value "Not Applicable" is reported for those patients who were not evaluated by a trauma surgeon within 24 hours of ED/Hospital arrival.
- The null value "Not Applicable" is reported if the data element *Highest Activation* is reported as *Element Value* "2. No."

### Data Source Hierarchy Guide

- 1 Triage/Trauma Flow Sheet
- 2 History & Physical
- 3 Physician Notes
- 4 Nursing Notes

### References to Other Databases

- NTDS 2026

## TRAUMA SURGEON ARRIVAL TIME

---

### Element Intent

To analyze provider response times.

### Description

The time the first trauma surgeon arrived at the patient's bedside.

### Element Values

1. Relevant value for data element

### Additional Information

- Reported as HHMM.
- Limit reporting to the 24 hours after ED/Hospital arrival.
- The trauma surgeon leads the trauma team and is responsible for the overall care of trauma patient, including coordinating care with other specialties and maintaining continuity of care.
- The null value "Not Applicable" is reported for those patients who were not evaluated by a trauma surgeon within 24 hours of ED/Hospital arrival.
- The null value "Not Applicable" is reported if the data element *Highest Activation* is reported as *Element Value* "2. No."

### Data Source Hierarchy Guide

- 1 Triage/Trauma Flow Sheet
- 2 History & Physical
- 3 Physician Notes
- 4 Nursing Notes

### References to Other Databases

- NTDS 2026

## INITIAL ED/HOSPITAL SYSTOLIC BLOOD PRESSURE

---

### Element Intent

A critical indicator of hemodynamic stability on arrival, which could impact care decisions, increase the risk of adverse outcomes, and prolong the length of stay.

### Description

*ED/Hospital Initial Systolic Blood Pressure* is the patient's first recorded systolic blood pressure within 30 minutes or less of ED/hospital arrival.

### Element Values

1. Relevant value for data element

### Common Null Values

- Accepted

### Additional Information

- Please note that first recorded/ hospital vitals do not need to be from the same assessment
- Measurement reported must be without the assistance of CPR or any type of mechanical chest compression device. For those patients who are receiving CPR or any type of mechanical chest compressions, report the value obtained while compressions are paused

### Data Source Hierarchy Guide

- 1 Triage/Trauma/Hospital Flow Sheet
- 2 Nurses Notes/Flow Sheet
- 3 Physician Notes
- 4 History & Physical

### References to Other Databases

- NTDS 2026

## INITIAL ED/HOSPITAL PULSE RATE

---

### Element Intent

A critical indicator of the body's response to injury and blood loss, which could impact care decisions, increase the risk of adverse outcomes, and prolong the length of stay.

### Description

*ED/Hospital Initial Pulse Rate* is the patient's first recorded pulse rate within 30 minutes or less of ED/hospital arrival (palpated or auscultated), expressed as a number per minute.

### Element Values

1. Relevant value for data element

### Common Null Values

- Accepted

### Additional Information

- Please note that first recorded/ hospital vitals do not need to be from the same assessment
- Measurement reported must be without the assistance of CPR or any type of mechanical chest compression device. For those patients who are receiving CPR or any type of mechanical chest compressions, report the value obtained while compressions are paused

### Data Source Hierarchy Guide

- 1 Triage/Trauma/Hospital Flow Sheet
- 2 Nurses Notes/Flow Sheet

### References to Other Databases

- NTDS 2026

## INITIAL ED/HOSPITAL RESPIRATORY RATE

---

### Element Intent

A critical indicator of the body's overall physiological condition, which could impact care decisions, increase the risk of adverse outcomes, and prolong the length of stay.

### Description

*ED/Hospital Initial Respiratory Rate* is the patient's first recorded respiratory rate within 30 minutes or less of ED/hospital arrival (expressed as a number per minute).

### Element Values

1. Relevant value for data element

### Common Null Values

- Accepted

### Additional Information

- If documented, report additional element *Initial ED/Hospital Respiratory Assistance*
- Please note that first recorded hospital vitals do not need to be from the same assessment

### Data Source Hierarchy Guide

- 1 Triage/Trauma/Hospital Flow Sheet
- 2 Nurses Notes/Flow Sheet
- 3 Respiratory Therapy Notes/Flow Sheet

### References to Other Databases

- NTDS 2026

## INITIAL ED/HOSPITAL RESPIRATORY ASSISTANCE

---

### Element Intent

A critical indicator of the patient's respiratory status on arrival, which could impact care decisions, increase the risk of adverse outcomes, and prolong the length of stay.

### Description

*ED/Hospital Initial Respiratory Assistance* documents whether the patient was receiving respiratory assistance within 30 minutes or less of ED/hospital arrival.

### Element Values

- 1 Unassisted Respiratory Rate
- 2 Assisted Respiratory Rate

### Common Null Values

- Accepted

### Additional Information

- Only reported if *Initial ED/Hospital Respiratory Rate* is documented
- Respiratory Assistance is defined as mechanical and/or external support of respiration
- Please note that first recorded/ hospital vitals do not need to be from the same assessment
- The null value "Not Applicable" is reported if "Initial ED/Hospital Respiratory Rate" is "Not Known/Not Recorded"

### Data Source Hierarchy Guide

- 1 Triage/Trauma/Hospital Flow Sheet
- 2 Nurses Notes/Flow Sheet
- 3 Respiratory Therapy Notes/Flow Sheet

### References to Other Databases

- NTDS 2026

## INITIAL ED/HOSPITAL OXYGEN SATURATION

---

### Element Intent

A critical indicator of the patient's ability to deliver oxygen to tissues, which could impact care decisions, increase the risk of adverse outcomes, and prolong the length of stay.

### Description

*ED/Hospital Initial Oxygen Saturation* is the patient's first recorded oxygen saturation within 30 minutes or less of ED/hospital arrival, expressed as a percentage.

### Element Values

1. Relevant value for data element

### Common Null Values

- Accepted

### Additional Information

- If documented, report additional element *Initial ED/Hospital Supplemental Oxygen*
- Please note that first recorded hospital vitals do not need to be from the same assessment

### Data Source Hierarchy Guide

- 1 Triage/Trauma/Hospital Flow Sheet
- 2 Nurses Notes/Flow Sheet
- 3 Respiratory Therapy Notes/Flow Sheet

### References to Other Databases

- NTDS 2026

## INITIAL ED/HOSPITAL SUPPLEMENTAL OXYGEN

---

### Element Intent

To determine whether the recorded initial ED/Hospital Oxygen Saturation value reflects the underlying patient condition alone or was influenced by oxygen therapy.

### Description

*ED/Hospital Supplemental Oxygen* is whether supplemental oxygen was provided to the patient during the assessment of *ED/Hospital Initial Oxygen Saturation Level* within 30 minutes or less of ED/hospital arrival.

### Element Values

- 1 No Supplemental Oxygen
- 2 Supplemental Oxygen

### Common Null Values

1. Accepted

### Additional Information

- The null value “Not Applicable” is reported if the *Initial ED/Hospital Oxygen Saturation* is “Not Known/Not Recorded”
- Please note that first recorded hospital vitals do not need to be from the same assessment

### Data Source Hierarchy Guide

- 1 Triage/Trauma/Hospital Flow Sheet
- 2 Nurses Notes/Flow Sheet

### References to Other Databases

- NTDS 2026



## INITIAL ED/HOSPITAL TEMPERATURE

---

### Element Intent

A critical indicator for the presence of hypothermia, which could impact care decisions, increase the risk of adverse outcomes, and prolong the length of stay.

### Description

*Initial ED/Hospital Temperature* is the patient's first recorded temperature within 30 minutes or less of ED/hospital arrival, documented in degrees Fahrenheit.

### Element Values

1. Relevant value for data element

### Common Null Values

- Accepted

### Additional Information

- Please note that first recorded hospital vitals do not need to be from the same assessment

### Data Source Hierarchy Guide

- 1 Triage/Trauma/Hospital Flow Sheet
- 2 Nurses Notes/Flow Sheet

### References to Other Databases

- NTDS 2026

## INITIAL ED/HOSPITAL GCS - EYE

---

### Element Intent

The GCS-Eye score is one component of the Total GCS and provides information on the severity of neurologic impairment. Collecting the initial provides identification of the patient's state on arrival.

### Description

*Initial ED/Hospital GCS Eye Opening* is the patient's first recorded Glasgow Coma Score (GCS) eye assessment documented within 30 minutes or less of ED/hospital arrival in your ED/hospital.

### Element Values

- 1 No eye movement when assessed
- 2 Opens eyes in response to painful stimulation
- 3 Opens eyes in response to verbal stimulation
- 4 Opens eyes spontaneously

### Common Null Values

- Accepted

### Additional Information

- If a patient does not have a numeric GCS score recorded, but written documentation closely (or directly) relates to verbiage describing a specific level of functioning within the GCS scale, the appropriate numeric score may be listed. E.g. the chart indicates: "patient's pupils are PERRL," an Eye GCS of 4 may be recorded, IF there is no other contradicting documentation.
- Please note that first recorded/ hospital vitals do not need to be from the same assessment.
- The null value "Not Known/Not Recorded" is reported if the patient's Initial ED/Hospital GCS – Eye was not measured within 30 minutes or less of ED/hospital arrival.

### Data Source Hierarchy Guide

- 1 Triage/Trauma/Hospital Flow Sheet
- 2 Nurses Notes/Flow Sheet
- 3 Physician Notes/Flow Sheet

### References to Other Databases

- NTDS 2026

## INITIAL ED/HOSPITAL GCS - VERBAL

---

### Element Intent

The GCS-Verbal score is one component of the Total GCS and provides information on the severity of neurologic impairment. Collecting the initial provides identification of the patient's state on arrival.

### Description

*ED/Hospital Initial GCS Verbal Response* is the patient's first recorded Glasgow Coma Score verbal assessment documented within 30 minutes or less of ED/hospital arrival.

### Element Values

- |  |  |
|--|--|
| <ul style="list-style-type: none"><li>• <u>Pediatric(&lt;= 2 years of age)</u><ol style="list-style-type: none"><li>1 No vocal response</li><li>2 Inconsolable, agitated</li><li>3 Inconsistently consolable, moaning</li><li>4 Cries but is consolable, inappropriate interactions</li><li>5 Smiles, oriented to sounds, follows objects, interacts</li></ol></li></ul> | <ul style="list-style-type: none"><li>• <u>Adult</u><ol style="list-style-type: none"><li>1 No verbal response</li><li>2 Incomprehensible sounds</li><li>3 Inappropriate words</li><li>4 Confused</li><li>5 Oriented</li></ol></li></ul> |
|--|--|

### Common Null Values

- Accepted

### Additional Information

- If patient is intubated then the GCS Verbal score is equal to 1
- If a patient does not have a numeric GCS score recorded, but written documentation closely (or directly) relates to verbiage describing a specific level of functioning within the GCS scale, the appropriate numeric score may be listed. E.g. the chart indicates: "patient is oriented to person place and time," a Verbal GCS of 5 may be recorded, IF there is no other contradicting documentation.
- Please note that first recorded/ hospital vitals do not need to be from the same assessment
- The null value "Not Known/Not Recorded" is reported if the patient's Initial ED/Hospital GCS – Verbal was not measured within 30 minutes or less of ED/hospital arrival

### Data Source Hierarchy Guide

- 1 Triage/Trauma/ Hospital Flow Sheet
- 2 Nurses Notes/Flow Sheet
- 3 Physician Notes/Flow Sheet

### References to Other Databases

- NTDS 2026

## INITIAL ED/HOSPITAL GCS - MOTOR

---

### Element Intent

The GCS-Motor score is one component of the Total GCS and provides information on the severity of neurologic impairment. Collecting the initial provides identification of the patient's state on arrival.

### Description

*ED/Hospital Initial GCS Motor Response* is the patient's first recorded Glasgow Coma Score motor assessment documented within 30 minutes or less of ED/hospital arrival.

### Element Values

- |  |                        |
|--|------------------------|
| • <u>Pediatric(&lt;= 2 years of age)</u> | • <u>Adult</u>         |
| 1 No motor response                      | 1 No motor response    |
| 2 Extension to pain                      | 2 Extension to pain    |
| 3 Flexion to pain                        | 3 Flexion to pain      |
| 4 Withdrawal from pain                   | 4 Withdrawal from pain |
| 5 Localizing pain                        | 5 Localizing pain      |
| 6 Appropriate response to stimulation    | 6 Obeys commands       |

### Common Null Values

- Accepted

### Additional Information

- If a patient does not have a numeric GCS score recorded, but written documentation closely (or directly) relates to verbiage describing a specific level of functioning within the GCS scale, the appropriate numeric score may be listed. E.g. the chart indicates: "patient withdraws from a painful stimulus," a Motor GCS of 4 may be recorded, IF there is no other contradicting documentation.
- Please note that first recorded hospital vitals do not need to be from the same assessment
- The null value "Not Known/Not Recorded" is reported if the patient's Initial ED/Hospital GCS – Motor was not measured within 30 minutes or less of ED/hospital arrival

### Data Source Hierarchy Guide

- 1 Triage/Trauma/ Hospital Flow Sheet
- 2 Nurses Notes/Flow Sheet
- 3 Physician Notes/Flow Sheet

### References to Other Databases

- NTDS 2026

## INITIAL ED/HOSPITAL GCS - TOTAL

---

### Element Intent

The GCS-Total score is used to gauge the severity of neurologic impairment. Collecting the initial provides identification of the patient's state on arrival.

### Description

*ED/Hospital Initial GCS Total Score* is the patient's first recorded Glasgow Coma Score documented within 30 minutes or less of ED/hospital arrival in your ED/hospital.

### Element Values

1. Relevant value for data element

### Common Null Values

- Accepted

### Additional Information

- If a patient does not have a numeric GCS recorded, but there is documentation related to their level of consciousness such as "AAOx3," "awake alert and oriented," or "patient with normal mental status," interpret this as GCS of 15 IF there is no other contradicting documentation.
- Please note that first recorded/hospital vitals do not need to be from the same assessment
- The null value "Not Known/Not Recorded" is reported if Initial ED/Hospital GCS – Eye, Initial ED/Hospital GCS – Motor, Initial ED/Hospital GCS – Verbal were not measured within 30 minutes or less of ED/Hospital arrival

### Data Source Hierarchy Guide

- 1 Triage/Trauma/ Hospital Flow Sheet
- 2 Nurses Notes/Flow Sheet
- 3 Physician Notes/Flow Sheet

### References to Other Databases

- NTDS 2026

## INITIAL ED/HOSPITAL GCS ASSESSMENT QUALIFIERS

---

### Element Intent

The GCS-Assessment Qualifier(s) indicate a GCS that might be altered due to a medical intervention.

### Description

*ED/Hospital Initial GCS Qualifiers* are factors that potentially affected the patient's first Glasgow Coma Score assessment within 30 minutes or less of ED/hospital arrival.

### Element Values

- 1 Patient Chemically Sedated or Paralyzed
- 2 Obstruction to the Patient's Eye
- 3 Patient Intubated
- 4 Valid GCS: Patient was not sedated, not intubated, and did not have obstruction to the eye

### Common Null Values

- Accepted

### Additional Information

- Report all that apply.
- Identifies treatments given to the patient that may affect the first GCS assessment. This element does not apply to self-medications the patient may administer (i.e., ETOH, prescriptions, etc.).
- *Element Value* "1. Patient Chemically Sedated or Paralyzed" is reported if an intubated patient has recently received an agent that results in neuromuscular blockade such that a motor or eye response is not possible.
- Neuromuscular blockade is typically induced following the administration of agent like succinylcholine, mivacurium, rocuronium, (cis)atracurium, vecuronium, or pancuronium. While these are the most common agents, please review what might be typically used in your center so it can be identified in the medical record.
- Each of these agents has a slightly different duration of action, so their effect on the GCS depends on when they were given. For example, succinylcholine's effects last for only 5-10 minutes.
- Please note that the first recorded hospital vitals do not need to be from the same assessment.
- The null value "Not Known/Not Recorded" is reported if **Initial ED/Hospital GCS-40** is reported.
- The null value "Not Known/Not Recorded" is reported if the **Initial ED/Hospital GCS Assessment Qualifiers** are not documented within 30 minutes of ED/hospital arrival.

### Data Source Hierarchy Guide

- 1 Triage/Trauma/ Hospital Flow Sheet
- 2 Nurses Notes/Flow Sheet
- 3 Physician Notes/Flow Sheet

### References to Other Databases

- NTDS 2026

# HEIGHT

---

## Element Intent

To calculate body mass index (BMI) which could impact care decisions and increase the risk of adverse outcomes.

## Description

*Height* is the patient's height in centimeters.

## Element Values

1. Height in centimeters

## Common Null Values

- Accepted

## Additional Information

- Recorded in centimeters
- May be based on family or self-report
- Please note that first recorded/hospital vitals do not need to be from the same assessment
- The null value "Not Known/Not Recorded" is reported if the patient's Initial ED/Hospital Height was not measured

## Data Source Hierarchy Guide

- 1 Triage/Trauma/Hospital Flow Sheet
- 2 Nurses Notes/Flow Sheet
- 3 Pharmacy Record

## References to Other Databases

- NTDS 2026

**NOTE:** HEIGHT differs from NTDS. Please refer to the section "DIFFERENCES BETWEEN OHIO AND NATIONAL TRAUMA DATA STANDARD (NTDS)" starting on Page 12 for more information.

## WEIGHT

---

### Element Intent

To calculate body mass index (BMI) which could impact care decisions and increase the risk of adverse outcomes.

### Description

*Weight* is the patient's weight in kilograms.

### Element Values

1. Weight in kilograms

### Common Null Values

- Accepted

### Additional Information

- Recorded in kilograms
- May be based on family or self-report
- Please note that first recorded/hospital vitals do not need to be from the same assessment
- The null value "Not Known/Not Recorded" is reported if the patient's Initial ED/Hospital Weight was not measured

### Data Source Hierarchy Guide

- 1 Triage/Trauma/Hospital Flow Sheet
- 2 Nurses Notes/Flow Sheet
- 3 Pharmacy Record

### References to Other Databases

- NTDS 2026

**NOTE:** WEIGHT differs from NTDS. Please refer to the section "DIFFERENCES BETWEEN OHIO AND NATIONAL TRAUMA DATA STANDARD (NTDS)" starting on Page 12 for more information.



## ED DISCHARGE ORDER WRITTEN DATE

---

### Element Intent

Provides information about the patient's ED disposition, which helps track the level of care the patient received.

### Description

*ED Discharge Order Written Date* is the date that the order was written for the patient to be discharged from your ED.

### Element Values

1. Relevant value for data element

### Common Null Values

- Accepted

### Additional Information

- The null value "Not Applicable" is reported if the patient is directly admitted to the hospital.
- If ED Discharge Disposition is "5. Deceased/Expired," then ED Discharge Date is the date of death as indicated on the patient's death certificate
- Collected as YYYY-MM-DD

### Data Source Hierarchy Guide

- 1 Hospital Discharge Summary
- 2 Billing Sheet/Medical Records Coding Summary Sheet
- 3 Physicians' Progress Notes

### References to Other Databases

- Not an NTDS element

**NOTE:** ED DISCHARGE ORDER WRITTEN DATE differs from NTDS. Please refer to the section "DIFFERENCES BETWEEN OHIO AND NATIONAL TRAUMA DATA STANDARD (NTDS)" starting on Page 12 for more information.

## ED DISCHARGE ORDER WRITTEN TIME

---

### Element Intent

Provides information about the patient's ED disposition, which helps track the level of care the patient received.

### Description

*ED Discharge Order Written Time* is the time that the order was written for the patient to be discharged from your ED.

### Element Values

1. Relevant value for data element

### Common Null Values

- Accepted

### Additional Information

- The null value "Not Applicable" is reported if the patient is directly admitted to the hospital.
- If ED Discharge Disposition is "5. Deceased/Expired," then ED Discharge Time is the time of death as indicated on the patient's death certificate
- Collected as HHMM military time

### Data Source Hierarchy Guide

- 1 Hospital Discharge Summary
- 2 Billing Sheet/Medical Records Coding Summary Sheet
- 3 Physicians' Progress Notes

### References to Other Databases

- Not an NTDS element

**NOTE:** ED DISCHARGE ORDER WRITTEN TIME differs from NTDS. Please refer to the section "DIFFERENCES BETWEEN OHIO AND NATIONAL TRAUMA DATA STANDARD (NTDS)" starting on Page 12 for more information.

## ED DISCHARGE DATE

---

### Element Intent

To calculate metrics such as hospital length of stay and to inform the care timeline.

### Description

*ED Discharge Date* is the date that the patient was physically discharged from your ED.

### Element Values

1. Relevant value for data element

### Common Null Values

- Accepted

### Additional Information

- Collected as YYYY-MM-DD
- The null value "Not Applicable" is reported if the patient is directly admitted to the hospital.
- If ED Discharge Disposition is "5. Deceased/Expired," then ED Discharge Date is the date of death as indicated on the patient's death certificate

### Data Source Hierarchy Guide

- 1 Physician Order
- 2 ED Record
- 3 Triage/Trauma/Hospital Flow Sheet
- 4 Nursing Notes/Flow Sheet
- 5 Discharge Summary
- 6 Billing Sheet
- 7 Progress Notes

### References to Other Databases

- NTDS 2026 (element name only)

**NOTE:** ED DISCHARGE DATE differs from NTDS. Please refer to the section "DIFFERENCES BETWEEN OHIO AND NATIONAL TRAUMA DATA STANDARD (NTDS)" starting on Page 12 for more information.

## ED DISCHARGE TIME

---

### Element Intent

To calculate metrics such as hospital length of stay and to inform the care timeline.

### Description

*ED Discharge Time* is the time that the patient was physically discharged from your ED.

### Element Values

1. Relevant value for data element

### Common Null Values

- Accepted

### Additional Information

- Collected as HHMM military time
- The null value "Not Applicable" is reported if the patient is directly admitted to the hospital.
- If ED Discharge Disposition is "5. Deceased/Expired," then ED Discharge Time is the time of death as indicated on the patient's death certificate

### Data Source Hierarchy Guide

- 1 Physician Order
- 2 ED Record
- 3 Triage/Trauma/Hospital Flow Sheet
- 4 Nursing Notes/Flow Sheet
- 5 Discharge Summary
- 6 Billing Sheet
- 7 Progress Notes

### References to Other Databases

- NTDS 2026 (element name only)

**NOTE:** ED DISCHARGE TIME differs from NTDS. Please refer to the section "DIFFERENCES BETWEEN OHIO AND NATIONAL TRAUMA DATA STANDARD (NTDS)" starting on Page 12 for more information.

## ED DISCHARGE DISPOSITION

---

### Element Intent

To indicate the patient's medical needs after their ED phase of care is complete.

### Description

The disposition unit the order was written for the patient to be discharged from the ED.

### Element Values

- |   |   |
|---|---|
| 1 Floor bed (general admission, non-specialty unit bed) | 7 Operating Room (Hybrid OR)                                    |
| 2 Observation unit                                      | 8 Intensive Care Unit (ICU)                                     |
| 3 Telemetry/step-down unit (less acuity than ICU)       | 9 Home without services   |
| 4 Home with services                                    | 10 Left against medical advice                                  |
| 5 Deceased/Expired                                      | 11 Transferred to another hospital                              |
| 6 Other (jail, institutional care, mental health, etc.) | 12 Interventional Radiology Suite                               |
|   | 13 Hospice (e.g., hospice facility, hospice unit, Home hospice) |

### Common Null Values

- Accepted

### Additional Information

- The null value "Not Applicable" is reported if the patient is directly admitted to the hospital
- If ED Discharge Disposition is 4, 5, 6, 9, 10, 11 or 13 then Hospital Discharge Date, Hospital Discharge Time and Hospital Discharge Disposition must be "Not Applicable"
- If the patient was boarded in the ED, the disposition must be the location the patient was ordered to go when their ED workup was complete.

### Data Source Hierarchy Guide

- 1 Physician Order
- 2 Discharge Summary
- 3 Nursing Notes/Flow Sheet
- 4 Case Management/Social Services Notes
- 5 ED Record
- 6 History & Physical

### References to Other Databases

- NTDS 2026

**NOTE:** ED DISCHARGE DISPOSITION differs from NTDS. Please refer to the section "DIFFERENCES BETWEEN OHIO AND NATIONAL TRAUMA DATA STANDARD (NTDS)" starting on Page 12 for more information.

## ED TRANSFER TO HOSPITAL

---

### Element Intent

Provides information about the patient's ED disposition, which helps track the level of care the patient received.

### Description

*ED Transfer to Hospital* is a subsequent hospital destination of the patient upon discharge from your ED.

### Element Values

1. Four-digit hospital code assigned by the Ohio Department of Public Safety.

### Common Null Values

- Accepted

### Additional Information

- The null value "Not Applicable" is reported if the patient is directly admitted to the hospital
- If ED Discharge Disposition is 4, 5, 6, 9, 10, 11 the Hospital Discharge date, Time, Disposition and Inpatient Transfer to Hospital should be "Not Applicable"

### Data Source Hierarchy Guide

- 1 ED Record
- 2 History & Physical

### References to Other Databases

- Not an NTDS element

## PRIMARY TRAUMA SERVICE TYPE

---

### Element Intent

To indicate the service primarily responsible for the patient's care because physiological, psychological, and developmental needs differ based on age, which helps to analyze resource utilization and outcomes.

### Description

The primary service type responsible for the care of this patient.

### Element Values

- 1 Adult
- 2 Pediatric

### Additional Information

- The primary service type responsible for trauma evaluation and care of the patient.
- This element will be used to determine which eligible Trauma Quality Programs report [adult or pediatric] the patient will appear; report age criteria will still apply.
- Adult trauma centers that do not have a separate pediatric service must report *Element Value* "1. Adult."
- Pediatric trauma centers that do not have a separate adult service must report *Element Value* "2. Pediatric."
- (OH Specific) Non-trauma facilities and Free-Standing Emergency Departments (FSEDs) should report *Element Value* "1. Adult" for patients age 16 and over; and *Element Value* "2. Pediatric" for patients age 15 and under.

### Data Source Hierarchy Guide

- 1 Triage/Trauma Flow Sheet
- 2 History and Physical
- 3 Discharge Summary

### References to Other Databases

- NTDS 2026

## PRIMARY MEDICAL EVENT

---

### Element Intent

To indicate pre-injury medical conditions that have a high risk of permanent disability or death which could impact care decisions and influence outcomes.

### Description:

The patient experienced a documented primary medical event (stroke, myocardial infarction, cardiac arrest, intracranial bleeding, sepsis) that immediately preceded the traumatic injury.

### Element Values

1. Yes
2. No

### Additional Information :

- Element Value “1. Yes” is reported if the patient experienced a medical event immediately preceding the trauma.
- The null value “Not Known/Not Recorded” is reported if it is unknown the primary medical event immediately preceded the traumatic injury.

### Data Source Hierarchy Guide

1. Physician’s Notes
2. History & Physical
3. Progress Notes
4. Case Management/Social Services
5. Nursing Notes/Flow Sheet
6. Triage/Trauma Flow Sheet
7. Discharge Summary
8. Transfer Facility Records

### References to Other Databases

- NTDS 2026



## ALCOHOL SCREEN

---

### Element Intent

To identify potential risks that could impact care decisions, increase the risk of adverse outcomes, prolong the length of stay, and to identify those that may benefit from intervention programs.

### Description

*Alcohol Screen* is a blood alcohol concentration (BAC) test was performed on the patient within 24 hours after first hospital encounter.

### Element Values

- 1 Yes
- 2 No

### Common Null Values

- a.* Not Accepted

### Additional Information

- Alcohol screen may be administered at any facility, unit or setting treating this patient event

### Data Source Hierarchy Guide

- 1 Lab Results
- 2 Transferring Facility Records

### References to Other Databases

- NTDS 2026

## ALCOHOL SCREEN RESULTS

---

### Element Intent

To identify potential risks that could impact care decisions, increase the risk of adverse outcomes, prolong the length of stay, and to identify those that may benefit from intervention programs.

### Description

*Alcohol Screen Results* is the first recorded blood alcohol concentration (BAC) results within 24 hours after first hospital encounter.

### Element Values

1. Relevant value for data element

### Common Null Values

- Accepted

### Additional Information

- Collect as X.XX grams per deciliter (g/dl)
- Record BAC results within 24 hours after first hospital encounter at either your facility or the transferring facility
- The null value “Not Applicable” is used for those patients who were not tested

### Data Source Hierarchy Guide

- 1 Lab Results
- 2 Transferring Facility Records

### References to Other Databases

- NTDS 2026

## DRUG SCREEN

---

### Element Intent

To identify potential risks that could impact care decisions, increase the risk of adverse outcomes, prolong the length of stay, and to identify those that may benefit from intervention programs.

### Description

*Drug Screen* is the first recorded positive drug screen within 24 hours after first hospital encounter (select all that apply).

### Element Values

- |                          |                                   |
|--------------------------|-----------------------------------|
| 1 AMP (Amphetamine)      | 9 OXY (Oxycodone)                 |
| 2 BAR (Barbiturate)      | 10 PCP (Phencyclidine)            |
| 3 BZO (Benzodiazepines)  | 11 TCA (Tricyclic Antidepressant) |
| 4 COC (Cocaine)          | 12 THC (Cannabinoid)              |
| 5 mAMP (Methamphetamine) | 13 Other                          |
| 6 MDMA (Ecstasy)         | 14 None                           |
| 7 MTD (Methadone)        | 15 Not Tested                     |
| 8 OPI (Opioid)           |                                   |

### Common Null Values

- β. Not Accepted

### Additional Information

- Report positive drug screen results within 24 hours after first hospital encounter, at either your facility or transferring facility
- “None” is reported for patients whose only positive results are due to drugs administered at any facility (or setting) treating this patient event, or for patients who were tested and had no positive results
- If multiple drugs are detected, only report drugs that were not administered at any facility (or setting) treating this patient event

### Data Source Hierarchy Guide

- 1 Lab Results
- 2 Transferring Facility Records

### References to Other Databases

- NTDS 2026

## ICD-10 HOSPITAL PROCEDURES

---

### Element Intent

To identify the types of hospital interventions.

### Description

*Hospital Procedures* are all operative and selected non-operative procedures conducted during hospital stay.

Operative and selected non-operative procedures are those that were essential to the diagnosis, stabilization, or treatment of the patient's specific injuries or complications. The list of procedures below should be used as a guide to non-operative procedures that should be provided to the OTR.

### Element Values

1. Major and minor procedure ICD-10 PCS procedure codes
2. The maximum number of procedures that may be reported for a patient is 200

### Common Null Values

- Accepted

### Additional Information

- The null value "Not Applicable" is reported if the patient did not have procedures
- Include only procedures performed at your institution
- Report all procedure performed in the operating room
- Report all procures in the ED, ICU, ward, or radiology department that were essential to the diagnosis, stabilization, or treatment of the patient's specific injuries or their complications
- Procedures with an asterisk have the potential to be performed multiple times during one episode of hospitalization. In this case, report only the first event. If there is no asterisk, report each event even if there is more than one.
- Plain radiography of whole body, Plain radiography of whole skeleton, and Plain radiography of infant whole body to the Diagnostic and Therapeutic Imaging.
- Note that the hospital may capture additional procedures

### Data Source Hierarchy Guide

- 1 Operative Reports
- 2 Procedure Notes
- 3 Trauma Flow Sheet
- 4 ED Record
- 5 Nursing Notes/Flow Sheet
- 6 Radiology Reports
- 7 Discharge Summary

### References to Other Databases

- NTDS 2026

## PROCEDURE LIST FOR *HOSPITAL PROCEDURES* ELEMENT

### **DIAGNOSTIC & THERAPEUTIC IMAGING**

Computerized tomographic studies\* (Head, Chest, Abdomen, Pelvis, C-Spine, T-Spine, L-Spine)  
Diagnostic ultrasound (includes FAST)\*  
Doppler ultrasound of extremities\*  
Angiography  
Angioembolization  
REBOA  
Inferior vena cava (IVC) filter  
Diagnostic imaging interventions on the total body  
Plain radiography of whole body  
Plain radiography of whole skeleton  
Plain radiography of infant whole body

### **CARDIOVASCULAR**

Open cardiac massage  
Cardiopulmonary Resuscitation (CPR)

### **CENTRAL NERVOUS SYSTEM**

Insertion of ICP monitor\*  
Ventriculostomy  
Cerebral oxygen monitoring\*

### **GASTROINTESTINAL**

Endoscopy (includes gastroscopy, sigmoidoscopy, colonoscopy)  
Gastrostomy/jejunostomy (percutaneous/or endoscopic)  
Percutaneous (endoscopic) gastrojejunoscopy

### **GENITOURINARY**

Ureteric catheterization (i.e. ureteric stent)  
Suprapubic cystostomy

### **MUSCULOSKELETAL**

Soft tissue/bony debridement\*  
Closed reduction fractures  
Skeletal (and halo) traction  
Fasciotomy

### **RESPIRATORY**

Insertion of endotracheal tube\* (Exclude intubations performed in the OR)  
Continuous invasive mechanical ventilation\*  
Chest tube\*  
Bronchoscopy\*  
Tracheostomy

### **TRANSFUSION**

The following blood products should be captured over first 24 hours after hospital arrival:  
Transfusion of red cells \*  
Transfusion of platelets \*  
Transfusion of plasma \*

**\*May be performed multiple times during hospitalization**

## PROCEDURE EPISODE

---

### Element Intent

Provides information about patient's care in the hospital.

### Description

*Procedure Episode* documents the frequency of operative visits. Each trip to the operating room should be identified in sequential order (regardless of number of procedures completed at that time).

### Element Values

- 1 First Operative Episode
- 2 Second Operative Episode
- 3 Third Operative Episode
- 4 Fourth Operative Episode
- 5 Fifth Operative Episode
- 6 Sixth Operative Episode
- 7 Seventh Operative Episode
- 8 Eighth Operative Episode
- 9 Ninth Operative Episode
- 10 Tenth or More Operative Episode

### Common Null Values

- Accepted

### Additional Information

- Include only those operative procedures performed at your hospital
- This element is linked to the *Hospital Procedures* element
- Leave element blank if procedure was not performed in the Operating Room
- All of the procedures done in the first OR visit would be Episode 1, all in visit 2 would be Episode 2, and so forth.

### Data Source Hierarchy Guide

- 1 Operative Reports

### References to Other Databases

- Not an NTDS element

## HOSPITAL PROCEDURE START DATE

---

### Element Intent

To determine the timeliness of hospital interventions.

### Description

*The date operative and selected non-operative procedures were performed.*

### Element Values

1. Relevant value for data element

### Common Null Values

- Accepted

### Additional Information

- This element is linked to the Hospital Procedures element
- Collected as YYYY-MM-DD

### Data Source Hierarchy Guide

- 1 Operative Reports
- 2 Procedure Notes
- 3 Trauma Flow Sheet
- 4 ED Record
- 5 Nursing Notes/Flow Sheet
- 6 Radiology Report
- 7 Discharge Summary

### References to Other Databases

- NTDS 2026

**NOTE:** HOSPITAL PROCEDURE START DATE differs from NTDS. Please refer to the section “DIFFERENCES BETWEEN OHIO AND NATIONAL TRAUMA DATA STANDARD (NTDS)” starting on Page 12 for more information.

## HOSPITAL PROCEDURE START TIME

---

### Element Intent

To determine the timeliness of hospital interventions.

### Description

*The time* operative and selected non-operative procedures were performed.

### Element Values

1. Relevant value for data element

### Common Null Values

- Accepted

### Additional Information

- This element is linked to the Hospital Procedures element
- Collected as HHMM military time
- Procedure start time is defined as the time that the incision was made (or the procedure started).
- If distinct procedures with the same procedure code are performed, their start time must be different.

### Data Source Hierarchy Guide

- 1 Operative Reports
- 2 Anesthesia Record
- 3 Procedure Notes
- 4 Trauma Flow Sheet
- 5 ED Record
- 6 Nursing Notes/Flow Sheet
- 7 Radiology Reports
- 8 Discharge Summary

### References to Other Databases

- NTDS 2026

**NOTE:** HOSPITAL PROCEDURE START TIME differs from NTDS. Please refer to the section “DIFFERENCES BETWEEN OHIO AND NATIONAL TRAUMA DATA STANDARD (NTDS)” starting on Page 12 for more information.



## ADVANCE DIRECTIVE LIMITING CARE

---

### Element Intent

Implementation of a previously signed advanced directive impacts care and influences outcomes.

### Description

The patient had a written request to limit life-sustaining treatment that restricted the care for the patient during this patient care event.

### Element Values

- 1 Yes
- 2 No

### Common Null Values

- Accepted

### Additional Information

- The written request was signed/dated by the patient and/or his/her designee prior to arrival at your center
- Report Element Value "2. No" for patients with Advanced Directives that did not limit life-sustaining treatments during this patient care event.
- Life-sustaining treatments include but are not limited to intubation, ventilator support, CPR, transfusion of blood products, dialysis or other forms of renal support, institution of medications to support blood pressure or cardiac function, or a specific surgical, interventional or radiological procedure (e.g. decompressive craniectomy, operation for hemorrhage control, angiography)
- The null value "Not Known/Not Recorded" is only reported if no past medical history is available.

### Data Source Hierarchy Guide


- 1 History & Physical
- 2 Physician's Notes
- 3 Progress Notes
- 4 Case Management/Social Services
- 5 Nursing Notes/Flow Sheet
- 6 Triage/Trauma Flow Sheet
- 7 Discharge Summary
- 8 Transfer Facility Records

### References to Other Databases

- NTDS 2026

## Advance Directive Limiting Care

2026 NTDS Data Dictionary, Released July 2025



1

Life-sustaining treatments include but are not limited to intubation, ventilator support, CPR, transfusion of blood products, dialysis or other forms of renal support, institution of medications to support blood pressure or cardiac function, or a specific surgical, interventional or radiological procedure (e.g., decompressive craniectomy, operation for hemorrhage control, angiography).

## ALCOHOL USE DISORDER

---

### Element Intent

Consumption of high levels of alcohol can affect the immune system, negatively affect wound healing, and increase the risk of developing infection, which could impact care decisions, increase the risk of adverse outcomes and prolong length of stay.

### Description

Descriptors documented in the medical record consistent with the diagnostic criteria of alcohol use disorder OR a diagnosis of alcohol use disorder documented in the patient's medical record.

### Element Values

- 1 Yes
- 2 No

### Common Null Values

- Accepted

### Additional Information

- Present prior to injury.
- Only report on patients  $\geq 15$  years-of-age.
- The null value "Not Applicable" must be reported for patients  $< 15$  years-of-age.
- The null value "Not Known/Not Recorded" is only reported if no past medical history is available for patients  $\geq 15$  years-of-age.
- Consistent with American Psychiatric Association (APA) DSM 5, 2013.
- Based on the patient's age on the day of arrival at your hospital.

### Data Source Hierarchy Guide


- 1 History & Physical
- 2 Physician's Notes
- 3 Progress Notes
- 4 Case Management/Social Services
- 5 Nursing Notes/Flow Sheet
- 6 Triage/Trauma Flow Sheet
- 7 Discharge Summary

### References to Other Databases

- NTDS 2026

## Alcohol Use Disorder

2026 NTDS Data Dictionary, Released July 2025



**1** Based on the patient's age on the day of arrival at your hospital.

**2** The NTDS definition is consistent with the American Psychological Association (APA) DSM 5, 2013. Refer to the APA and/or the TPM/TMD for more information.

## ANTICOAGULANT THERAPY

---

### Element Intent

Anticoagulants could induce greater risk of bleeding and increase the risk of adverse outcomes.

### Description

Documentation in the medical record of the administration of medication (anticoagulants, antiplatelet agents, thrombin inhibitors, thrombolytic agents) that interferes with blood clotting.

ANTICOAGULANTS	ANTIPLATELET AGENTS	THROMBIN INHIBITORS	THROMBOLYTIC AGENTS
Fondaparinux	Tirofiban	Bevalirudin	Alteplase
Warfarin	Dipyridamole	Argatroban	Reteplase
Dalteparin	Anagrelide	Lepirudin, Hirudin	Tenecteplase
Lovenox	Eptifibatide	Drotrecogin alpha	kabikinase
Pentasaccaride	Dipyridamole	Dabigatran	tPA
APC	Clopidogrel		
Ximelagatran	Cilostazol		
Pentoxifylline	Abciximab		
Rivaroxaban	Ticlopidine		
Apixaban	Prasugrel		
Heparin	Ticagrelor		

### Element Values

- 1 Yes
- 2 No

### Common Null Values

- Accepted

### Additional Information

- Present prior to injury.
- Anticoagulant must be part of the patient's active medication.
- Exclude patients whose only anticoagulant therapy is chronic Aspirin.
- The null value "Not Known/Not Recorded" is only reported if no past medical history is available.

### Data Source Hierarchy Guide


- 1 History & Physical
- 2 Physician's Notes
- 3 Progress Notes
- 4 Case Management/Social Services
- 5 Nursing Notes/Flow Sheet
- 6 Triage/Trauma Flow Sheet
- 7 Discharge Summary

### References to Other Databases

- NTDS 2026

# Anticoagulant Therapy

2026 NTDS Data Dictionary, Released July 2025



1

Examples of commonly prescribed medications that interfere with blood clotting can be found on the definition page. This list is not all-inclusive. If medication meets the definition criteria and is not included in the list, report *Element Value "1. Yes."*

## ATTENTION DEFICIT DISORDER/ATTENTION DEFICIT HYPERACTIVITY DISORDER (ADD/ADHD)

---

### Element Intent

Patients with ADD/ADHD experience impulsiveness, restlessness, and difficulty focusing on tasks which could impact care decisions, increase the risk of adverse outcomes, and prolong the length of stay.

### Description

A disorder involving inattention, hyperactivity, or impulsivity requiring medication for treatment.

### Element Values

- 1 Yes
- 2 No

### Common Null Values

- Accepted

### Additional Information

- Present prior to injury.
- A diagnosis of ADD/ADHD must be documented in the patient's medical record.
- The null value "Not Known/Not Recorded" is only reported if no past medical history is available.

### Data Source Hierarchy Guide


- 1 History & Physical
- 2 Physician's Notes
- 3 Progress Notes
- 4 Case Management/Social Services
- 5 Nursing Notes/Flow Sheet
- 6 Triage/Trauma Flow Sheet
- 7 Discharge Summary

### References to Other Databases

- NTDS 2026

## Attention Deficit Disorder/Attention Deficit Hyperactivity Disorder (ADD/ADHD)

2026 NTDS Data Dictionary, Released July 2025





## AUTISM SPECTRUM DISORDER (ASD) (PRE-EXISTING CONDITIONS)

---

### Element Intent

Patients with ASD experience problems with social communications and interaction, restricted or repetitive behaviors or interest, and/or different ways of learning, moving, or paying attention, which could impact care decisions and increase the risk of adverse outcomes.

### Description:

A disorder involving problems with social communication and interaction, and restricted or repetitive behaviors or interests as well as different ways of learning, moving, or paying attention.

### Element Values

1. Yes
2. No

### Additional Information :

- Present prior to injury.
- A diagnosis of ASD must be documented in the patient's medical record (e.g., autism, autism spectrum disorder, or Asperger's syndrome/disorder).
- The null value "Not Known/Not Recorded" is only reported if no past medical history is available.

### Data Source Hierarchy Guide


1. Physician Notes/Flow Sheet
2. History and Physical
3. Progress Notes
4. Case Management/Social Services Notes
5. Nursing Notes/Flow Sheet
6. Triage/Trauma Flow Sheet
7. Discharge Summary

### References to Other Databases

- NTDS 2026

## Autism Spectrum Disorder (ASD)

2026 NTDS Data Dictionary, Released July 2025



Might also be referred to as autism, ASD, or Asperger's syndrome/disorder.

## BIPOLAR I/II DISORDER

---

### Element Intent

Patients with Bipolar Disorder experience severe mood disturbances that cause clinically significant distress or impairment in social, occupational, or other important areas of functioning, which could impact care decisions and increase the risk of adverse outcomes.

### Description

A bipolar I/II disorder diagnosis documented in the medical record.

### Element Values

- 1 Yes
- 2 No

### Additional Information

- Present prior to injury.
- Only report on patients  $\geq 15$  years-of-age.
- The null value "Not Applicable" must be reported for patients  $< 15$  years-of-age.
- The null value "Not Known/Not Recorded" is only reported if no past medical history is available for patients  $\geq 15$  years-of-age.
- Based on patient's age on the day of arrival at your hospital.

### Data Source Hierarchy Guide


- 1 History and Physical
- 2 Physician Notes/Flow Sheet
- 3 Progress Notes
- 4 Case Management/Social Services Notes
- 5 Nursing Notes/Flow Sheet
- 6 Triage/Trauma Flow Sheet
- 7 Discharge Summary

### References to Other Databases

- NTDS 2026

# Bipolar I/II Disorder

2026 NTDS Data Dictionary, Released July 2025



**1** Based on the patient's age on the day of arrival at your hospital.

## BLEEDING DISORDER

---

### Element Intent

Underlying hematologic disorders result in a greater risk of bleeding which could increase the risk of adverse outcomes.

### Description

A group of conditions that result when the blood cannot clot properly.

### Element Values

- 1 Yes
- 2 No

### Common Null Values

- Accepted

### Additional Information

- Present prior to injury.
- A Bleeding Disorder diagnosis must be documented in the patient's medical record (e.g., Hemophilia, von Willenbrand Disease, Factor V Leiden).
- Consistent with American Society of Hematology, 2015.
- The null value "Not Known/Not Recorded" is only reported if no past medical history is available.

### Data Source Hierarchy Guide


- 1 History & Physical
- 2 Physician's Notes
- 3 Progress Notes
- 4 Case Management/Social Services
- 5 Nursing Notes/Flow Sheet
- 6 Triage/Trauma Flow Sheet
- 7 Discharge Summary

### References to Other Databases

- NTDS 2026

# Bleeding Disorder

2026 NTDS Data Dictionary, Released July 2025



**1** The NTDS definition is consistent with the American Society of Hematology, 2015.

**2** Some examples of bleeding disorders that affect blood clotting are hemophilia, von Willebrand Disease, and Factor V Leiden.  
If the bleeding disorder does not cause the patient's blood to clot improperly, e.g., sickle cell disease, report *Element Value* "2. No." Consult with the TPM or TMD if questioning a specific diagnosis.

## BRONCHOPULMONARY DYSPLASIA/CHRONIC LUNG DISEASE

---

### Element Intent

Bronchopulmonary Dysplasia/Chronic Lung Disease could induce negative respiratory and pulmonary function, which could impact care decisions and increase the risk of adverse outcomes.

### Description

The disorders which constitute Chronic Lung Disease (CLD) generally have a slow tempo of progression over many months or even years. The most common causes of CLD in children are Cystic Fibrosis (CF), and other causes of bronchiectasis (such as immunodeficiency, and in the third world, post -infective bronchiectasis (e.g., measles), Bronchopulmonary Dysplasia (BPD), or lung disease of prematurity).

### INCLUDE:

- Patients with a diagnosis of Cystic Fibrosis with pulmonary involvement.

### EXCLUDE:

- Patients with a diagnosis of Cystic Fibrosis with no documentation of lung disease.

### Element Values

1. Yes
2. No

### Additional Information

- Present prior to injury.
- Based on the patient's age on the day of arrival at your hospital.
- Only report on patients <15 years -of-age.
- The null value "Not Applicable" must be reported for patients >15 years-of-age.
- The null value "Not Known/Not Recorded" is only reported if no past medical history is available for patients <15 years -of-age.
- Examples of evidence of Cystic Fibrosis-associated pulmonary disease included, but are not limited to:
- Use of Chest Physiotherapy (CPT) or other airway clearing techniques.
- Vest therapy or intrapulmonary percussive ventilator.
- Intravenous, inhaled, or oral antibiotics to treat chronic respiratory infections related to Cystic Fibrosis.
- Consistent with the [ncbi.nlm.nih.gov](http://ncbi.nlm.nih.gov).

### Data Source Hierarchy Guide


1. History and Physical
2. Physician Notes/Flow Sheet
3. Progress Notes
4. Case Management/Social Services Notes
5. Nursing Notes/Flow Sheet
6. Triage/Trauma Flow Sheet
7. Discharge Summary

### References to Other Databases

- NTDS 2026

## Bronchopulmonary Dysplasia/Chronic Lung Disease

2026 NTDS Data Dictionary, Released July 2025



1

Based on the patient's age on the day of arrival at your hospital.

2

Include patients with a diagnosis of Cystic Fibrosis with pulmonary involvement.

Exclude patients with a diagnosis of Cystic Fibrosis with no documentation of lung disease.



## CEREBRAL VASCULAR ACCIDENT (CVA)

---

### Element Intent

Persistent residual motor sensory or cognitive deficits could impact care decisions and increase the risk of adverse outcomes.

### Description

A history prior to injury of a cerebrovascular accident (embolic, thrombotic, or hemorrhagic) with persistent residual motor sensory or cognitive dysfunction (e.g., hemiplegia, hemiparesis, aphasia, sensory deficit, impaired memory).

### Element Values

- 1 Yes
- 2 No

### Common Null Values

- Accepted

### Additional Information

- Present prior to injury.
- A diagnosis of CVA must be documented in the patient's medical record.
- The null value "Not Known/Not Recorded" is only reported if no past medical history is available.

### Data Source Hierarchy Guide


- 1 History & Physical
- 2 Physician's Notes
- 3 Progress Notes
- 4 Case Management/Social Services
- 5 Nursing Notes/Flow Sheet
- 6 Triage/Trauma Flow Sheet
- 7 Discharge Summary

### References to Other Databases

- NTDS 2026

## Cerebral Vascular Accident (CVA)

2026 NTDS Data Dictionary, Released July 2025



## CHRONIC OBSTRUCTIVE PULMONARY DISEASE (COPD)

---

### Element Intent

COPD limits respiratory reserve and prolongs the duration of mechanical ventilation, which could increase the risk of adverse outcomes.

### Description

Chronic obstructive pulmonary disease (COPD) is a lung disease characterized by chronic obstruction of lung airflow that interferes with normal breathing and is not fully reversible. The more familiar terms 'chronic bronchitis' and emphysema' are no longer used but are now included within the COPD diagnosis.

### EXCLUDE:

- Patients whose only pulmonary disease is asthma.
- Patients with diffuse interstitial fibrosis or sarcoidosis.

### Element Values

- 1 Yes
- 2 No

### Common Null Values

- Accepted

### Additional Information

- Present prior to injury.
- A diagnosis of COPD must be documented in the patient's medical record.
- Only report on patients ≥15-years-of-age.
- The null value "Not Applicable" must be reported for patients <15 years-of-age.
- The null value "Not Known/Not Recorded" is only reported if no past medical history is available for patients ≥15-years-of-age.
- Consistent with World Health Organization (WHO), 2019.
- Based on the patient's age on the day of arrival at your hospital.

### Data Source Hierarchy Guide


- 1 History & Physical
- 2 Physician's Notes
- 3 Progress Notes
- 4 Case Management/Social Services
- 5 Nursing Notes/Flow Sheet
- 6 Triage/Trauma Flow Sheet
- 7 Discharge Summary

### References to Other Databases

- NTDS 2026

# Chronic Obstructive Pulmonary Disease (COPD)

2026 NTDS Data Dictionary, Released July 2025



**1** Based on the patient's age on the day of arrival at your hospital.

**2** The NTDS definition is consistent with the World Health Organization (WHO), 2019.

The terms 'chronic bronchitis' and 'emphysema' are included in the COPD diagnosis.

Asthma, diffuse interstitial fibrosis, and/or sarcoidosis are excluded from the NTDS definition.

## CHRONIC RENAL FAILURE

---

### Element Intent

Chronic renal failure reflects limited renal reserve, which increases the risk of adverse outcomes.

### Description

Chronic renal failure prior to injury that was requiring periodic peritoneal dialysis, hemodialysis, hemofiltration, or hemodiafiltration.

### Element Values

- 1 Yes
- 2 No

### Common Null Values

- Accepted

### Additional Information

- Present prior to injury.
- A diagnosis of chronic renal failure must be documented in the patient's medical record.
- The null value "Not Known/Not Recorded" is only reported if no past medical history is available.

### Data Source Hierarchy Guide


- 1 History & Physical
- 2 Physician's Notes
- 3 Progress Notes
- 4 Case Management/Social Services
- 5 Nursing Notes/Flow Sheet
- 6 Triage/Trauma Flow Sheet
- 7 Discharge Summary

### References to Other Databases

- NTDS 2026

# Chronic Renal Failure

2026 NTDS Data Dictionary, Released July 2025



**1** Include patients with chronic renal failure that was present prior to injury that required renal replacement therapy but were not compliant or declined therapy.

# CIRRHOSIS

---

## Element Intent

Cirrhosis/end state liver disease reflects limited hepatic reserve, which could impact care decisions and increase the risk of adverse outcomes.

## Description

Cirrhosis is the replacement of normal liver tissue with non-living scar tissue related to other liver diseases. Must have documentation in the medical record of cirrhosis, which might also be referred to as end-stage liver disease.

## Element Values

- 1 Yes
- 2 No

## Common Null Values

- Accepted

## Additional Information

- Present prior to injury.
- A diagnosis of cirrhosis, or documentation of cirrhosis by diagnostic imaging studies or a laparotomy/laparoscopy, must be in the patient's medical record.
- Documentation in the medical record may include CHILD or MELD scores that support evidence of cirrhosis.
- The null value "Not Known/Not Recorded" is only reported if no past medical history is available.
- Patients who no longer have cirrhosis due to a successful liver transplant should be excluded.

## Data Source Hierarchy Guide


- 1 History & Physical
- 2 Physician's Notes
- 3 Progress Notes
- 4 Case Management/Social Services
- 5 Nursing Notes/Flow Sheet
- 6 Triage/Trauma Flow Sheet
- 7 Discharge Summary

## References to Other Databases

- NTDS 2026

## Cirrhosis

2026 NTDS Data Dictionary, Released July 2025



**1** Might also be referred to as end-stage liver disease.

**2** Exclude patients who no longer have cirrhosis due to a successful liver transplant.

**3** In lieu of a diagnosis of cirrhosis, documentation of cirrhosis by diagnostic imaging studies or a laparotomy/laparoscopy may be used.



## CONGENITAL ANOMALIES

---

### Element Intent

Congenital anomalies have a multitude of risks, which could impact care decisions and increase the risk of adverse outcomes.

### Description

Documentation of a cardiac, pulmonary, airway, body wall, CNS/spinal, GI, renal, orthopedic, or metabolic anomaly.

### Element Values

- 1 Yes
- 2 No

### Common Null Values

- Accepted

### Additional Information

- Present prior to injury.
- A diagnosis of a congenital anomaly must be documented in the patient's medical record.
- Only report on patients <15 years-of-age.
- The null value "Not Applicable" must be reported for patients ≥15-years-of-age.
- The null value "Not Known/Not Recorded" is only reported if no past medical history is available for patients <15 years-of-age.
- Based on the patient's age on the day of arrival at your hospital.

### Data Source Hierarchy Guide


- 1 History & Physical
- 2 Physician's Notes
- 3 Progress Notes
- 4 Case Management/Social Services
- 5 Nursing Notes/Flow Sheet
- 6 Triage/Trauma Flow Sheet
- 7 Discharge Summary

### References to Other Databases

- NTDS 2026

## Congenital Anomalies

2026 NTDS Data Dictionary, Released July 2025



**1** Based on the patient's age on the day of arrival at your hospital.

**2** Congenital anomalies are limited to those listed in the *Description*. Consult with the TPM or TMD if questioning a specific diagnosis.

## CONGESTIVE HEART FAILURE (CHF)

---

### Element Intent

CHF reflects limited cardiac reserve, leading to a higher risk of adverse outcomes

### Description

The inability of the heart to pump a sufficient quantity of blood to meet the metabolic needs of the body or can do so only at an increased ventricular filling pressure.

### Element Values

- 1 Yes
- 2 No

### Common Null Values

- Accepted

### Additional Information

- Present prior to injury.
- A diagnosis of CHF must be documented in the patient's medical record.
- To be included, this condition must be noted in the medical record as CHF, congestive heart failure, or pulmonary edema with onset of increasing symptoms within 30 days prior to injury.
- Common manifestations are:
- Abnormal limitation in exercise tolerance due to dyspnea or fatigue
- Orthopnea (dyspnea or lying supine)
- Paroxysmal nocturnal dyspnea (awakening from sleep with dyspnea)
- Increased jugular venous pressure
- Pulmonary rales on physical examination
- Cardiomegaly
- Pulmonary vascular engorgement
- The null value "Not Known/Not Recorded" is only reported if no past medical history is available.

### Data Source Hierarchy Guide


- 1 History & Physical
- 2 Physician's Notes
- 3 Progress Notes
- 4 Case Management/Social Services
- 5 Nursing Notes/Flow Sheet
- 6 Triage/Trauma Flow Sheet
- 7 Discharge Summary

### References to Other Databases

- NTDS 2026

## Congestive Heart Failure (CHF)

2026 NTDS Data Dictionary, Released July 2025



## CURRENT SMOKER

---

### Element Intent

Inhaling nicotine could induce negative cardiopulmonary effects, increase risk for stroke, negatively affect wound healing, increase anesthesia risk and the development of a venous thromboembolism (VTE), which could impact care decisions and increase the risk of adverse outcomes.

### Description

A patient who reports inhaling nicotine by smoking cigars, pipes, cigarettes, e-cigarettes, vaping, or juuling every day or some days within the last 30-days.

### EXCLUDE:

- Patients who chew tobacco or snuff.

### Element Values

- 1 Yes
- 2 No

### Common Null Values

- Accepted

### Additional Information

- Present prior to injury.
- The null value “Not Known/Not Recorded” is only reported if no past medical history is available.
- Vaping and juuling includes vape pens, dab pens, dab rings, mods, pod-mods, or any other electronic delivery system used to inhale nicotine.

### Data Source Hierarchy Guide


- 1 History & Physical
- 2 Physician's Notes
- 3 Progress Notes
- 4 Case Management/Social Services
- 5 Nursing Notes/Flow Sheet
- 6 Triage/Trauma Flow Sheet
- 7 Discharge Summary

### References to Other Databases

- NTDS 2026

## Current Smoker

2026 NTDS Data Dictionary, Released July 2025



1

Vaping and juuling includes vape pens, dab pens, dab rings, mods, pod-mods, or any other electronic delivery system used to inhale nicotine.

Exclude patients who chew tobacco or snuff.

## CURRENTLY RECEIVING CHEMOTHERAPY FOR CANCER

---

### Element Intent

The effects of chemotherapy increase the risk of infection, and could limit physiologic reserve, which together increases the risk of adverse outcomes.

### Description

A patient who is currently receiving any chemotherapy treatment for cancer prior to injury.

### Element Values

- 1 Yes
- 2 No

### Common Null Values

- Accepted

### Additional Information

- Present prior to injury.
- Chemotherapy may include, but is not restricted to, oral and parenteral treatment with chemotherapeutic agents for malignancies such as colon, breast, lung, head and neck, and gastrointestinal solid tumors as well as lymphatic and hematopoietic malignancies such as lymphoma, leukemia, and multiple myeloma.
- The null value “Not Known/Not Recorded” is only reported if no past medical history is available.

### Data Source Hierarchy Guide


- 1 History & Physical
- 2 Physician's Notes
- 3 Progress Notes
- 4 Case Management/Social Services
- 5 Nursing Notes/Flow Sheet
- 6 Triage/Trauma Flow Sheet
- 7 Discharge Summary

### References to Other Databases

- NTDS 2026

## Currently Receiving Chemotherapy for Cancer

2026 NTDS Data Dictionary, Released July 2025



1

Limited to patients who were in active chemotherapy treatment for cancer and does not include patients with a history of receiving chemotherapy for cancer.

Chemotherapy may include, but is not restricted to, oral and parenteral treatment with chemotherapeutic agents for malignancies such as colon, breast, lung, head and neck, and gastrointestinal solid tumors as well as lymphatic and hematopoietic malignancies such as lymphoma, leukemia, and multiple myeloma.



# DEMENTIA

---

## Element Intent

Patients with dementia experience forgetfulness, limited social skills and impaired thinking that could impact care decisions and prolong the length of stay.

## Description

Documentation in the patient's medical record of dementia including senile or vascular dementia (e.g., Alzheimer's).

## Element Values

- 1 Yes
- 2 No

## Common Null Values

1. Accepted

## Additional Information

- Present prior to injury.
- A diagnosis of dementia including Alzheimer's, Lewy Body Dementia, frontotemporal dementia (Pick's Disease), or vascular dementia must be documented in the patient's medical record.
- The null value "Not Known/Not Recorded" is only reported if no past medical history is available.
- Consistent with the National Institute on Aging December 2017.

## Data Source Hierarchy Guide


- 1 History & Physical
- 2 Physician's Notes
- 3 Progress Notes
- 4 Case Management/Social Services
- 5 Nursing Notes/Flow Sheet
- 6 Triage/Trauma Flow Sheet
- 7 Discharge Summary

## References to Other Databases

- NTDS 2026

# Dementia

2026 NTDS Data Dictionary, Released July 2025



Documentation of Alzheimer's, Lewy Body Dementia, frontotemporal dementia (Pick's Disease) or vascular dementia.

## DIABETES MELLITUS

---

### Element Intent

Diabetes can increase risk for infection, negatively affect wound healing, and contribute to renal and cardiac dysfunction, which could impact care decisions and increase the risk of adverse outcomes.

### Description

Diabetes mellitus that requires exogenous parenteral insulin or an oral hypoglycemic agent.

### Element Values

- 1 Yes
- 2 No

### Common Null Values

- Accepted

### Additional Information

- Present prior to injury.
- A diagnosis of diabetes mellitus must be documented in the patient's medical record.
- Report *Element Value* "1. Yes" for patients who were non-compliant with their prescribed exogenous parenteral insulin or oral hypoglycemic agent.
- The null value "Not Known/Not Recorded" is only reported if no past medical history is available.

### Data Source Hierarchy Guide


- 1 History & Physical
- 2 Physician's Notes
- 3 Progress Notes
- 4 Case Management/Social Services
- 5 Nursing Notes/Flow Sheet
- 6 Triage/Trauma Flow Sheet
- 7 Discharge Summary

### References to Other Databases

- NTDS 2026

## Diabetes Mellitus

2026 NTDS Data Dictionary, Released July 2025



**1** Include patients whose diabetes mellitus required exogenous parenteral insulin or an oral hypoglycemic agent but were non-compliant with treatment.

## DISSEMINATED CANCER

---

### Element Intent

Advanced malignancy reflecting serious physiologic compromise has a multitude of risks, which could impact care decisions and increase the risk of adverse outcomes.

### Description

Cancer that has spread to one or more sites in addition to the primary site AND in the presence of multiple metastases indicates the cancer is widespread, fulminant, or near terminal.

### Element Values

- 1 Yes
- 2 No

### Common Null Values

- Accepted

### Additional Information

- Present prior to injury.
- Another term describing disseminated cancer is "metastatic cancer."
- A diagnosis of cancer that has spread to one or more sites must be documented in the patient's medical record.
- The null value "Not Known/Not Recorded" is only reported if no past medical history is available.

### Data Source Hierarchy Guide


- 1 History & Physical
- 2 Physician's Notes
- 3 Progress Notes
- 4 Case Management/Social Services
- 5 Nursing Notes/Flow Sheet
- 6 Triage/Trauma Flow Sheet
- 7 Discharge Summary

### References to Other Databases

- NTDS 2026

## Disseminated Cancer

2026 NTDS Data Dictionary, Released July 2025



**1** Include patients who have cancer that has spread to one or more sites in addition to the primary site AND in whom the presence of multiple metastases indicates the cancer is widespread, fulminant, or near terminal. Another term describing disseminated cancer is "metastatic cancer."

## FUNCTIONALLY DEPENDENT HEALTH STATUS

---

### Element Intent

Pre-injury functional status could indicate a chronic/underlying disease state, which could impact care decisions and increase the risk of adverse outcomes.

### Description

Pre-injury functional status may be represented by the ability of the patient to complete age appropriate activities of daily living (ADL).

### Element Values

- 1 Yes
- 2 No

### Common Null Values

- Accepted

### Additional Information

- Present prior to injury.
- Activities of Daily Living include: bathing, feeding, dressing, toileting, and walking.
- Include patients whom prior to injury, and as a result of cognitive or physical limitations relating to a pre-existing medical condition, was partially dependent or completely dependent upon equipment, devices or another person to complete some or all activities of daily living.
- The null value "Not Known/Not Recorded" is only reported if no past medical history is available.
- If Ventilator Dependence is Element Value "1. Yes", Functionally Dependent Health Status must be Element Value "1. Yes."

### Data Source Hierarchy Guide


- 1 History & Physical
- 2 Physician's Notes
- 3 Progress Notes
- 4 Case Management/Social Services
- 5 Nursing Notes/Flow Sheet
- 6 Triage/Trauma Flow Sheet
- 7 Discharge Summary

### References to Other Databases

- NTDS 2026

## Functionally Dependent Health Status

2026 NTDS Data Dictionary, Released July 2025



**1** Activities of daily living include bathing, feeding, dressing, toileting, and walking. Consult with the TPM or TMD if questioning a specific pre-injury functional status.



## HYPERTENSION

---

### Element Intent

Hypertension that requires medication increases risk for cerebrovascular, renal, and cardiac disease, which could impact care decisions and increase the risk of adverse outcomes.

### Description

History of persistent elevated blood pressure requiring antihypertensive medication.

### Element Values

- 1 Yes
- 2 No

### Common Null Values

- Accepted

### Additional Information

- Present prior to injury.
- A diagnosis of Hypertension must be documented in the patient's medical record.
- The null value "Not Known/Not Recorded" is only reported if no past medical history is available.
- Report Element Value '1. Yes' for patients who were non-compliant with their prescribed antihypertensive medication.

### Data Source Hierarchy Guide


- 1 History & Physical
- 2 Physician's Notes
- 3 Progress Notes
- 4 Case Management/Social Services
- 5 Nursing Notes/Flow Sheet
- 6 Triage/Trauma Flow Sheet
- 7 Discharge Summary

### References to Other Databases

- NTDS 2026

# Hypertension

2026 NTDS Data Dictionary, Released July 2025



**1** Include patients who were non-compliant with prescribed antihypertensive medication to treat their hypertension.

## MAJOR DEPRESSIVE DISORDER

---

### Element Intent

Patients with Major Depressive Disorder experience depressed mood, loss of interest/pleasure, weight issues, fatigue, insomnia or hypersomnia, psychomotor agitation or retardation, decreased concentration, delusional guilt, and suicidal ideation which cause clinically significant distress or impairment in social, occupational, or other important areas of functioning, which could impact care decisions and increase the risk of adverse outcomes.

### Description

A major depressive disorder diagnosis documented in the medical record.

### Element Values

- 1 Yes
- 2 No

### Additional Information

1. Present prior to injury.
2. Only report on patients ≥15 years-of-age.
3. The null value "Not Applicable" must be reported for patients <15 years-of-age.
4. The null value "Not Known/Not Recorded" is only reported if no past medical history is available for patients ≥15 years-of- age.
5. Based on the patient's age on the day of arrival at your hospital.

### Data Source Hierarchy Guide


- 1 History and Physical
- 2 Physician Notes/Flow Sheet
- 3 Progress Notes
- 4 Case Management/Social Services Notes
- 5 Nursing Notes/Flow Sheet
- 6 Triage/Trauma Flow Sheet
- 7 Discharge Summary

### References to Other Databases

- NTDS 2026

## Major Depressive Disorder

2026 NTDS Data Dictionary, Released July 2025



**1** Based on the patient's age on the day of arrival at your hospital.

## MYOCARDIAL INFARCTION (MI)

---

### Element Intent

Myocardial infarction causes damage or death to the heart muscle, which could impact care decisions and increase the risk of adverse outcomes.

### Description

History of a MI in the six months prior to injury.

### Element Values

- 1 Yes
- 2 No

### Common Null Values

- Accepted

### Additional Information

- Present prior to injury.
- A diagnosis of MI must be documented in the patient's medical record.
- The null value "Not Known/Not Recorded" is only reported if no past medical history is available.

### Data Source Hierarchy Guide


- 1 History & Physical
- 2 Physician's Notes
- 3 Progress Notes
- 4 Case Management/Social Services
- 5 Nursing Notes/Flow Sheet
- 6 Triage/Trauma Flow Sheet
- 7 Discharge Summary

### References to Other Databases

- NTDS 2026

## Myocardial Infarction (MI)

2026 NTDS Data Dictionary, Released July 2025



## OTHER MENTAL/PERSONALITY DISORDERS

---

### Element Intent

Patients with these disorders experience significant distress or impairment in social, occupational, or other important areas of functioning, which could impact care decisions and increase the risk of adverse outcomes.

### Description

A diagnosis of any of the following documented in the medical record:

1. Antisocial personality disorder
2. Avoidant personality disorder
3. Borderline personality disorder
4. Dependent personality disorder
5. Generalized anxiety disorder
6. Histrionic personality disorder
7. Narcissistic personality disorder
8. Obsessive-compulsive disorder
9. Obsessive-compulsive personality disorder
10. Panic disorder
11. Paranoid personality disorder
12. Schizotypal personality disorder

### Element Values

- 1 Yes
- 2 No

### Additional Information

- Present prior to injury.
- Only report on patients ≥15 years-of-age.
- The null value "Not Applicable" must be reported for patients <15 years-of-age.
- The null value "Not Known/Not Recorded" is only reported if no past medical history is available for patients ≥15 years-of- age.
- Based on the patient's age on the day of arrival at your hospital.

### Data Source Hierarchy Guide


- 1 History and Physical
- 2 Physician Notes/Flow Sheet
- 3 Progress Notes
- 4 Case Management/Social Services Notes
- 5 Nursing Notes/Flow Sheet
- 6 Triage/Trauma Flow Sheet
- 7 Discharge Summary

### References to Other Databases

- NTDS 2026

## Other Mental/Personality Disorders

2026 NTDS Data Dictionary, Released July 2025



**1** Based on the patient's age on the day of arrival at your hospital.



## PERIPHERAL ARTERIAL DISEASE (PAD)

---

### Element Intent

PAD reflects cardiovascular risk, which itself is associated with adverse outcomes.

### Description

The narrowing or blockage of the vessels that carry blood from the heart to the legs. It is primarily caused by the buildup of fatty plaque in the arteries, which is called atherosclerosis. PAD can occur in any blood vessel, but it is more common in the legs than the arms.

### Element Values

- 1 Yes
- 2 No

### Common Null Values

- Accepted

### Additional Information

- Present prior to injury.
- A diagnosis of Peripheral Arterial Disease or Peripheral Vascular Disease must be documented in the patient's medical record.
- Only report on patients  $\geq 15$  years-of-age.
- The null value "Not Applicable" must be reported for patients  $< 15$  years-of-age.
- The null value "Not Known/Not Recorded" is only reported if no past medical history is available for patients  $\geq 15$  years-of-age.
- Consistent with Centers for Disease Control, 2014 Fact Sheet.
- Based on the patient's age on the day of arrival at your hospital.

### Data Source Hierarchy Guide


- 1 History & Physical
- 2 Physician's Notes
- 3 Progress Notes
- 4 Case Management/Social Services
- 5 Nursing Notes/Flow Sheet
- 6 Triage/Trauma Flow Sheet
- 7 Discharge Summary

### References to Other Databases

- NTDS 2026

## Peripheral Arterial Disease (PAD)

2026 NTDS Data Dictionary, Released July 2025



1

Based on the patient's age on the day of arrival at your hospital.

2

Consistent with the CDC 2014 Fact Sheet. The term "peripheral vascular disease (PVD)" can be used interchangeably with "PAD."

## POST-TRAUMATIC STRESS DISORDER

---

### Element Intent

Patients with PTSD experience intrusive symptoms, avoidance, altered mood, altered reactivity, causing clinically significant distress or impairment in social, occupational, or other important areas of functioning, which could impact care decisions and increase the risk of adverse outcomes.

### Description

A post-traumatic stress disorder diagnosis documented in the medical record.

### Element Values

- 1 Yes
- 2 No

### Additional Information

- Present prior to injury.
- Only report on patients  $\geq 15$  years-of-age.
- The null value "Not Applicable" must be reported for patients  $< 15$  years-of-age.
- The null value "Not Known/Not Recorded" is only reported if no past medical history is available for patients  $\geq 15$  years-of-age.
- Based on the patient's age on the day of arrival at your hospital.

### Data Source Hierarchy Guide


- 1 History and Physical
- 2 Physician Notes/Flow Sheet
- 3 Progress Notes
- 4 Case Management/Social Services Notes
- 5 Nursing Notes/Flow Sheet
- 6 Triage/Trauma Flow Sheet
- 7 Discharge Summary

### References to Other Databases

- NTDS 2026

## Post-Traumatic Stress Disorder (PTSD)

2026 NTDS Data Dictionary, Released July 2025



Based on the patient's age on the day of arrival at your hospital.

## PREGNANCY

---

### Element Intent

Trauma during pregnancy could cause pre-term labor and/or placental abruption, which could impact care decisions and increase the risk of adverse outcomes.

### Description

Pregnancy confirmed by lab, ultrasound, or other diagnostic tool OR diagnosis of pregnancy documented in the patient's medical record.

### Element Values

- 1 Yes
- 2 No

### Additional Information

- Present prior to arrival at your center
- The null value "Not Known/Not Recorded" is only reported if no past medical history is available.

### Data Source Hierarchy Guide


- 1 History & Physical
- 2 Physician's Notes
- 3 Progress Notes
- 4 Case Management/Social Services
- 5 Nursing Notes/Flow Sheet
- 6 Triage/Trauma Flow Sheet
- 7 Discharge Summary

### References to Other Databases

- NTDS 2026

# Pregnancy

2026 NTDS Data Dictionary, Released July 2025



**1** Pregnancy must be confirmed by lab, ultrasound, or other diagnostic tool if diagnosis was not documented.  
This data element must be reported for all records submitted to TQIP, males and females.

## PREMATURITY

---

### Element Intent

Prematurity can induce a multitude of risks, which could impact care decisions and increase the risk of adverse outcomes.

### Description

Babies born before 37 weeks of pregnancy are completed.

### Element Values

- 1 Yes
- 2 No

### Common Null Values

- Accepted

### Additional Information

- Present prior to injury.
- Only report on patients <15 years-of-age.
- A diagnosis of prematurity, or delivery before 37 weeks of pregnancy are completed, must be documented in the patient's medical record.
- The null value "Not Applicable" must be reported for patients ≥15-years-of-age.
- The null value "Not Known/Not Recorded" is only reported if no past medical history is available for patients <15 years-of-age.
- Based on the patient's age on the day of arrival at your hospital.

### Data Source Hierarchy Guide


- 1 History & Physical
- 2 Physician's Notes
- 3 Progress Notes
- 4 Case Management/Social Services
- 5 Nursing Notes/Flow Sheet
- 6 Triage/Trauma Flow Sheet
- 7 Discharge Summary

### References to Other Databases

- NTDS 2026

# Prematurity

2026 NTDS Data Dictionary, Released July 2025



Based on the patient's age on the day of arrival at your hospital.



## SCHIZOAFFECTIVE DISORDER

---

### Element Intent

Patients with Schizoaffective Disorder experience hallucinations, delusions, mania, depression, and disorganized thinking causing clinically significant distress or impairment in social, occupational, or other important areas of functioning, which could impact care decisions and increase the risk of adverse outcomes.

### Description

A schizoaffective disorder diagnosis documented in the medical record.

### Element Values

- 1 Yes
- 2 No

### Additional Information

- Present prior to injury.
- Only report on patients  $\geq 15$  years-of-age.
- The null value "Not Applicable" must be reported for patients  $< 15$  years-of-age.
- The null value "Not Known/Not Recorded" is only reported if no past medical history is available for patients  $\geq 15$  years-of-age.
- Based on the patient's age on the day of arrival at your hospital.

### Data Source Hierarchy Guide


- 1 History and Physical
- 2 Physician Notes/Flow Sheet
- 3 Progress Notes
- 4 Case Management/Social Services Notes
- 5 Nursing Notes/Flow Sheet
- 6 Triage/Trauma Flow Sheet
- 7 Discharge Summary

### References to Other Databases

- NTDS 2026

## Schizoaffective Disorder

2026 NTDS Data Dictionary, Released July 2025



Based on the patient's age on the day of arrival at your hospital.

# SCHIZOPHRENIA

---

## Element Intent

Patients with Schizophrenia experience hallucinations, delusions, disorganized speech, grossly disorganized or catatonic behavior, diminished emotional expression or avolition causing clinically significant distress or impairment in social, occupation, or other important areas of functioning, which could impact care decisions and increase the risk of adverse outcomes.

## Description

A schizophrenia diagnosis documented in the medical record.

## Element Values

- 1 Yes
- 2 No

## Additional Information

- Present prior to injury.
- Only report on patients ≥15 years-of-age.
- The null value "Not Applicable" must be reported for patients <15 years-of-age.
- The null value "Not Known/Not Recorded" is only reported if no past medical history is available for patients ≥15 years-of- age.
- Based on the patient's age on the day of arrival at your hospital.

## Data Source Hierarchy Guide


- 1 History and Physical
- 2 Physician Notes/Flow Sheet
- 3 Progress Notes
- 4 Case Management/Social Services Notes
- 5 Nursing Notes/Flow Sheet
- 6 Triage/Trauma Flow Sheet
- 7 Discharge Summary

## References to Other Databases

- NTDS 2026

# Schizophrenia

2026 NTDS Data Dictionary, Released July 2025



Based on the patient's age on the day of arrival at your hospital.

## STEROID USE

---

### Element Intent

Steroids negatively affect wound healing and increase the risk of infection, which could impact care decisions and increase the risk of adverse outcomes.

### Description

Regular administration of oral or parenteral corticosteroid medications within 30 days prior to injury for a chronic medical condition.

### EXCLUDE:

- Topical corticosteroids applied to the skin, and corticosteroids administered by inhalation or rectally.

### Element Values

- 1 Yes
- 2 No

### Common Null Values

- Accepted

### Additional Information

- Present prior to injury.
- Examples of oral or parenteral corticosteroid medications are: prednisone and dexamethasone.
- Examples of chronic medical conditions are: COPD, asthma, rheumatologic disease, rheumatoid arthritis, and inflammatory bowel disease.
- The null value "Not Known/Not Recorded" is only reported if no past medical history is available.

### Data Source Hierarchy Guide


- 1 History & Physical
- 2 Physician's Notes
- 3 Progress Notes
- 4 Case Management/Social Services
- 5 Nursing Notes/Flow Sheet
- 6 Triage/Trauma Flow Sheet
- 7 Discharge Summary

### References to Other Databases

- NTDS 2026

## Steroid Use

2026 NTDS Data Dictionary, Released July 2025



**1** A few examples of chronic medical conditions that could require corticosteroids are COPD, asthma, rheumatologic disease, rheumatoid arthritis, and inflammatory bowel disease.

**2** Exclude corticosteroids that are administered by inhalation, topically, or rectally.

Examples of oral or parenteral steroid medications are prednisone and dexamethasone.

## SUBSTANCE USE DISORDER

---

### Element Intent

Patients with substance use disorder are at increased risk of heart, lung, liver, and kidney diseases, as well as stroke, cancer, and mental health conditions, which could impact care decisions and increase the risk of adverse outcomes.

### Description

Descriptors documented in the patient's medical record consistent with the diagnostic criteria of substance use disorders specifically cannabis, hallucinogens, inhalants, opioids, sedative/hypnotics, and stimulants (e.g. patient has a history of drug use; patient has a history of opioid use) OR diagnosis of any of the following documented in the patient's medical record:

1. Cannabis Use Disorder; Other Cannabis-Induced Disorder; Unspecified Cannabis-Related Disorder
2. Phencyclidine Use Disorder; Other Hallucinogen Use Disorder; Hallucinogen Persisting Perception Disorder; Other Phencyclidine-Induced Disorder; Other Hallucinogen-Induced Disorder; Unspecified Phencyclidine-Related Disorder; Unspecified Hallucinogen-Related Disorder
3. Inhalant Use Disorder; Other Inhalant-Induced Disorder; Unspecified Inhalant-Related Disorder
4. Opioid Use Disorder; Other Opioid-Induced Disorder; Unspecified Opioid-Related Disorder
5. Sedative, Hypnotic, or Anxiolytic Use Disorder; Other Sedative, Hypnotic, or Anxiolytic-Induced Disorder; Unspecified Sedative, Hypnotic, or Anxiolytic-Related Disorder
6. Stimulant Use Disorder; Other Stimulant-Induced Disorder; Other Stimulant-Related Disorder
7. Based on the patient's age on the day of arrival at your hospital.

### Element Values

- 1 Yes
- 2 No

### Common Null Values

- Accepted

### Additional Information

- Present prior to injury.
- Only report on patients ≥15-years-of-age.
- The null value "Not Applicable" must be reported for patients <15 years-of-age.
- The null value "Not Known/Not Recorded" is only reported if no past medical history is available for patients ≥15-years-of-age.
- Consistent with the American Psychiatric Association (APA) DSM 5, 2013.

### Data Source Hierarchy Guide


- 1 History & Physical
- 2 Physician's Notes
- 3 Progress Notes
- 4 Case Management/Social Services
- 5 Nursing Notes/Flow Sheet
- 6 Triage/Trauma Flow Sheet
- 7 Discharge Summary

### References to Other Databases

- NTDS 2026

## Substance Use Disorder

2026 NTDS Data Dictionary, Released July 2025





--Page Blank--

## VENTILATOR DEPENDENCE

---

### Element Intent

The need for ventilator-assisted respirations reflects limited pulmonary reserve, which increases the risk of adverse outcomes.

### Description

Patients who are ventilator dependent with a tracheostomy prior to injury.

### Element Values

1. Yes
2. No

### Additional Information

- Present prior to injury.
- The null value "Not Known/Not Recorded" is only reported if no past medical history is available.
- If Ventilator Dependence is Element Value "1. Yes", Functionally Dependent Health Status must be Element Value "1. Yes."

### Data Source Hierarchy Guide


1. History and Physical
2. Physician Notes/Flow Sheet
3. Progress Notes
4. Case Management/Social Services Notes
5. Nursing Notes/Flow Sheet
6. Triage/Trauma Flow Sheet
7. Discharge Summary

### References to Other Databases

- NTDS 2026

## Ventilator Dependence

2026 NTDS Data Dictionary, Released July 2025



**1** If Ventilator Dependence is Element Value "1. Yes," Functionally Dependent Health Status must be Element Value "1. Yes."

## DNR STATUS

---

### Element Intent

Provides information about patient's care in the hospital

### Description

DNR Status documents the presence of signed DNR paperwork and/or written order from physician or mid-level provider to withhold select resuscitative efforts from the patient, and whether the order was issued prior to or during the patient's stay at your ED/hospital.

### Element Values

1. Not a DNR patient (patient is to receive all resuscitative efforts if needed)
2. DNR status ordered prior to patient's arrival at your hospital
3. DNR status ordered after patient's arrival to your hospital

### Common Null Values

- Not Accepted

### Additional Information

- This element is completed for each patient.
- DNR status is typically ordered for a patient who does not wish to be resuscitated in the event of a cardiac arrest (no palpable pulse) or respiratory arrest (no spontaneous respirations or the presence of labored breathing) near the end of life.
- A DNR status includes both *DNR-CC* (comfort care) and *DNR-CCA* (comfort care arrest) orders.
- DNR may also be referred to as Allow Natural Death (AND)
- Until DNR status is documented, the patient is considered to be "not a DNR patient".
- DNR Status is to be collected at time of discharge from Hospital.
- Refer to Ohio Department of Health for additional details: <https://odh.ohio.gov/wps/portal/gov/odh/know-our-programs/do-not-resuscitate-comfort-care>.

### Data Source Hierarchy Guide

- 1 Do Not Resuscitate Document
- 2 History and Physical
- 3 Discharge Sheet
- 4 Billing Sheet

### References to Other Databases

- Not an NTDS element

## ICD-10 INJURY DIAGNOSES

---

### Element Intent

To classify and quantify the severity of individual injuries, which is used to understand injury patterns, care plans, and outcomes.

### Description

*Injury Diagnoses* related to all identified injuries.

### Element Values

1. Injury diagnoses are defined by ICD-10-CM codes; refer to inclusion criteria
2. The maximum number of diagnoses that may be reported for an individual patient is 50.

### Common Null Values

- Not Accepted

### Additional Information

- ICD-10-CM codes pertaining to other medical conditions (e.g., CVA, MI, co-morbidities, etc.) may also be included in this element

### Data Source Hierarchy Guide

- 1 Autopsy/Medical Examiner Report
- 2 Operative Reports
- 3 Radiology Reports
- 4 Physician's Notes
- 5 Trauma Flow Sheet
- 6 History & Physical
- 7 Nursing Notes/Flow Sheet
- 8 Progress Notes
- 9 Discharge Summary

### References to Other Databases

- NTDS 2026

## AIS CODE

---

### Element Intent

To classify and quantify the severity of individual injuries, which is used to understand injury patterns, care plans, and outcomes.

### Description

The Abbreviated Injury Scale (AIS) code(s) that reflect the patient's injuries.

### Element Values

1. The code is the 8-digit AIS code

### Additional Information

None

### Data Source Hierarchy Guide

- 1 AIS Coding Manual

### References to Other Databases

- NTDS 2026

## AIS VERSION

---

### Element Intent

To indicate the AIS version used to code identified injuries.

### Description

*AIS version* is the software version used to calculate Abbreviated Injury Scale (AIS) severity codes for the patient's current injury event.

### Element Values

1. AIS 2015

### Additional Information

None

### Common Null Values

- Accepted

### Data Source Hierarchy Guide

- 1 AIS Coding Manual

### References to Other Databases

- NTDS 2026

# INJURY SEVERITY SCORE

---

**Element Intent**

To classify and quantify the severity of individual injuries, which is used to understand injury patterns, care plans, and outcomes.

**Description**

*Injury Severity Score (ISS)* is a nationally-accepted scoring system that reflects the patient’s injuries for this injury event.

**Element Values**

- 1. Relevant ISS value for the constellation of injuries

**Additional Information**

None

**Common Null Values**

- Accepted

**Data Source Hierarchy Guide**

- 1 AIS Coding Manual

**References to Other Databases**

- Not an NTDS element



## TOTAL ICU LENGTH OF STAY

---

### Element Intent

To indicate the patient's condition and the duration of their requirement for specialized monitoring and care.

### Description

The cumulative amount of time spent in the ICU. Each partial or full day should be measured as one calendar day.

### Element Values

1. Relevant value for data element

### Common Null Values

- Accepted

### Additional Information

- Reported in full day increments with any partial calendar day counted as a full calendar day.
- The calculation assumes that the date and time of starting and stopping an ICU episode are recorded in the patient's chart.
- The null value "Not Known / Not Recorded" is reported if any dates are missing.
- If patient has multiple ICU episodes on the same calendar day, count that day as one calendar day.
- At no time should the ICU LOS exceed the Hospital LOS.
- The null value "Not applicable" is reported if the patient had no ICU days according to the above definition.
- A '0' (zero) in this field is not an acceptable value.
- See Appendix B for examples of ICU LOS calculations

### Data Source Hierarchy Guide

- 1 ICU Flow Sheet
- 2 Nursing Notes/Flow Sheet

### References to Other Databases

- NTDS 2026

## TOTAL VENTILATOR DAYS

---

### Element Intent

To indicate the patient's respiratory condition and the duration of their requirement for specialized monitoring and care.

### Description

The cumulative amount of time spent on the ventilator. Each partial or full day should be measured as one calendar day.

### Element Values

1. Relevant value for data element

### Common Null Values

- Accepted

### Additional Information

- Excludes mechanical ventilation time associated with OR procedures.
- Non-invasive means of ventilatory support (CPAP or BIPAP) should not be considered in the calculation of ventilator days.
- Reported in full day increments with any partial calendar day counted as a full calendar day.
- The calculation assumes that the date and time of starting and stopping Ventilator episode are recorded in the patient's chart.
- The null value "Not known / Not Recorded" is reported if any dates are missing.
- At no time should the Total Vent Days exceed the Hospital LOS.
- The null value "Not Applicable" is reported if the patient was not on the ventilator according to the above definition.
- A '0' (zero) in this field is not an acceptable value.
- See Appendix B for examples of Total Ventilator Days calculations.

### Data Source Hierarchy Guide

- 1 Respiratory Therapy Notes/Flow Sheet
- 2 ICU Flow Sheet
- 3 Progress Notes

### References to Other Databases

- NTDS 2026

## HOSPITAL DISCHARGE ORDER WRITTEN DATE

---

### Element Intent

Provides information about the patient's hospital disposition, which helps track the level of care the patient received.

### Description

*Hospital Discharge Order Written Date* is the date that the order was written for the patient to be discharged from your hospital.

### Element Values

1. Relevant value for data element

### Common Null Values

- Accepted

### Additional Information

- Collected as YYYY-MM-DD
- The null value "Not Applicable" is reported if ED Discharge Disposition = 4, 6, 9, 10, or 11
- If Hospital Discharge Disposition is "5. Deceased/Expired," then Hospital Discharge Date is the date of death as indicated on the patient's death certificate
- The null value "Not Applicable" is reported if ED Discharge Disposition is 5. Deceased/Expired.
- The null value "Not Applicable" is reported if Hospital Discharge Disposition is reported as "Not Applicable"

### Data Source Hierarchy Guide

- 1 Physician Order
- 2 Discharge Instructions
- 3 Nursing Notes/Flow Sheet
- 4 Case Management/Social Services Notes
- 5 Discharge Summary

### References to Other Databases

- Not an NTDS element

**NOTE:** HOSPITAL DISCHARGE ORDER WRITTEN DATE differs from NTDS. Please refer to the section "DIFFERENCES BETWEEN OHIO AND NATIONAL TRAUMA DATA STANDARD (NTDS)" starting on Page 12 for more information.

## HOSPITAL DISCHARGE ORDER WRITTEN TIME

---

### Element Intent

Provides information about the patient's hospital disposition, which helps track the level of care the patient received.

### Description

*Hospital Discharge Order Written Time* is the time that the order was written for the patient to be discharged from your hospital.

### Element Values

1. Relevant value for data element

### Common Null Values

- Accepted

### Additional Information

- Collected as HHMM military time
- The null value "Not Applicable" is used if ED Discharge Disposition = 4, 6, 9, 10, or 11.
- If Hospital Discharge Disposition is "5. Deceased/Expired," then Hospital Discharge Date is the date of death as indicated on the patient's death certificate
- The null value "Not Applicable" is used if ED Discharge Disposition = 5 (Deceased/ expired).
- The null value "Not Applicable" is reported if Hospital Discharge Disposition is reported as "Not Applicable"

### Data Source Hierarchy Guide

- 1 Physician Order
- 2 Discharge Instructions
- 3 Nursing Notes/Flow Sheet
- 4 Case Management/Social Services Notes
- 5 Discharge Summary

### References to Other Databases

- Not an NTDS element

**NOTE:** HOSPITAL DISCHARGE ORDER WRITTEN TIME differs from NTDS. Please refer to the section "DIFFERENCES BETWEEN OHIO AND NATIONAL TRAUMA DATA STANDARD (NTDS)" starting on Page 12 for more information.

## HOSPITAL DISCHARGE DATE

---

### Element Intent

To calculate metrics such as hospital length of stay and to inform the care timeline.

### Description

*Hospital Discharge Date* is the date that the patient was physically discharged from your hospital.

### Element Values

1. Relevant value for data element

### Common Null Values

- Accepted

### Additional Information

- Collected as YYYY-MM-DD
- The null value “Not Applicable” is reported if ED Discharge Disposition = 4, 6, 9, 10, or 11
- The null value “Not Applicable” is reported if Hospital Discharge Disposition is reported as “Not Applicable”
- If Hospital Discharge Disposition is “5. Deceased/Expired,” then Hospital Discharge Date is the date of death as indicated on the patient’s death certificate

### Data Source Hierarchy Guide

1. Discharge Instructions
2. Nursing Notes/Flow Sheet
3. Case Management/Social Services Notes
4. Discharge Summary

### References to Other Databases

- NTDS 2026 (element name only)

**NOTE:** HOSPITAL DISCHARGE DATE differs from NTDS. Please refer to the section “DIFFERENCES BETWEEN OHIO AND NATIONAL TRAUMA DATA STANDARD (NTDS)” starting on Page 12 for more information.

## HOSPITAL DISCHARGE TIME

---

### Element Intent

To calculate metrics such as hospital length of stay and to inform the care timeline.

### Description

*Hospital Discharge Time* is the time of day that the patient was physically discharged from your hospital.

### Element Values

1. Relevant value for data element

### Common Null Values

- Accepted

### Additional Information

- Collected as HHMM military time
- The null value “Not Applicable” is used if ED Discharge Disposition = 4, 6, 9, 10, or 11.
- The null value “Not Applicable” is reported if Hospital Discharge Date is reported as “Not Applicable”.
- If Hospital Discharge Disposition is “5. Deceased/Expired,” then Hospital Discharge Date is the date of death as indicated on the patient’s death certificate

### Data Source Hierarchy Guide

- 1 Discharge Instructions
- 2 Nursing Notes/Flow Sheet
- 3 Case Management/Social Services Notes
- 4 Discharge Summary

### References to Other Databases

- NTDS 2026 (element name only)

**NOTE:** HOSPITAL DISCHARGE TIME differs from NTDS. Please refer to the section “DIFFERENCES BETWEEN OHIO AND NATIONAL TRAUMA DATA STANDARD (NTDS)” starting on Page 12 for more information.

## HOSPITAL DISCHARGE DISPOSITION

---

### Element Intent

To indicate the patient's medical and support needs after their acute care is complete, including functional dependence and ongoing medical requirements.

### Description

*Hospital Discharge Disposition* documents in general terms where the patient went after discharge from your hospital.

### Element Values

- 1 Discharged/Transferred to another hospital for ongoing acute inpatient care
- 2 Discharged to an intermediate care facility (ICF)/long term care facility (LTCF)
- 3 Discharged/Transferred to home under the care of an organized home health service
- 4 Left against medical advice (AMA) or discontinued care
- 5 Died
- 6 Discharged home or self-care (routine discharge)
- 7 Discharged to a skilled nursing facility (SNF)
- 8 Discharged to hospice care
- 9 [Value 9 not used]
- 10 Discharged to court/law enforcement/jail
- 11 Discharged to another type of inpatient rehabilitation facility (IRF)
- 12 Discharged to a long term acute care hospital (LTACH)
- 13 Discharged/transferred to psychiatric hospital/psychiatric unit
- 14 Discharged/transferred to other type of institution not listed here

### Common Null Values

- Accepted

### Additional Information

- Element value "6. Home" refers to the patient's current place of residence (e.g., Prison, Child Protective Services etc.).
- Element values adapted from UB-04 disposition coding.
- Disposition to any other non-medical facility should be coded as 6.
- Disposition to any other medical facility should be coded as 14.
- The null value "Not Applicable" is reported if ED Discharge Disposition = 4, 6, 9, 10, or 11.
- The null value "Not Applicable" is reported if ED Discharge Disposition is "5, Deceased/Expired."
- Hospital Discharge Dispositions which were retired greater than 2 years before the current NTDS version are no longer listed under Element Values above, which is why there are numbering gaps.
- Refer to the NTDS Change Log for a full list of retired Hospital Discharge Dispositions.
- Disposition to any Federal Health Care facility must be reported by selecting the option that most closely aligns to the needs of the patient (e.g., patients discharged to a Veteran's hospital skilled nursing facility must be reported as Element Value "7 Discharged/Transferred to Skilled Nursing Facility.")

### Data Source Hierarchy Guide

- 1 Physician Order
- 2 Discharge Instructions
- 3 Nursing Notes/Flow Sheet
- 4 Case Management/Social Services Notes
- 5 Discharge Summary

### References to Other Databases

- NTDS 2026

## INPATIENT TRANSFER TO HOSPITAL

---

### Element Intent

Provides information about the patient's hospital disposition, which helps track the level of care the patient received.

### Description

*Inpatient Transfer to Hospital* documents a subsequent hospital destination for the patient after inpatient admission at your hospital. This includes transfers to inpatient rehabilitation facilities.

### Element Values

1. Four-digit hospital code assigned by the Ohio Department of Public Safety.

### Common Null Values

- Accepted

### Additional Information

None

### Data Source Hierarchy Guide

- 1 Discharge Summary
- 2 Progress Notes
- 3 Billing/Registration Sheet

### References to Other Databases

- Not an NTDS element



## DISCHARGE STATUS

---

### Element Intent

Provides information about patient's vital status at the time of discharge

### Description

*Discharge Status* is whether the patient left your hospital alive or dead.

### Element Values

- 1 Alive
- 2 Dead

### Common Null Values

- Not Accepted

### Additional Information

None

### Data Source Hierarchy Guide

- 1 Discharge Summary
- 2 Progress Notes
- 3 Billing Sheet

### References to Other Databases

- Not an NTDS element

## DATE OF DEATH

---

### Element Intent

If the patient is deceased, provides additional information regarding the patient's disposition.

### Description

*Date of Death* is the date that the patient was pronounced dead or time of declaration of brain death.

### Element Values

1. Relevant value for data element

### Common Null Values

- Accepted

### Additional Information

- Collected as YYYY-MM-DD
- *Date of Death* must be  $\leq$  *Hospital Discharge Date*
- Only complete element when *Discharge Status* is completed as *Dead*
- This may differ from the date of discharge

### Data Source Hierarchy Guide

- 1 Hospital Record
- 2 Billing Sheet/Medical Records Coding Summary Sheet
- 3 Physician Discharge Summary

### References to Other Databases

- Not an NTDS element

## PRIMARY METHOD OF PAYMENT

---

### Element Intent

to investigate disparities in care, access to services, and other relevant healthcare issues.

### Description

*Primary Method of Payment* is the primary source of payment for hospital care.

### Element Values

- 1 Medicaid
- 2 Not Billed (for any reason)
- 3 Self-Pay
- 4 Private/Commercial Insurance
- 6 Medicare
- 7 Other Government Payer Source
- 8 Workers Compensation
- 10 Other

### Common Null Values

- Accepted

### Additional Information

- No Fault Automobile, Workers Compensation, and Blue Cross/Blue Shield should be reported as “4. Private/Commercial Insurance”.
- Primary methods of payments which were retired greater than 2 years before the current NTDS version are no longer listed under Element Values. Refer to the NTDS Change Log for a full list of retired Primary Methods of Payments.
- Examples of “Other Government Payer Source”: Veterans Affairs (VA), TRICARE, CHAMPVA
- Charity or HCAP should be coded under “Not Billed”

### Data Source Hierarchy Guide

- 1 Billing Sheet
- 2 Admission Form
- 3 Face Sheet

### References to Other Databases

- NTDS 2026

**NOTE:** PRIMARY METHOD OF PAYMENT differs from NTDS. Please refer to the section “DIFFERENCES BETWEEN OHIO AND NATIONAL TRAUMA DATA STANDARD (NTDS)” starting on Page 12 for more information.

## AUTOPSY PERFORMED

---

### Element Intent

If the patient is deceased, provides additional information regarding the patient's disposition.

### Description

*Autopsy Performed* documents whether an internal organ exam was performed on the patient by a trained pathologist.

### Element Values

- 1 Yes, an autopsy was performed
- 2 No, an autopsy was not performed

### Common Null Values

- Accepted

### Additional Information

- Select *NA* if the patient is alive
- If only an external or visual-type exam was done and no internal organs were surgically explored, element value #2, *No, an autopsy was not performed*, should be selected.

### Data Source Hierarchy Guide

- 1 Autopsy Report
- 2 Discharge Summary

### References to Other Databases

- Not an NTDS element

## ACUTE KIDNEY INJURY (AKI)

---

### Element Intent

A potentially preventable event often induced by sepsis, hypotension, drug toxicity and/or renal trauma; advancement to stage 3 requires treatment which could increase the hospital stay and the likelihood of mortality.

### Description

Acute kidney injury, AKI (stage 3), is an abrupt decrease in kidney function that occurred during the patient's stay at your hospital.

### EXCLUDE:

1. Patients with renal failure that were requiring chronic renal replacement therapy such as periodic peritoneal dialysis, hemodialysis, hemofiltration, or hemodiafiltration prior to injury.

### KDIGO (Stage 3) Table:

(SCr) 3 times baseline

**OR**

Increase in SCr to  $\geq 4.0$  mg/dl ( $\geq 353.6$   $\mu$ mol/l)

**OR**

Initiation of renal replacement therapy OR, in patients < 18 years, decrease in eGFR to <35 ml/min per 1.73 m<sup>2</sup>

**OR**

Urine output <0.3 ml/kg/h for  $\geq 24$  hours

**OR**

Anuria for  $\geq 12$  hours

### Element Values

- 1 Yes
- 2 No

### Additional Information

- Onset of AKI stage 3 began after arrival to your ED/hospital.
- A diagnosis of AKI must be documented in the patient's medical record.
- Consistent with the March 2012 Kidney Disease Improving Global Outcome (KDIGO) Guideline.

### Data Source Hierarchy Guide


- 1 History and Physical
- 2 Physician's Notes
- 3 Progress Notes
- 4 Case Management/Social Services
- 5 Nursing Notes/Flow Sheet
- 6 Triage/Trauma Flow Sheet
- 7 Discharge Summary

### References to Other Databases

- NTDS 2026

# Acute Kidney Injury (AKI)

2026 NTDS Data Dictionary, Released July 2025



## ACUTE RESPIRATORY DISTRESS SYNDROME (ARDS)

---

### Element Intent

A potentially preventable event often induced by pneumonia, viral infection, sepsis, blood transfusion, pancreatitis, fat emboli, trauma, or other injuries, requiring treatment which could increase the hospital length of stay and likelihood of mortality.

### Description

Timing:	Within 1 week of known clinical insult or new or worsening respiratory symptoms.
Chest imaging:	Bilateral opacities – not fully explained by effusions, lobar/lung collapse, or nodules
Origin of edema:	Respiratory failure not fully explained by cardiac failure or fluid overload. Need objective assessment (e.g., echocardiography) to exclude hydrostatic edema if no risk factor present.
Oxygenation:	
Mild	200 mm Hg < PaO <sub>2</sub> /FIO <sub>2</sub> < 300 mm Hg With PEEP or CPAP ≥ 5 cm H <sub>2</sub> O
Moderate	100 mm Hg < PaO <sub>2</sub> /FIO <sub>2</sub> < 200 mm Hg With PEEP > 5 cm H <sub>2</sub> O
Severe	PaO <sub>2</sub> /FIO <sub>2</sub> < 100 mm Hg With PEEP or CPAP > 5 cm H <sub>2</sub> O

### Element Values

- 1 Yes
- 2 No

### Additional Information

- Onset of symptoms began after arrival to your ED/hospital.
- A diagnosis of ARDS must be documented in the patient's medical record.
- Consistent with the 2012 New Berlin Definition.

### Data Source Hierarchy Guide


- 1 History and Physical
- 2 Physician's Notes
- 3 Progress Notes
- 4 Case Management/Social Services
- 5 Nursing Notes/Flow Sheet
- 6 Triage/Trauma Flow Sheet
- 7 Discharge Summary

### References to Other Databases

- NTDS 2026

# Acute Respiratory Distress Syndrome (ARDS)

2026 NTDS Data Dictionary, Released July 2025





--Page Blank--

## ALCOHOL WITHDRAWAL SYNDROME

---

### Element Intent

A potentially preventable event often associated with infectious complications, requiring treatment which could increase the hospital length of stay and the likelihood of mortality.

### Description

Characterized by tremor, sweating, anxiety, agitation, depression, nausea, and malaise. It occurs 6-48 hours after cessation of alcohol consumption and, when uncomplicated, abates after 2-5 days. It may be complicated by grand mal seizures and may progress to delirium (known as delirium tremens).

### Element Values

- 1 Yes
- 2 No

### Additional Information

- Onset of symptoms began after arrival to your ED/hospital.
- Documentation of alcohol withdrawal must be in the patient's medical record.
- Consistent with the 2019 World Health Organization (WHO) definition of Alcohol Withdrawal Syndrome.

### Data Source Hierarchy Guide


- 1 History and Physical
- 2 Physician's Notes
- 3 Progress Notes
- 4 Case Management/Social Services
- 5 Nursing Notes/Flow Sheet
- 6 Triage/Trauma Flow Sheet
- 7 Discharge Summary

### References to Other Databases

- NTDS 2026

# Alcohol Withdrawal Syndrome

2026 NTDS Data Dictionary, Released July 2025



**1** Note that a diagnosis of alcohol withdrawal syndrome is not required by the NTDS definition.

## CARDIAC ARREST WITH CPR

---

### Element Intent

A potentially preventable event often associated with either a medical or trauma-related condition, requiring treatment which could increase the hospital length of stay and the likelihood of mortality.

### Description

Cardiac arrest is the sudden cessation of cardiac activity after hospital arrival. The patient becomes unresponsive with no normal breathing and no signs of circulation. If corrective measures are not taken rapidly, this condition progresses to sudden death.

### INCLUDE:

- Patients who, after arrival at your hospital, have had an episode of cardiac arrest evaluated by hospital personnel, and received compressions or defibrillation or cardioversion or cardiac pacing to restore circulation.

### EXCLUDE:

- Patients whose ONLY episode of cardiac arrest with CPR was on arrival to your hospital.

### Element Values

- 1 Yes
- 2 No

### Additional Information

- Onset of symptoms began after arrival to your ED/hospital.
- Cardiac Arrest must be documented in the patient's medical record.

### Data Source Hierarchy Guide


- 1 History and Physical
- 2 Physician's Notes
- 3 Progress Notes
- 4 Case Management/Social Services
- 5 Nursing Notes/Flow Sheet
- 6 Triage/Trauma Flow Sheet
- 7 Discharge Summary

### References to Other Databases

- NTDS 2026

## Cardiac Arrest with CPR

2026 NTDS Data Dictionary, Released July 2025



1

Cardiac arrest is the sudden cessation of cardiac activity after hospital arrival. The patient becomes unresponsive with no normal breathing and no signs of circulation. If corrective measures are not taken rapidly, this condition progresses to sudden death.

EXCLUDE patients whose ONLY episode of cardiac arrest with CPR was on arrival to your hospital.

## CATHETER-ASSOCIATED URINARY TRACT INFECTION (CAUTI)

---

### Element Intent

A potentially preventable event often induced by bacteria entering the urinary tract through the catheter, requiring treatment which could increase the hospital length of stay and the likelihood of mortality.

### Description

A UTI where an indwelling urinary catheter was in place for > 2 calendar days on the date of the event, with day of device placement being day 1,

### AND

An indwelling urinary catheter was in place on the date of event or the day before. If an indwelling urinary catheter was in place for more than 2 consecutive days in an inpatient location and then removed, the date of the event for the UTI must be day of device discontinuation or the next day for the UTI to be catheter-associated.

### January 2019 CDC CAUTI Criterion SUTI 1a:

Patient must meet 1, 2, **and** 3 below:

1. Patient had an indwelling urinary catheter that had been in place for more than 2 consecutive days in an inpatient location on the date of the event AND was either:
  - Present for any portion of the calendar day on the date of event, OR
  - Removed the day before the date of event
2. Patient has at least one of the following signs or symptoms:
  - Fever ( $\geq 38^{\circ}\text{C}$ ): Reminder: to use fever in a patient >65 years of age, the IUC needs to be in place for more than 2 consecutive days in an inpatient location on date of event and is either still in place OR was removed the day before the DOE.
  - Suprapubic tenderness with no other recognized cause
  - Costovertebral angle pain or tenderness
  - Urinary urgency
  - Urinary frequency
  - dysuria
3. Patient has a urine culture with no more than two species of organisms, at least one of which is a bacteria  $> 10^5$  CFU/ml.

### January 2019 CDC CAUTI Criterion SUTI 2:

Patient must meet 1, 2, **and** 3 below:

1. Patient is  $\leq 1$  year of age
2. Patient has at least one of the following signs or symptoms:
  - Fever ( $> 38.0^{\circ}\text{C}$ )
  - Hypothermia ( $< 36.0^{\circ}\text{C}$ )
  - Apnea
  - Bradycardia
  - Lethargy
  - Vomiting
  - Suprapubic tenderness

3. Patient has a urine culture with no more than two species of organisms, at least one of which is bacteria of  $\geq 10^5$  CFU/ml.

**Element Values**

- 1 Yes
- 2 No

**Additional Information**

- Onset of symptoms began after arrival to your ED/hospital.
- A diagnosis of UTI must be documented in the patient's medical record.
- Consistent with the January 2019 CDC defined CAUTI.

**Data Source Hierarchy Guide**


- 1 History and Physical
- 2 Physician's Notes
- 3 Progress Notes
- 4 Case Management/Social Services
- 5 Nursing Notes/Flow Sheet
- 6 Triage/Trauma Flow Sheet
- 7 Discharge Summary

**References to Other Databases**

- NTDS 2026


## Catheter-Associated Urinary Tract Infection (CAUTI) SUTI 1a (Patients >1 year of age)

2026 NTDS Data Dictionary, Released July 2025





## 2026 NTDS Data Dictionary, Released July 2025



## CENTRAL LINE-ASSOCIATED BLOODSTREAM INFECTION (CLABSI)

---

### Element Intent

A potentially preventable event often induced by bacteria entering the bloodstream through the central line, requiring treatment which could increase the hospital length of stay and the likelihood of mortality.

### Description

A laboratory-confirmed bloodstream infection (LCBI) where central line (CL) or umbilical catheter (UC) was in place for > 2 calendar days on the date of event, with day of device placement being Day 1,

#### AND

The line was also in place on the date of event or the day before. If a CL or UC was in place for > 2 calendar days and then remove, the date of event of the LCBI must be the day of discontinuation or the next day to be a CLABSI. If the patient is admitted or transferred into a facility with an implanted central line (port) in place, and that is the patient's central line, day of first access in an inpatient location is considered Day. "Access" is defined as line placement, infusion or withdrawal through the line. Such lines continue to be eligible for CLABSI once they are accessed until they are either discontinued or the day after patient discharge (as per the Transfer Rule). Note that the "de-access" of a port does not result in the patient's removal from CLABSI surveillance.

### January 2016 CDC Criterion LCBI 1:

Patient has a recognized pathogen identified from one or more blood specimens by a culture or non-culture based microbiologic testing method which is performed for purposes of clinical diagnosis or treatment (e.g., not Active Surveillance Culture/Testing (ASC/AST)).

#### AND

Organism(s) identified in blood is not related to an infection at another site.

#### OR

### January 2016 CDC Criterion LCBI 2:

Patient has at least one of the following signs or symptoms:

- Fever (>38°C)
- Chills
- Hypotension

#### AND

Organism(s) identified from blood is not related to an infection at another site

#### AND

The same common commensal (i.e., diphtheroids [Corynebacterium spp. Not C. diphtheria], Bacillus spp. [not B. anthracis], Propionibacterium spp., coagulase-negative staphylococci [including S. epidermidis], viridans group streptococci, Aerococcus spp., and Micrococcus spp.) is identified from two or more blood specimens drawn on separate occasions, by a culture or non-culture based microbiologic testing method which is performed for purposes of clinical diagnosis or treatment (e.g., not Active Surveillance Culture/Testing (ASC/AST)). Criterion elements must occur within the Infection Window Period, the 7-day time period which includes the collection date of the positive blood, the 3 calendar days before and the 3 calendar days after.

#### OR

### **January 2016 CDC Criterion LCBI 3:**

Patient  $\leq 1$  year of age has at least one of the following signs or symptoms:

- Fever ( $>38^{\circ}\text{C}$ )
- Hypothermia ( $<36^{\circ}\text{C}$ )
- Apnea
- Bradycardia

**AND**

Organism(s) identified from blood is not related to an infection at another state

**AND**

The same common commensal (i.e., diphtheroids [*Corynebacterium* spp. Not *C. diphtheria*], *Bacillus* spp. [not *B. anthracis*], *Propionibacterium* spp., coagulase-negative staphylococci [including *S. epidermidis*], viridans group streptococci, *Aerococcus* spp., and *Micrococcus* spp.) is identified from two or more blood specimens drawn on separate occasions, by a culture or non-culture based microbiologic testing method which is performed for purposes of clinical diagnosis or treatment (e.g., not Active Surveillance Culture/Testing (ASC/AST). Criterion elements must occur within the Infection Window Period, the 7-day time period which includes the collection date of the positive blood, the 3 calendar days before and the 3 calendar days after.

### **Element Values**

- 1 Yes
- 2 No

### **Additional Information**

- Onset of symptoms began after arrival to your ED/hospital.
- A diagnosis of CLABSI must be documented in the patient's medical record.
- Consistent with the January 2016 CDC defined CLABSI.

### **Data Source Hierarchy Guide**


- 1 History and Physical
- 2 Physician's Notes
- 3 Progress Notes
- 4 Case Management/Social Services
- 5 Nursing Notes/Flow Sheet
- 6 Triage/Trauma Flow Sheet
- 7 Discharge Summary

### **References to Other Databases**

- NTDS 2026

## Central Line-Associated Blood Stream Infection (CLABSI) (pg. 1 of 3)

2026 NTDS Data Dictionary, Released July 2025




**1** The NTDS definition is consistent with the January 2016 CDC CLABSI definition. If information not contained in the NTDS definition is needed, please refer to the CDC or your hospital's infection control department.

**2** The Date of Event is the date the first element used to meet an NHSN site-specific infection criterion occurs for the first time within the seven-day infection window period. The day of device placement is Day 1.

**3** If a CL or UC was in place for >2 calendar days and then removed, the date of event of the CLABSI must be the day of discontinuation or the next day to be a CLABSI. If the patient is admitted or transferred into a facility with an implanted central line (port) in place, and that is the patient's only central line, day of first access in an inpatient location is considered Day 1. "Access" is defined as line placement, infusion or withdrawal through the line. Such lines continue to be eligible for CLABSI once they are accessed until they are either discontinued or the day after patient discharge (as per the Transfer Rule). Note that the "de-access" of a port does not result in the patient's removal from CLABSI surveillance.

## Central Line-Associated Blood Stream Infection (CLABSI) (pg. 2 of 3)

2026 NTDS Data Dictionary, Released July 2025




4

Criterion elements must occur within the Infection Window Period, the 7-day time period which includes the collection date of the positive blood, the 3 calendar days before and the 3 calendar days after.

## Central Line-Associated Blood Stream Infection (CLABSI) (pg. 3 of 3)

2026 NTDS Data Dictionary, Released July 2025



5

Criterion elements must occur within the Infection Window Period, the 7-day time period which includes the collection date of the positive blood, the 3 calendar days before and the 3 calendar days after.

## DEEP SURGICAL SITE INFECTION

### Element Intent

A potentially preventable event often induced by bacteria, viruses, or endogenous flora contacting a surgical wound, requiring treatment which could increase the hospital length of stay and the likelihood of mortality.

### Description

Must meet the following criteria:

Infection occurs within 30 or 90 days after the NHSN operative procedure (where day 1 = the procedure date) According to list in Table 2

### AND

Patient has at least one of the following:

- Purulent drainage from the deep incision
- A deep incision that spontaneously dehisces, or is deliberately opened or aspirated by a surgeon, attending physician\*\* or other designee and organism is identified by a culture or non-culture based microbiologic testing method which is performed for purposed of clinical diagnosis or treatment (e.g., not Active Surveillance Culture/Testing (ACS/AST) or culture or non-culture based microbiologic test method is not performed

### AND

Patient has at least one of the following signs or symptoms:

- Fever (>38°C)
- Localized pain or tenderness
- A culture or non-culture based test that has a negative finding does not meet this criterion
- An abscess or other evidence of infection involving the deep incision that is detected on gross anatomical or histopathologic exam, or imaging test

COMMENTS: There are two specific types of deep incisional SSIs:

- Deep Incisional Primary (DIP): a deep incisional SSI that is identified in a primary incision in a patient that has had an operation with one or more incisions (e.g., C-section incision or chest incision for CBGB)
- Deep Incisional Secondary (DIS): a deep incisional SSI that is identified in the secondary incision in a patient that has had an operation with more than one incision (e.g., donor site [leg] incision for CBGB.)

**Table 2. Surveillance Period for Deep Incisional or Organ/Space SSI Following Selected NHSN Operative Procedure Categories. Day 1 = the date of the procedure.**

30- day Surveillance			
Code	Operative Procedure	Code	Operative Procedure
AAA	Abdominal Aortic Aneurysm repair	LAM	Laminectomy
AMP	Limb Amputation	LTP	Liver transplant
APPY	Appendix Surgery	NECK	Neck surgery
AVSD	Shunt for dialysis	NEPH	Kidney surgery
BIBL	Bile duct, liver or pancreatic surgery	OVRV	Ovarian surgery
CEA	Carotid endarterectomy	PRST	Prostate surgery
CHOL	Gallbladder Surgery	REC	Rectal surgery

COLO	Colon Surgery	SB	Small bowel surgery
CSEC	Cesarean Section	SPLE	Spleen surgery
GAST	Gastric surgery	THOR	Thoracic surgery

HTP	Heart transplant	THUR	Thyroid and/or parathyroid surgery
HYST	Abdominal hysterectomy	VHYS	Vaginal hysterectomy
KTP	Kidney transplant	XLAP	Exploratory Laparotomy

#### 90- day Surveillance

Code	Operative Procedure
BRST	Breast surgery
CARD	Cardiac surgery
CBGB	Coronary artery bypass graft with both chest and donor site incisions
CBGC	Coronary artery bypass graft with check incision only
CRAN	Craniotomy
FUSN	Spinal fusion
FX	Open reduction of fracture
HER	Herniorrhaphy
HPRO	Hip prosthesis
KPRO	Knee prosthesis
PACE	Pacemaker surgery
PVBY	Peripheral vascular bypass surgery
VSHN	Ventricular shunt

#### Element Values

- 1 Yes
- 2 No

#### Additional Information

- Onset of symptoms began after arrival to your ED/hospital.
- A diagnosis of SSI must be documented in the patient's medical record.
- Consistent with the CDC January 2024 defined SSI.

#### Data Source Hierarchy Guide

- 1 History and Physical
- 2 Physician's Notes
- 3 Progress Notes
- 4 Case Management/Social Services
- 5 Nursing Notes/Flow Sheet
- 6 Triage/Trauma Flow Sheet
- 7 Discharge Summary


#### References to Other Databases

- NTDS 2026



## Deep Surgical Site Infection (Deep SSI) (pg. 1 of 2)

2026 NTDS Data Dictionary, Released July 2025




1

The NTDS definition is consistent with the January 2024 CDC SSI definition. If information not contained in the NTDS definition is needed, please refer to the CDC or your hospital's infection control department.

## Deep Surgical Site Infection (Deep SSI) (pg. 2 of 2)

2026 NTDS Data Dictionary, Released July 2025



**2**

The term **attending physician** for the purposes of application of the NHSN SSI criteria may be interpreted to mean the surgeon(s), infectious disease, other physician on the case, emergency physician, or physician's designee (nurse practitioner or physician's assistant).

## DEEP VEIN THROMBOSIS (DVT)

---

### Element Intent

A potentially preventable event often induced by immobility, anesthesia, stroke, venous catheters, dehydration, and/or thrombocytosis, requiring treatment which could increase the hospital length of stay and the likelihood of mortality.

### Description

The formation, development, or existence of a blood clot or thrombus within the venous system, which may be coupled with inflammation.

### Element Values

- 1 Yes
- 2 No

### Additional Information

- Onset of symptoms began after arrival to your ED/hospital.
- The patient must be treated with anticoagulation therapy and/or placement of a vena cava filter or clipping of the vena cava.
- A diagnosis of DVT must be documented in the patient's medical record, which may be confirmed by venogram, ultrasound, or CT.

### Data Source Hierarchy Guide


- 1 History and Physical
- 2 Physician's Notes
- 3 Progress Notes
- 4 Case Management/Social Services
- 5 Nursing Notes/Flow Sheet
- 6 Triage/Trauma Flow Sheet
- 7 Discharge Summary

### References to Other Databases

- NTDS 2026

## Deep Vein Thrombosis (DVT)

2026 NTDS Data Dictionary, Released July 2025



# DELIRIUM

---

## Element Intent

A potentially preventable event often induced by infection, stroke, lung or liver disease, medications, low sodium, low blood sugar, urinary retention, dehydration, low oxygen, or an unfamiliar environment, requiring treatment which could increase the hospital length of stay and the likelihood of mortality.

## Description

Acute onset of behaviors characterized by restlessness, illusions, and incoherence of thought and speech. Delirium can often be traced to one or more contributing factors, such as severe or chronic medical illness, changes in your metabolic balance (such as low sodium), medication, infection, surgery, or alcohol or drug withdrawal.

### OR

Patient tests positive after using an objective screening tool like the Confusion Assessment Method (CAM or the Intensive Care Delirium Screening Checklist (ICDSC).

### OR

A diagnosis of delirium documented in the patient's medical record.

## EXCLUDE:

- Patients whose delirium is due to alcohol withdrawal.

## Element Values

- 1 Yes
- 2 No

## Additional Information

- Onset of symptoms began after arrival to your ED/hospital.

## Data Source Hierarchy Guide


- 1 History and Physical
- 2 Physician's Notes
- 3 Progress Notes
- 4 Case Management/Social Services
- 5 Nursing Notes/Flow Sheet
- 6 Triage/Trauma Flow Sheet
- 7 Discharge Summary

## References to Other Databases

- NTDS 2026

## Delirium

2026 NTDS Data Dictionary, Released July 2025



Delirium can often be traced to one or more contributing factors, such as a severe or chronic medical illness, changes in your metabolic balance (such as low sodium), medication, infection, surgery, or alcohol or drug withdrawal.

## MYOCARDIAL INFARCTION (MI)

---

### Element Intent

A potentially preventable event often induced by coronary artery disease, medications, emotional stress, or pain, requiring treatment which could increase the hospital length of stay and the likelihood of mortality.

### Description

An acute myocardial infarction must be noted with documentation of ECG changes indicative of acute MI

### AND

New elevation in troponin greater than three times upper level of the reference range in the setting of suspected myocardial ischemia

### AND

Physician diagnosis of an acute myocardial infarction that occurred subsequent to arrival at your center

### Element Values

- 1 Yes
- 2 No

### Additional Information

- Onset of symptoms began after arrival to your ED/hospital.

### Data Source Hierarchy Guide


- 1 History and Physical
- 2 Physician's Notes
- 3 Progress Notes
- 4 Case Management/Social Services
- 5 Nursing Notes/Flow Sheet
- 6 Triage/Trauma Flow Sheet
- 7 Discharge Summary

### References to Other Databases

- NTDS 2026

## Myocardial Infarction (MI)

2026 NTDS Data Dictionary, Released July 2025





## ORGAN/SPACE SURGICAL SITE INFECTION

### Element Intent

A potentially preventable event often induced by bacteria or endogenous flora contacting a surgical wound, requiring treatment which could increase the hospital length of stay and the likelihood of mortality.

### Description

Must meet the following criteria:

Infection that occurs within 30 or 90 days after the NHS operative procedure (where da 1 = the procedure date) according to the list in Table 2

### AND

Infection involves any part of the body deeper than the fascial/muscle layers, that is opened or manipulated during the operative procedure

### AND

Patient has at least **one** of the following:

- Purulent drainage from a drain that is placed into the organ/space (e.g., closed suction drainage system, open drain, T-tube drain, CT guided drainage)
- Organisms are identified from an aseptically-obtained fluid or tissue in the organ/space by a culture or non-culture based microbiologic testing method which is performed for purposes of clinical diagnosis or treatment) e.g., not Active Surveillance Culture/Testing (ASC/AST).
- An abscess or other evidence of infection involving the organ/space that is detected on gross anatomical or histopathologic exam, or imaging test

### AND

Meets at least one criterion for a specific organ/space infection site listed in Table 3. These criteria are found in the Surveillance Definitions for Specific Types of Infections chapter.

**Table 2. Surveillance Period for Deep Incisional or Organ/Space SSI Following Selected NHSN Operative Procedure Categories. Day 1 = the date of the procedure.**

30- day Surveillance			
Code	Operative Procedure	Code	Operative Procedure
AAA	Abdominal Aortic Aneurysm repair	LAM	Laminectomy
AMP	Limb Amputation	LTP	Liver transplant
APPY	Appendix Surgery	NECK	Neck surgery
AVSD	Shunt for dialysis	NEPH	Kidney surgery
BIBL	Bile duct, liver or pancreatic surgery	OVRV	Ovarian surgery
CEA	Carotid endarterectomy	PRST	Prostate surgery
CHOL	Gallbladder Surgery	REC	Rectal surgery
COLO	Colon Surgery	SB	Small bowel surgery
CSEC	Cesarean Section	SPLE	Spleen surgery
GAST	Gastric surgery	THOR	Thoracic surgery
HTP	Heart transplant	THUR	Thyroid and/or parathyroid surgery
HYST	Abdominal hysterectomy	VHYS	Vaginal hysterectomy
KTP	Kidney transplant	XLAP	Exploratory Laparotomy
90- day Surveillance			
Code	Operative Procedure		
BRST	Breast surgery		

CARD	Cardiac surgery
CBGB	Coronary artery bypass graft with both chest and donor site incisions
CBGC	Coronary artery bypass graft with check incision only
CRAN	Craniotomy
FUSN	Spinal fusion
FX	Open reduction of fracture
HER	Herniorrhaphy
HPRO	Hip prosthesis
KPRO	Knee prosthesis
PACE	Pacemaker surgery
PVBY	Peripheral vascular bypass surgery
VSHN	Ventricular shunt

**Table 3. Specific Sites of an Organ/Space SSI**

Code	Site	Code	Site
BONE	Osteomyelitis	LUNG	Other infections of respiratory tract
BRST	Breast abscess mastitis	MED	Mediastinitis
CARD	Myocarditis or Pericarditis	MEN	Meningitis or ventriculitis
DISC	Disc space	ORAL	Oral cavity (mouth, tongue, or gums)
EAR	Ear, Mastoid	OREP	Other infections of the male or female reproductive tract
EMET	Endometritis	PJI	Periprosthetic Joint Infection
ENDO	Endocarditis	SA	Spinal abscess without meningitis
EYE	Eye, other than conjunctivitis	SINU	Sinusitis
GIT	GI Tract	UR	Upper respiratory tract
HEP	Hepatitis	USI	Urinary System Infection
IAB	Intraabdominal, not specified	VASC	Arterial or venous infection
IC	Intracranial, brain abscess or dura	VCUF	Vaginal cuff
JNT	Joint or bursa		

#### Element Values

- 1 Yes
- 2 No

#### Additional Information

- Onset of symptoms began after arrival to your ED/hospital
- A diagnosis of SSI must be documented in the patient's medical record.
- Consistent with the January 2019 CDC defined SSI.

#### Data Source Hierarchy Guide


- 1 History and Physical
- 2 Physician's Notes
- 3 Progress Notes
- 4 Case Management/Social Services
- 5 Nursing Notes/Flow Sheet
- 6 Triage/Trauma Flow Sheet
- 7 Discharge Summary

#### References to Other Databases

- NTDS 2026

## Organ/Space Surgical Site Infection (O/S SSI)

2026 NTDS Data Dictionary, Released July 2025



# OSTEOMYELITIS

---

## Element Intent

A potentially preventable event often induced by bacteria or fungi, diabetes, and/or a weakened immune system, requiring treatment which could increase the hospital length of stay and the likelihood of mortality.

## Description

Osteomyelitis must meet at least one of the following criteria:

1. Patient has organisms identified by culture or non-cultured based microbiologic testing method which is performed for purposes of clinical diagnosis and treatment (e.g., not Active Surveillance Culture/Testing (ASC/ASST)).
2. Patient has evidence of osteomyelitis on gross anatomic or histopathologic examination.
3. Patient has at least two of the following localized signs or symptoms:
  - Fever ( $>38^{\circ}\text{C}$ )
  - Swelling\*
  - Pain or Tenderness\*
  - Heat\*
  - Drainage\*

**AND at least one of the following:**

- a. Organisms identified from blood by culture or non-culture based microbiologic testing method, which is performed for purposes of clinical diagnosis and treatment, for example, not Active Surveillance Culture/Testing (ASC/AST) AND Imaging test evidence suggestive of infection (for example, x-ray, CT scan, MRI, radiolabel scan [gallium, technetium, etc.]), which if equivocal is supported by clinical correlation, specifically, physician documentation of antimicrobial treatment for osteomyelitis.
- b. Imaging test evidence suggestive of infection (for example, x-ray, CT scan, MRI, radiolabel scan [gallium, technetium, etc.]), which if equivocal is supported by clinical correlation, specifically, physician documentation of antimicrobial treatment for osteomyelitis).

\*With no other recognized cause

## Element Values

- 1 Yes
- 2 No

## Additional Information

- Onset of symptoms began after arrival to your ED/hospital.
- A diagnosis of osteomyelitis must be documented in the patient's medical record.
- Consistent with the January 2020 CDC definition of Bone and Joint Infection.

## Data Source Hierarchy Guide


- 1 History and Physical
- 2 Physician's Notes
- 3 Progress Notes
- 4 Case Management/Social Services
- 5 Nursing Notes/Flow Sheet
- 6 Triage/Trauma Flow Sheet
- 7 Discharge Summary

## References to Other Databases

- NTDS 2026

## Osteomyelitis

2026 NTDS Data Dictionary, Released July 2025



**1** The NTDS definition is consistent with the January 2020 CDC Bone and Joint infection definition. If information not contained in the NTDS definition is needed, please refer to the CDC or your hospital's infection control department.

## PULMONARY EMBOLISM (PE)

---

### Element Intent

A potentially preventable event requiring treatment which could increase the hospital length of stay and the likelihood of mortality.

### Description

A lodging of a blood clot in a pulmonary artery with subsequent obstruction of blood supply to the lung parenchyma. The blood clots usually originate from the deep leg veins or the pelvic venous system.

### EXCLUDE:

- Subsegmental PEs.

### Element Values

- 1 Yes
- 2 No

### Additional Information

- Onset of symptoms began after arrival to your ED/hospital.
- Consider the condition present if the patient has a V-Q scan interpreted as high probability of pulmonary embolism or a positive pulmonary arteriogram or positive CT angiogram and/or a diagnosis of PE is documented in the patient's medical record.

### Data Source Hierarchy Guide


- 1 History and Physical
- 2 Physician's Notes
- 3 Progress Notes
- 4 Case Management/Social Services
- 5 Nursing Notes/Flow Sheet
- 6 Triage/Trauma Flow Sheet
- 7 Discharge Summary

### References to Other Databases

- NTDS 2026

## Pulmonary Embolism (PE)

2026 NTDS Data Dictionary, Released July 2025



Subsegmental PEs are excluded from the NTDS definition of PE.

## PRESSURE ULCER

---

### Element Intent

A potentially preventable event often induced by pressure or friction, moisture or other medical factors; advancement to stage II or greater requires treatment which could increase the hospital length of stay and the likelihood of mortality.

### Description

A localized injury to the skin and/or underlying tissue usually over a bony prominence, as a result of pressure, or pressure in combination with shear. A number of contributing or confounding factors are also associated with pressure ulcers; the significance of these factors is yet to be elucidated. Equivalent to NPUAP Stages II-IV, Unstageable/Unclassified, and Suspected Deep Tissue Injury.

### Element Values

- 1 Yes
- 2 No

### Additional Information

- Onset of NPUAP Stage II began after arrival to your ED/hospital.
- Pressure ulcer documentation must be in the patient's medical record.
- Consistent with the NPUAP 2014.

### Data Source Hierarchy Guide

- 1 History and Physical
- 2 Physician's Notes
- 3 Progress Notes
- 4 Case Management/Social Services
- 5 Nursing Notes/Flow Sheet
- 6 Triage/Trauma Flow Sheet
- 7 Discharge Summary


### References to Other Databases

- NTDS 2026



## Pressure Ulcer

2026 NTDS Data Dictionary, Released July 2025



1

A pressure ulcer is a localized injury to the skin and/or underlying tissue usually over a bony prominence, as a result of pressure, or pressure in combination with shear. A number of contributing or confounding factors are also associated with pressure ulcers; the significance of these factors is yet to be elucidated.

2

The NTDS definition is consistent with the NPUAP 2014.

## SEVERE SEPSIS

---

### Element Intent

A potentially preventable event often induced by bacterial, viral or fungal infections, requiring treatment which could increase the hospital length of stay and the likelihood of mortality.

### Description

Severe sepsis: sepsis plus organ dysfunction, hypotension (low blood pressure), or hypoperfusion (insufficient blood flow) to 1 or more organs.

Septic shock: sepsis with persisting arterial hypotension or hypoperfusion despite adequate fluid resuscitation.

### Element Values

- 1 Yes
- 2 No

### Additional Information

- Onset of symptoms began after arrival to your ED/hospital.
- A diagnosis of sepsis must be documented in the patient's medical record.
- Consistent with the American College of Chest Physicians and the Society of Critical Care Medicine October 2010.

### Data Source Hierarchy Guide


- 1 History and Physical
- 2 Physician's Notes
- 3 Progress Notes
- 4 Case Management/Social Services
- 5 Nursing Notes/Flow Sheet
- 6 Triage/Trauma Flow Sheet
- 7 Discharge Summary

### References to Other Databases

- NTDS 2026

## Severe Sepsis

2026 NTDS Data Dictionary, Released July 2025



**1** The NTDS definition is consistent with the American College of Chest Physicians and the Society of Critical Care Medicine October 2010.

## STROKE/CVA

---

### Element Intent

A potentially preventable event often induced by obstruction of blood flow or a ruptured blood vessel in the brain, requiring treatment which could increase the hospital length of stay and the likelihood of mortality.

### Description

A focal or global neurological deficit of rapid onset and NOT present on admission. The patient must have at least one of the following symptoms:

- Change in level of consciousness
- Hemiplegia
- Hemiparesis
- Numbness or sensory loss affecting on side of the body
- Dysphasia or aphasia
- Hemianopia
- Amaurosis fugax
- Other neurological signs or symptoms consistent with stroke

#### AND

- Duration of neurological deficit  $\geq 24$  h

#### OR

- Duration of deficit  $< 24$  h, if neuroimaging (MR, CT, or cerebral angiography) documents a new hemorrhage or infarct consistent with stroke, or therapeutic intervention(s) were performed for stroke, or the neurological deficit results in death

#### AND

- No other readily identifiable non-stroke cause, e.g., progression of existing traumatic brain injury, seizure, tumor, metabolic or pharmacologic etiologies, is identified

#### AND

- Diagnosis is confirmed by neurology or neurosurgical specialist or neuroimaging procedure (MR, CT, angiography,) or lumbar puncture (CSF demonstrating intracranial hemorrhage that was not present on admission.)

### Element Values

- 1 Yes
- 2 No

### Additional Information

- Onset of symptoms began after arrival to your ED/hospital.
- A diagnosis of stroke/CVA must be documented in the patient's medical record.
- Although the neurologic deficit must not present on admission, risk factors predisposing to stroke (e.g., blunt cerebrovascular injury, dysrhythmia) may be present on admission.

### Data Source Hierarchy Guide

- 1 History and Physical
- 2 Physician's Notes
- 3 Progress Notes
- 4 Case Management/Social Services


- 5 Nursing Notes/Flow Sheet
- 6 Triage/Trauma Flow Sheet
- 7 Discharge Summary

#### **References to Other Databases**

- NTDS 2026

## Stroke/CVA

2026 NTDS Data Dictionary, Released July 2025



--Page Blank--

## SUPERFICIAL INCISIONAL SURGICAL SITE INFECTION

---

### Element Intent

A potentially preventable event often induced by endogenous flora or exogenous contamination contacting a surgical site, requiring treatment which could increase the hospital length of stay and the likelihood of mortality.

### Description

Must meet the following criteria:

Infection occurs within 30 days after any NHSN operative procedure (where day 1 = the procedure date)

### AND

Involves only skin or subcutaneous tissue of the incision

### AND

Patient has at least **one** of the following:

- a. Purulent drainage from the superficial incision.
- b. Organisms identified from an aseptically-obtained specimen from the superficial incision or subcutaneous tissue by a culture or non-culture based microbiologic testing method which is performed for purposes of clinical diagnosis or treatment (e.g., not Active Surveillance Culture/Testing (ASC/AST).
- c. Superficial incision is deliberately opened by the surgeon, attending physician\*\* or other designee and culture or non-culture based testing is not performed

### AND

Patient has at least one of the following signs or symptoms:

- Pain or tenderness
  - Localized swelling
  - Erythema
  - Heat
  - A culture or non-culture based test that has a negative finding does not meet this criterion
- d. Diagnosis of Superficial incisional SSI by the surgeon or attending physician\*\* or other designee.

COMMENTS: There are two specific types of superficial incisional SSIs:

1. Superficial Incisional Primary (SIP)- a superficial incisional SSI that is identified in the primary incision in a patient that has had an operation with one or more incisions (e.g., C-section incision or chest incision for CBGB)
2. Superficial Incisional Secondary (SIS)- a superficial incisional SSI that is identified in the secondary incision in a patient that has had an operation with more than one incision (e.g., donor site incision for CBGB)

### Element Values

- 1 Yes
- 2 No

### Additional Information

- Onset of symptoms began after arrival to your ED/hospital.
- A diagnosis of SSI must be documented in the patient's medical record.
- Consistent with the January 2019 CDC defined SSI.



**Data Source Hierarchy Guide**


- 1 History and Physical
- 2 Physician's Notes
- 3 Progress Notes
- 4 Case Management/Social Services
- 5 Nursing Notes/Flow Sheet
- 6 Triage/Trauma Flow Sheet
- 7 Discharge Summary

**References to Other Databases**

- NTDS 2026

## Superficial Incisional Surgical Site Infection (S/I SSI)

2026 NTDS Data Dictionary, Released July 2025



## UNPLANNED ADMISSION TO ICU

---

### Element Intent

A potentially preventable event that highlights possible gaps in the assessment of the severity of the patient's condition or the application of appropriate treatment plans.

### Description

Patients admitted to the ICU after initial transfer to the floor, and/or patients with an unplanned return to the ICU after initial ICU discharge.

#### INCLUDE:

- Patients who required ICU care due to an event that occurred during surgery or in the PACU.

#### EXCLUDE:

- Patients with a planned post-operative ICU stay.

### Element Values

- 1 Yes
- 2 No

### Additional Information

- Must have occurred during the patient's initial stay at your hospital.

### Data Source Hierarchy Guide


- 1 History and Physical
- 2 Physician's Notes
- 3 Progress Notes
- 4 Case Management/Social Services
- 5 Nursing Notes/Flow Sheet
- 6 Triage/Trauma Flow Sheet
- 7 Discharge Summary

### References to Other Databases

- NTDS 2026

## Unplanned Admission to the ICU

2026 NTDS Data Dictionary, Released July 2025



1

Floor is defined as any other ward that is not an ICU. "Floor" also includes step-down units and the operating room for the purposes of this definition.

EXCLUDE: Patients with a planned post-operative ICU stay. This means that it was known prior to surgery that the patient would require postoperative ICU care.

INCLUDE: Patients who required ICU care due to an event that occurred during surgery or in the PACU.

## UNPLANNED INTUBATION

---

### Element Intent

A potentially preventable event that highlights possible gaps in the assessment of the severity of the patient's condition or the application of appropriate treatment plans.

### Description

Patient requires placement of an endotracheal tube and mechanical or assisted ventilation manifested by severe respiratory distress, hypoxia, hypercarbia, or respiratory acidosis.

### Element Values

- 1 Yes
- 2 No

### Additional Information

- Must have occurred during the patient's initial stay at your hospital.
- In patients who were intubated in the field or Emergency Department, or those intubated for surgery, unplanned intubation occurs if they require reintubation > 24 hours after extubation.

### Data Source Hierarchy Guide


- 1 History and Physical
- 2 Physician's Notes
- 3 Progress Notes
- 4 Case Management/Social Services
- 5 Nursing Notes/Flow Sheet
- 6 Triage/Trauma Flow Sheet
- 7 Discharge Summary

### References to Other Databases

- NTDS 2026

## Unplanned Intubation

2026 NTDS Data Dictionary, Released July 2025



## UNPLANNED RETURN TO THE OPERATING ROOM

---

### Element Intent

A potentially preventable event that highlights possible opportunities for improvements in care.

### Description

The patient underwent a subsequent operative procedure at the same operative site as the initial operative procedure. Both procedures must have been performed in the operating room at your center.

### EXCLUDE:

- Planned return to the operating room after damage control surgery or staged surgical interventions.
- Procedures performed in an interventional radiology suite.
- Procedures performed in a hybrid operating room where the intervention is limited to a percutaneous approach.
- Pre-planned multiple-stage approach procedures.

### Element Values

- 1 Yes
- 2 No

### Additional Information

- The same operative site usually (but not exclusively) implies there was a need to re-open the previous incision.
- *Element Value "1. Yes"* is reported whether the initial intervention was related to the injuries (e.g., anastomotic leak after laparotomy, hardware failure/infection after ORIF of fractures) OR if there is a return to the operating room for an unplanned intervention related to a secondary procedure (e.g., return to the OR for bleeding after tracheostomy).
- *Element Value "2. No"* is reported if there is intent to return to the operating room for a two-stage approach.

### Data Source Hierarchy Guide


- 1 Operative Report
- 2 Physician Notes/Flow Sheet
- 3 Progress Notes
- 4 Nursing Notes/Flow Sheet
- 5 Discharge Summary

### References to Other Databases

- NTDS 2026

## Unplanned Return to the Operating Room

2026 NTDS Data Dictionary, Released July 2025



1

The same operative site usually (but not exclusively) implies there was the need to re-open the previous incision.

### EXCLUDE:

- Planned return to the operating room after damage control surgery or staged surgical interventions.
- Procedures performed in an interventional radiology suite.
- Procedures performed in a hybrid operating room where the intervention is limited to a percutaneous approach.
- Pre-planned multiple-stage approach procedures.



VENTILATOR-ASSOCIATED PNEUMONIA (VAP)

Element Intent

A potentially preventable event often induced by bacteria or virus entering the lungs, requiring treatment which could increase the hospital length of stay and the likelihood or mortality.

Description

A pneumonia where the patient is on mechanical ventilation for > 2 calendar days on the date of event, with day of ventilator placement being Day 1,

AND

The ventilator was in place on the date of event or the day before.

VAP Algorithm (PNU2 Bacterial or Filamentous Fungal Pathogens):

IMAGING TEST EVIDENCE	SIGNS/SYMPTOMS	LABORATORY
<p>Two or more serial chest imaging test results with at least <b>one</b> of the following:</p> <ul style="list-style-type: none"><li>• New or progressive <b>and</b> persistent infiltrate</li><li>• Consolidation</li><li>• Cavitation</li><li>• Pneumatocoles, in infants ≤1 year old</li></ul> <p>NOTE: In patients <b>without</b> underlying pulmonary or cardiac disease (e.g., respiratory distress syndrome, bronchopulmonary dysplasia, pulmonary edema, or chronic obstructive pulmonary disease), <b>one definitive</b> chest imaging test result is acceptable.</p>	<p>At least <b>one</b> of the following:</p> <ul style="list-style-type: none"><li>• Fever (&gt;38°C or &gt;100.4°F)</li><li>• Leukopenia (&lt;4000 WBC/mm³) or leukocytosis (≥12,000 WBC/mm³)</li><li>• For adults ≥70 years old, altered mental status with no other recognized cause</li></ul> <p><b>AND at least two of the following:</b></p> <ul style="list-style-type: none"><li>• New onset of purulent sputum, or change in character of sputum, or increased respiratory secretions, or increased suctioning requirements</li><li>• New onset or worsening cough, or dyspnea, or tachypnea</li><li>• Rales or bronchial breath sounds</li><li>• Worsening gas exchange (e.g., O<sub>2</sub> desaturations (e.g., PaO<sub>2</sub>/FiO<sub>2</sub>≤240), increased oxygen requirements, or increased ventilator demand)</li></ul>	<p>At least <b>one</b> of the following:</p> <ul style="list-style-type: none"><li>• Organism identified from blood</li><li>• Organism identified from pleural fluid</li><li>• Positive quantitative culture from minimally-contaminated LRT specimen (e.g., BAL or protected specimen brushing)</li><li>• ≥5% BAL-obtained cells contain intracellular bacteria on direct microscopic exam (e.g., Gram’s stain)</li><li>• Positive quantitative culture of lung tissue</li><li>• Histopathologic exam shows at least <b>one</b> of the following evidences of pneumonia:<ul style="list-style-type: none"><li>○ Abscess formation or foci of consolidation with intense PMN accumulation in bronchioles and alveoli</li><li>○ Evidence of lung parenchyma invasion by fungal hyphae or pseudohyphae</li></ul></li></ul>

**VAP Algorithm (PNU2 Viral, Legionella, and other Bacterial Pneumonias):**

IMAGING TEST EVIDENCE	SIGNS/SYMPTOMS	LABORATORY
<p>Two or more serial chest imaging test results with at least <b>one</b> of the following:</p> <ul style="list-style-type: none"> <li>• New or progressive <b>and</b> persistent infiltrate</li> <li>• Consolidation</li> <li>• Cavitation</li> <li>• Pneumatoceles, in infants <math>\leq 1</math> year old</li> </ul> <p>NOTE: In patients <b>without</b> underlying pulmonary or cardiac disease (e.g., respiratory distress syndrome, bronchopulmonary dysplasia, pulmonary edema, or chronic obstructive pulmonary disease), <b>one definitive</b> chest imaging test result is acceptable</p>	<p>At least <b>one</b> of the following:</p> <ul style="list-style-type: none"> <li>• Fever (<math>&gt;38^{\circ}\text{C}</math> or <math>&gt;100.4^{\circ}\text{F}</math>)</li> <li>• Leukopenia (<math>&lt;4000 \text{ WBC/mm}^3</math>) or leukocytosis (<math>\geq 12,000 \text{ WBC/mm}^3</math>)</li> <li>• For adults <math>\geq 70</math> years old, altered mental status with no other recognized cause</li> </ul> <p><b>AND at least two of the following:</b></p> <ul style="list-style-type: none"> <li>• New onset of purulent sputum, or change in character of sputum, or increased respiratory secretions, or increased suctioning requirements</li> <li>• New onset or worsening cough, or dyspnea, or tachypnea</li> <li>• Rales or bronchial breath sounds</li> <li>• Worsening gas exchange (e.g., <math>\text{O}_2</math> desaturations (e.g., <math>\text{PaO}_2/\text{FiO}_2 \leq 240</math>), increased oxygen requirements, or increased ventilator demand)</li> </ul>	<p>At least <b>one</b> of the following:</p> <ul style="list-style-type: none"> <li>• Virus, <i>Bordetella</i>, <i>Legionella</i>, <i>Chlamydia</i> or <i>Mycoplasma</i> identified from respiratory secretions or tissue by a culture or non-culture based microbiologic testing method which is performed for purposes of clinical diagnosis or treatment (e.g., not Active Surveillance Culture/Testing (ASC/AST).</li> <li>• Fourfold rise in paired sera (IgG) for pathogen (e.g., influenza viruses, Chlamydia)</li> <li>• Fourfold rise in <i>Legionella pneumophila</i> serogroup 1 antibody titer to <math>\geq 1:128</math> in paired acute and convalescent sera by indirect IFA.</li> <li>• Detection of <i>L. pneumophila</i> serogroup 1 antigens in urine by RIA or EIA</li> </ul>

**VAP Algorithm (PNU3 Immunocompromised Patients):**

IMAGING TEST EVIDENCE	SIGNS/SYMPTOMS	LABORATORY
<p>Two or more serial chest radiographs with at least <b>one</b> of the following:</p> <ul style="list-style-type: none"> <li>• New or progressive <b>and</b> persistent infiltrate</li> <li>• Consolidation</li> <li>• Cavitation</li> <li>• Pneumatoceles, in infants <math>\leq 1</math> year old</li> </ul> <p>NOTE: In patients <b>without</b> underlying pulmonary or cardiac disease (e.g., respiratory distress syndrome, bronchopulmonary dysplasia, pulmonary edema, or chronic obstructive pulmonary disease), <b>one definitive</b> chest imaging test result is acceptable</p>	<p>Patient who is immunocompromised has at least <b>one</b> of the following:</p> <ul style="list-style-type: none"> <li>• Fever (<math>&gt;38^{\circ}\text{C}</math> or <math>&gt;100.4^{\circ}\text{F}</math>)</li> <li>• For adults <math>\geq 70</math> years old, altered mental status with no other recognized cause</li> <li>• New onset of purulent sputum, or change in character of sputum, or increased respiratory secretions, or increased suctioning requirements</li> <li>• New onset or worsening cough, or dyspnea, or tachypnea</li> <li>• Rales or bronchial breath sounds</li> <li>• Worsening gas exchange (e.g., <math>\text{O}_2</math> desaturations (e.g., <math>\text{PaO}_2/\text{FiO}_2 \leq 240</math>), increased oxygen requirements, or increased ventilator demand)</li> <li>• Hemoptysis</li> <li>• Pleuritic chest pain</li> </ul>	<p>At least <b>one</b> of the following:</p> <ul style="list-style-type: none"> <li>• Identification of matching <i>Candida</i> spp. from blood and sputum, endotracheal aspirate, BAL or protected specimen brushing.<sup>11,12,13</sup></li> <li>• Evidence of fungi from minimally contaminated LRT specimen (e.g., BAL or protected specimen brushing) from one of the following: <ul style="list-style-type: none"> <li>○ Direct microscopic exam</li> <li>○ Positive culture of fungi</li> <li>○ Non-culture diagnostic laboratory test</li> </ul> </li> </ul> <p>Any of the following from: <b>LABORATORY CRITERIA DEFINED UNDER PNU2</b></p>

**VAP Algorithm ALTERNATE CRITERIA (PNU1), for infants ≤1 year old:**

IMAGING TEST EVIDENCE	SIGNS/SYMPTOMS/LABORATORY
<p>Two or more serial chest imaging test results with at least <b>one</b> of the following:</p> <ul style="list-style-type: none"> <li>• New or progressive <b>and</b> persistent infiltrate</li> <li>• Consolidation</li> <li>• Cavitation</li> <li>• Pneumatoceles, in infants ≤1 year old</li> </ul> <p>NOTE: In patients without underlying pulmonary or cardiac disease (e.g., respiratory distress syndrome, bronchopulmonary dysplasia, pulmonary edema, or chronic obstructive pulmonary disease), <b>one definitive</b> imaging test result is acceptable.</p>	<p>Worsening gas exchange (e.g., O<sub>2</sub> desaturation [e.g. pulse oximetry &lt;94%], increased oxygen requirements, or increased ventilator demand)</p> <p><b>AND</b> at least <b>three</b> of the following:</p> <ul style="list-style-type: none"> <li>• Temperature instability</li> <li>• Leukopenia (&lt;4000 WBC/mm<sup>3</sup>) <b>or</b> leukocytosis (≥15,000 WBC/mm<sup>3</sup>) and left shift (≥10% band forms)</li> <li>• New onset of purulent sputum, or change in character of sputum, or increased respiratory secretions, or increased suctioning requirements</li> <li>• Apnea, tachypnea, nasal flaring with retraction of chest wall, or nasal flaring with grunting</li> <li>• Wheezing, rales, or rhonchi</li> <li>• Cough</li> <li>• Bradycardia (&lt;100 beats/min) or tachycardia (&gt;170 beats/min)</li> </ul>

**VAP Algorithm ALTERNATE CRITERIA (PNU1), for children >1 year old or ≤12 years old:**

IMAGING TEST EVIDENCE	SIGNS/SYMPTOMS/LABORATORY
<p>Two or more serial chest imaging test results with at least <b>one</b> of the following:</p> <ul style="list-style-type: none"> <li>• New or progressive <b>and</b> persistent infiltrate</li> <li>• Consolidation</li> <li>• Cavitation</li> <li>• Pneumatoceles, in infants ≤1 year old</li> </ul> <p>NOTE: In patients without underlying pulmonary or cardiac disease (e.g., respiratory distress syndrome, bronchopulmonary dysplasia, pulmonary edema, or chronic obstructive pulmonary disease), <b>one definitive</b> chest radiograph is acceptable</p>	<p>At least <b>three</b> of the following:</p> <ul style="list-style-type: none"> <li>• Fever (&gt;38.0°C or &gt;100.4°F) or hypothermia (&lt;36.0°C or &lt;96.8°F)</li> <li>• Leukopenia (&lt;4000 WBC/mm<sup>3</sup>) <b>or</b> leukocytosis (≥15,000 WBC/mm<sup>3</sup>)</li> <li>• New onset of purulent sputum, or change in character of sputum, or increased respiratory secretions, or increased suctioning requirements</li> <li>• New onset or worsening cough, or dyspnea, apnea, or tachypnea</li> <li>• Rales or bronchial breath sounds</li> <li>• Worsening gas exchange (e.g., O<sub>2</sub> desaturations [e.g., pulse oximetry &lt;94%], increased oxygen requirements, or increased ventilator demand)</li> </ul>

**Element Values**

- 1 Yes
- 2 No

**Additional Information**

- Onset of symptoms began after arrival to your ED/hospital.
- A diagnosis of pneumonia must be documented in the patient's medical record.
- Consistent with the January 2019 CDC defined VAP.

**Data Source Hierarchy Guide**


- 1 History and Physical
- 2 Physician's Notes
- 3 Progress Notes
- 4 Case Management/Social Services
- 5 Nursing Notes/Flow Sheet
- 6 Triage/Trauma Flow Sheet
- 7 Discharge Summary

**References to Other Databases**

- NTDS 2026

## Ventilator-Associated Pneumonia (VAP) PNU2 Bacterial of Filamentous Fungal Pathogens (pg. 1 of 2)

2026 NTDS Data Dictionary, Released July 2025




1

NOTE: In patients without underlying pulmonary or cardiac disease (for example: respiratory distress syndrome, bronchopulmonary dysplasia, pulmonary edema, or chronic obstructive pulmonary disease), one definitive chest imaging test result is acceptable.


## Ventilator-Associated Pneumonia (VAP) PNU2 Bacterial of Filamentous Fungal Pathogens (pg. 2 of 2)

2026 NTDS Data Dictionary, Released July 2025



## Ventilator-Associated Pneumonia (VAP) PNU2 Viral, Legionella, and other Bacterial Pneumonias (pg. 1 of 2)

2026 NTDS Data Dictionary, Released July 2025




**1** NOTE: In patients without underlying pulmonary or cardiac disease (for example: respiratory distress syndrome, bronchopulmonary dysplasia, pulmonary edema, or chronic obstructive pulmonary disease), one definitive chest imaging test result is acceptable.




## Ventilator-Associated Pneumonia (VAP) PNU2 Viral, Legionella, and other Bacterial Pneumonias (pg. 2 of 2)

2026 NTDS Data Dictionary, Released July 2025



## Ventilator-Associated Pneumonia (VAP) PNU3 Immunocompromised Patients (pg. 1 of 2)

2026 NTDS Data Dictionary, Released July 2025




1

NOTE: In patients without underlying pulmonary or cardiac disease (for example: respiratory distress syndrome, bronchopulmonary dysplasia, pulmonary edema, or chronic obstructive pulmonary disease), one definitive chest imaging test result is acceptable.


## Ventilator-Associated Pneumonia (VAP) PNU3 Immunocompromised Patients (pg. 2 of 2)

2026 NTDS Data Dictionary, Released July 2025



## Ventilator-Associated Pneumonia (VAP) ALTERNATE CRITERIA (PNU1), for infants $\leq 1$ -year-old (pg. 1 of 2)

2026 NTDS Data Dictionary, Released July 2025




1

NOTE: In patients without underlying pulmonary or cardiac disease (for example: respiratory distress syndrome, bronchopulmonary dysplasia, pulmonary edema, or chronic obstructive pulmonary disease), one definitive chest imaging test result is acceptable.


## Ventilator-Associated Pneumonia (VAP) ALTERNATE CRITERIA (PNU1), for infants $\leq 1$ -year-old (pg. 2 of 2)

2026 NTDS Data Dictionary, Released July 2025



## Ventilator-Associated Pneumonia (VAP) ALTERNATE CRITERIA (PNU1), for children >1-year-old or ≤12-years-old (pg. 1 of 2)

2026 NTDS Data Dictionary, Released July 2025



## Appendix A - Discharge Disposition Definitions

Element Value	Variable	Definition
2	Intermediate Care Facility (ICF)	A nursing home providing long-term care less than a skilled level, usually custodial care only.
7	Skilled Nursing Facility (SNF)	A nursing home or unit which provides skilled nursing or rehabilitation care, less than the level of an inpatient rehabilitation facility.
8	Hospice	A special way of caring for persons who are terminally ill. Hospice services can be provided in the home or at a nursing facility.
9	Inpatient Rehabilitation Facility (IRF)	A hospital or part of a hospital which provides intensive (3 hours per day) of rehabilitation therapies to persons with disability from recent injury or illness.
10	Long Term Acute Care Hospital (LTACH)	A special hospital or part of a hospital that provides treatment for patients who stay, on average, more than 25 days for extended acute care. Most patients are transferred from an intensive or critical care unit.

## Appendix B - Calculating ICU Length of Stay and Ventilator Days

Example #	Start Date	Start Time	Stop Date	Stop Time	LOS
A.	01/01/11	01:00	01/01/11	04:00	1 day (one calendar day)
B.	01/01/11	01:00	01/01/11	04:00	
	01/01/11	16:00	01/01/11	18:00	1 day (2 episodes within one calendar day)
C.	01/01/11	01:00	01/01/11	04:00	
	01/02/11	16:00	01/02/11	18:00	2 days (episodes on 2 separate calendar days)
D.	01/01/11	01:00	01/01/11	16:00	
	01/02/11	09:00	01/02/11	18:00	2 days (episodes on 2 separate calendar days)
E.	01/01/11	01:00	01/01/11	16:00	
	01/02/11	09:00	01/02/11	21:00	2 days (episodes on 2 separate calendar days)
F.	01/01/11	Unknown	01/01/11	16:00	1 day
G.	01/01/11	Unknown	01/02/11	16:00	2 days (patient was in ICU on 2 separate calendar days)
H.	01/01/11	Unknown	01/02/11	16:00	
	01/02/11	18:00	01/02/11	Unknown	2 days (patient was in ICU on 2 separate calendar days)
I.	01/01/11	Unknown	01/02/11	16:00	
	01/02/11	18:00	01/02/11	20:00	2 days (patient was in ICU on 2 separate calendar days)
J.	01/01/11	Unknown	01/02/11	16:00	
	01/03/11	18:00	01/03/11	20:00	3 days (patient was in ICU on 3 separate calendar days)
K.	Unknown	Unknown	01/02/11	16:00	
	01/03/11	18:00	01/03/11	20:00	Unknown (can't compute total)



## Appendix C - Glossary of Abbreviations

ACE	Angiotensin Converting Enzyme
ACS	Abdominal compartment syndrome; American College of Surgeons
ADL	Activities of daily living
AIS	Abbreviated Injury Scale
ARDS	Acute respiratory distress syndrome
ARF	Acute Renal Failure
BMI	Body mass index
BP	Blood pressure
CDC	Centers for Disease Control and Prevention
CHF	Congestive heart failure
CPAP/BIPAP	Continuous positive airway pressure/variable bi-level positive airway pressure
CT	Computerized topography
CVA	Cerebral vascular accident
DNR	Do not resuscitate
DNR-CC	Do not resuscitate; comfort care only
DNR-CCA	Do not resuscitate; comfort care arrest
DVT	Deep vein thrombosis
EOA	Esophageal Obturator Airway
ED	Emergency department
EMS	Emergency medical services
FAST	Focused assessment with sonography for trauma
FIPS	Federal Information Processing Standard codes
GCS	Glasgow Coma Score
ICD-9-CM	International Classification of Diseases, Ninth Revision, Clinical Modification
ICD-10-CM	International Classification of Diseases, Tenth Revision, Clinical Modification
IgG	Immunoglobulin G
ISS	Injury Severity Score
LMA	Laryngeal Mask Airway
MI	Myocardial infarction
MRI	Magnetic resonance imaging
NTDS	National Trauma Data Standard
OPO	Organ Procurement Organization
OR	Operating Room
OTR	Ohio Trauma Registry
PT	Prothrombin time
PTT	Partial thromboplastin time
PVD	Peripheral vascular disease
SaO <sub>2</sub>	Saturation of oxygen in arterial blood
TACR	Trauma Acute Care Registry
UB-04	Uniform Billing Form-04
XSD	XML (Extensible Markup Language) Schema definition

## Appendix D – Acronyms

- AIS: Abbreviated Injury Scale
- CDC: Centers for Disease Control
- CPR: cardiopulmonary resuscitation
- CT: computerized tomography
- ED: emergency department
- EMS: emergency medical service
- GCS: Glasgow Coma Scale
- ICD-10: International Classification of Diseases, Tenth Revision
- ICD-10-CA: International Classification of Diseases, Tenth Revision, Canada
- ICD-10-CM: International Classification of Diseases, Tenth Revision, Clinical Modification
- ICD-10-PCS: International Classification of Diseases, Tenth Revision, Procedure Coding System
- ICU: intensive care unit
- LOS: length of stay
- NA: not applicable
- NEMSIS: National Emergency Medical Services Information System
- NK/NR: not known/not recorded
- NTDB: National Trauma Data Bank
- NTDS: National Trauma Data Standard
- OR: operating room
- PACU: post-anesthesia care unit
- TQIP: Trauma Quality Improvement Program
- TQP: Trauma Quality Programs

### References to Other Databases

- NTDS 2026

## **Appendix E – Ohio Regional Trauma System Data Dictionary**

Ohio contains several regional trauma systems. These are organized, coordinated efforts in a defined geographic area that deliver the full range of care to all injured patients and work together with emergency services and disaster preparedness making efficient use of health care resources to improve patient outcomes in the state of Ohio. Membership in a regional trauma system is voluntary and not generally restricted by a facility's location.

This "Ohio Regional Data Dictionary" is an effort to collapse individual regional dictionaries into a single unified regional dictionary to improve state, regional and vendor responsiveness during the annual reconciliation with the changes issued by the American College of Surgeons (ACS).

It has been included as a reference in the State of Ohio Trauma Acute Care Registry's data dictionary and intended as a shared reference and data set common to all regional trauma systems. Specific questions about its contents should be directed to the regional trauma system to which you are a member.

If you are not a member of a regional trauma system, then you are not required to collect the items in these appendices. These items are for regional trauma system use only and should not be submitted to the state unless otherwise directed by the Division of EMS.

## Appendix F – Change LOG for the Ohio Trauma Acute Care Data (TACR)

### STATE OF OHIO CHANGES:

TACR INCLUSION/EXCLUSION CRITERIA	Additional Information	Added ***Initial hospital encounter is referring to the patient's first encounter for this injury event at your hospital.
TACR INCLUSION/EXCLUSION CRITERIA	Additional Information	<p>Added to patient inclusion criteria –</p> <ul style="list-style-type: none"> <li>• <b>NON-TRAUMA CENTERS:</b> S00, S10, S20, S30, S40, S50, S60, S70, S80, S90 (Patients with these isolated injuries that were transferred in/ out (FOR TRAUMA EVALUATION) or died) – Centers only capture patients transferred for trauma care. This would exclude patients being transferred to another non-trauma center/ acute care facility for medical purposes.</li> <li>• <b>TRAUMA CENTERS:</b> S00, S10, S20, S30, S40, S50, S60, S70, S80, S90 (Patients with these isolated injuries that were transferred in or died THAT WERE EVALUATED BY TRAUMA SERVICE) –Centers only capture patients transferred for trauma care.</li> </ul>
TACR INCLUSION/EXCLUSION CRITERIA	Additional Information	<p>Added to patient exclusion criteria –</p> <ul style="list-style-type: none"> <li>• <b>NON-TRAUMA CENTERS:</b> S00, S10, S20, S30, S40, S50, S60, S70, S80, S90 (Patients with these isolated injuries that were transferred in/ out (<b>NOT</b> FOR TRAUMA EVALUATION) or died) –Centers only capture patients transferred for trauma care. This would exclude patients being transferred to another non-trauma center/ acute care facility for medical purposes.</li> <li>• <b>TRAUMA CENTERS:</b> S00, S10, S20, S30, S40, S50, S60, S70, S80, S90 (Patients with these isolated injuries that were transferred in or died THAT WERE <b>NOT</b> EVALUATED BY TRAUMA SERVICE) –Centers only capture patients transferred for trauma care.</li> </ul>
Incident City Description	Description	CHANGED: Incident County is the county or parish where the incident occurred
Incident County	Description	CHANGED: Incident County is the county or parish where the incident occurred

DNR Status	Additional Information	CHANGED: DNR Status is to be collected at time of discharge from Hospital
Highest Activation	Additional Information	ADDED: Non-Trauma Center Facilities report the null value "Not Applicable" (NA)

### **NTDS CHANGES:**

Element Name	Change Location	Change Text
NTDS PATIENT INCLUSION CRITERIA	Text	CHANGED: *Exclude patient injuries sustained at your facility after initial ED/Hospital arrival and before hospital discharge, and all data associated with that injury event.
GENDER-AFFIRMING HORMONE THERAPY	Description	CHANGED: Is the patient currently (i.e., within the past 30 days) taking gender-affirming hormone therapy?
PROTECTIVE DEVICES	Element Value	CHANGED: 6. Child Restraint (child car seat, infant car seat, or child booster seat)
CHILD SPECIFIC RESTRAINT	Additional Information	ADDED: Report Element Value "1. Child Car Seat" for forward-facing child seats.
CHILD SPECIFIC RESTRAINT	Additional Information	ADDED: Report Element Value "2. Infant Car Seat" for rear-facing child seats.
OTHER TRANSPORT MODE	Data Source Hierarchy Guide	ADDED: Transfer Facility Records
INTER-FACILITY TRANSFER	Data Source Hierarchy Guide	ADDED: Transfer Facility Records
PRE-HOSPITAL CARDIAC ARREST	Data Source Hierarchy Guide	CHANGED: Transfer Facility Records
INTUBATION PRIOR TO ARRIVAL	Description	CHANGED: The patient is intubated with a definitive airway due to this injury prior to arrival at your hospital.
INTUBATION PRIOR TO ARRIVAL	Data Source Hierarchy Guide	ADDED: Transfer Facility Records
INTUBATION LOCATION	Data Source Hierarchy Guide	ADDED: Transfer Facility Records
TRAUMA SURGEON ARRIVAL DATE	Additional Information	CHANGED: Reported as YYYY-MM-DD.
TRAUMA SURGEON ARRIVAL TIME	Additional Information	CHANGED: Reported as HHMM.
PRIMARY MEDICAL EVENT	Description	CHANGED: The patient experienced a documented primary medical event (stroke, myocardial infarction, cardiac arrest, intracranial bleeding, sepsis) that immediately preceded the traumatic injury.
PRIMARY MEDICAL EVENT	Data Source Hierarchy Guide	ADDED: Transfer Facility Records
ADVANCE DIRECTIVE LIMITING CARE (Pre-Existing Condition)	Data Source Hierarchy Guide	ADDED: Transfer Facility Records
CURRENT SMOKER (Pre-Existing Condition)	Description	CHANGED: A patient who reports inhaling nicotine by smoking cigars, pipes, cigarettes, e-cigarettes, vaping, or juuling every day or some days within the last 30-days. EXCLUDE: Patients who chew tobacco or snuff.
CURRENT SMOKER (Pre-Existing Condition)	Additional Information	ADDED: Vaping and juuling includes vape pens, dab pens, dab rings, mods, pod-mods, or any other electronic delivery system used to inhale nicotine.

FUNCTIONALLY DEPENDENT HEALTH STATUS (Pre-Existing Condition)	Additional Information	ADDED: If Ventilator Dependence is Element Value "1. Yes," Functionally Dependent Health Status must be Element Value "1. Yes."
PERIPHERAL ARTERIAL DISEASE (Pre-Existing Condition)	Additional Information	CHANGED: A diagnosis of Peripheral Arterial Disease or Peripheral Vascular Disease must be documented in the patient's medical record.
VENTILATOR DEPENDENCE (Pre-Existing Conditions)	Additional Information	ADDED: If Ventilator Dependence is Element Value "1. Yes," Functionally Dependent Health Status must be Element Value "1. Yes."
DEEP SURGICAL SITE INFECTION (Hospital Event)	Additional Information	CHANGED: Consistent with the CDC January 2024 defined SSI.
UNPLANNED VISIT TO THE OPERATING ROOM (Hospital Event)	ELEMENT	RETIRED
UNPLANNED RETURN TO THE OPERATING ROOM (Hospital Event)	ELEMENT	NEW
ALL ELEMENTS	Element Intent	ADDED

# Ohio Regional Data Dictionary (ORDD) 2026

COTS, NORTN, NORTR, NOTS, SORTS, TRISTATE

## CAUSE CODE

---

### Element Intent

Utilized for reporting and regional validation processes

### Description

*Cause Code* is the code for the cause or mechanism of injury.

### Element Values

ANIMAL	Animal injury (includes bite and struck by)	INHAL	Inhalation
ASSAULT	Assault by person (blunt mechanism)	MACHINE	Machine
BIKE	Bicycle	MCC	Motorcycle Crash
BITING	Biting (human)	MVC	Motor Vehicle Crash
BURN	Burns (Chemical, Thermal, Electrical)	OV	Other Vehicle/ Off road (ATV, Animal, Golf Cart, Dirt bike)
CRUSH	Crush Injury	OVEREX	Overexertion
CUT	Cut	PED	Pedestrian
DROWN	Drowning/ Submersion	STAB	Stabbing/ Pierce/ Impalement
EXPOSURE	Exposure effects of heat or cold only	STRUCK	Struck by or against
FALL.SL	Fall same Level	SUFF	Suffocation/ Hanging/- Asphyxiation
FALL.MINOR	Fall < 10 feet (not same level fall)	WATERCRAFT	Watercraft
FALL.MAJOR	Fall > 10 feet	UNK	Unknown (Found down)
FALL.NFS	Fall NFS (unwitnessed fall)		
GSW	Gun Shot Wound		

### Common Null Values

- Not Accepted

### Additional Information

- The Primary E-Code assigned should correlate with the patient's cause code.
- See pages 15-16 of ORDD for additional clarifications
- Intention of the cause of injury will be captured using the software provided fields that include ICD-10 code intent classifications based off of the ICD-10 – Cause of Injury Matrix.
- The definition of a physical crush differs from that of the AIS injury definition of crush. Do not use AIS dictionary definition.

### Data Source Hierarchy Guide

- 1 EMS Run Sheet
- 2 Triage Form/Trauma Flow Sheet
- 3 ED Documentation



# INJURY DETAILS

---

## Element intent

Provides additional details of injury event, utilized for E-Code validation processes

## Description

*Injury Details* is a free text description that describes the circumstances of how the patient was injured.

## Element Values

- Relevant value for data element

## Additional Information

- Include as many details as possible
- Recommended examples:
  - 23- year old male, restrained driver, was T-boned by a tractor-trailer on the driver's side of the car, positive LOC, from Scene
  - 56- year old female fell down a flight of basement stairs and struck her head on the concrete floor, denies LOC, transfer by EMS from OSH

## Data Source Hierarchy Guide

- 1 EMS Run Sheet
- 2 Triage Form/Trauma Flow Sheet
- 3 ED Documentation

# TRAUMA TYPE

---

## Element Intent

Utilized for reporting and regional validation processes

## Description

*Trauma Type* is injury to human tissues or organs resulting from the transfer of energy from the environment to the human body, in which the human body lacks resilience to resist the energy transference. Trauma refers to critical injury that threatens life or permanent loss of function of a body part. There are five classifications of trauma, also referred to as trauma type. *Trauma Type* is the classification of the trauma.

## Element Values

A	Asphyxia
B	Blunt Trauma
P	Penetrating Trauma
TH	Thermal
OTHER	Other

## Common Null Values

- Not Accepted

## Additional Information

- Enter the trauma type which causes the highest injury severity
- *Penetrating Trauma*: Injury resulting from a projectile or thrust foreign object with perforation of tissues and underlying structures.
- *Blunt Trauma*: Injury secondary to a violent diffuse force that displaces tissues and or underlying structures.
- *Thermal*: Injury as a result of exposure to extreme temperatures of heat or cold, including chemical and electrical burns.
- *Asphyxia*: Injury as a result of inhalation, drowning, asphyxiation, hanging, strangulation, or suffocation.
- Other: Injury as a result of none of the above choices, such as overexertion resulting in injury
- Enter the injury type that causes the most serious injury as determined by the attending physician.

## Data Source Hierarchy Guide

- 1 EMS Run Sheet
- 2 Triage Form/Trauma Flow Sheet
- 3 ED Documentation
- 4 E-Code Matrix
- 5 Discharge Summary

# TRAUMA ACTIVATION LEVEL\*

---

## Element Intent

Utilized for reporting and regional validation processes

## Description

*Trauma Activation Level* is the highest level of trauma activation called for the patient when at your hospital emergency department.

## Element Values

- 1 Highest Level of Activation
- 2 Intermediate Level of Activation
- 3 Lowest Level of Activation (includes consults)
- 4 No Trauma Activation
- 5 Direct Admission

## Common Null Value

- Accepted

## Additional Information

- Enter a common null value of “Not Applicable” if your facility does not have a trauma service and is NOT a verified trauma center.
- Highest level of activation is defined by your hospital’s criteria.
- INCLUDE: patients who received the highest level of trauma activation initiated by emergency medical services (EMS) or by emergency department (ED) personnel at your hospital.
- INCLUDE: patients who received the highest level of trauma activation initiated by emergency medical services (EMS) or by emergency department (ED) personnel at your hospital and were downgraded after arrival to your center.
- INCLUDE: patients who received a lower level of trauma activation initiated by emergency medical services (EMS) or by emergency department (ED) personnel at your hospital and were upgraded to the highest level of trauma activation.
- EXCLUDE: patients who received the highest level of trauma activation after emergency department (ED) discharge.

## Data Source Hierarchy Guide

- 1 Triage/Trauma Flow Sheet
- 2 ED Record
- 3 History & Physical
- 4 Physician Notes
- 5 Discharge Summary

## References to Other Databases

- NTDS 2022
- Ohio Trauma Registry, Trauma Acute Care Registry 2022

*\*Element Values different than NTDS and OTR*

# ADMITTING SPECIALTY

---

## Element Intent

Utilized for reporting

## Description

*Admitting Specialty* is the medical specialty of the attending physician who admits the patient to your hospital.

## Element Values

0	Not Admitted (Died in your ED, transferred to another facility or discharged home)	13	Microvascular Surgery
1	General Adult Surgery	14	OBGYN Surgery
2	Neurosurgery	15	Ophthalmology
3	Orthopedic Surgery	16	Otolaryngology
4	General Pediatric Surgery	17	Urology
5	Burn Service	18	Intensivist/ Critical Care
6	Thoracic Surgery	19	Geriatrics
7	Plastic Surgery	20	Endocrinology
8	All Other Surgical Services	21	Trauma - Pediatric
9	All Other Non-Surgical Services	22	Trauma – Adult
10	Cardio Thoracic Surgery	23	Oral-Maxillofacial Surgery
11	Vascular Surgery	24	Pediatrics
12	Hand Surgery		

## Additional Information

- This is not necessarily the service to which the patient is designated upon admission to the hospital, but the medical specialty of the patient's attending physician
- #9, All Other Non-Surgical Services would include specified observation service line if it pertains to your facility

## Common Null Value

- Not Accepted

## Data Source Hierarchy Guide

- 1 ED Record
- 2 Trauma Flow Sheet
- 3 Billing/Registration Sheet
- 4 History & Physical

## PROCEDURE LOCATION

---

### Element Intent

Utilized for reporting

### Description

*Procedure Location* documents the location of the procedures performed while the patient was in your hospital.

### Element Values

- |   |                      |    |                               |
|---|----------------------|----|-------------------------------|
| 1 | Emergency Department | 6  | Other Specialty Area          |
| 2 | Operating Room       | 7  | Interventional radiology (IR) |
| 3 | ICU                  | 8  | Stepdown/Telemetry Unit       |
| 4 | Floor                | 9  | Observation Unit              |
| 5 | Radiology            | 10 | Post Anesthesia Care Unit     |

### Additional Information

- Include only those procedures performed at your hospital.
- This field is linked to the *Hospital Procedures* Field
- Other Specialty Area includes: Endo, cardiac cath lab, dialysis, etc.
- Scene Procedures are not part of this data variable but are collected elsewhere

### Common Null Value

- Accepted

### Data Source Hierarchy Guide

- 1 Operative Reports
- 2 Procedure Notes
- 3 ED and ICU Records
- 4 Trauma Flow Sheet
- 5 Nursing Notes
- 6 Radiology Reports
- 7 Anesthesia Record
- 8 Billing Sheet/Medical Records Coding Summary Sheet
- 9 Hospital Discharge Summary

## HOSPITAL PROCEDURE CODE

### Element Intent

Utilized for reporting and regional validation processes

### Description

*Hospital Procedure Code* is all operative or essential procedures conducted on the patient during his/her stay at your hospital.

### Element Values

- All values for data element

At minimum:

<b>AGRAM</b>	<b>Arteriograms (includes angiography)</b>	<b>FASC</b>	<b>Fasciotomy</b>
<b>ANGIOEMB</b>	<b>Angioembolization</b>	<b>FAST</b>	<b>FAST Exam (includes diagnostic Ultrasound)*</b>
<b>BRONCH</b>	<b>Bronchoscopy*</b>	<b>GAST</b>	<b>Gastrostomy/ jejunostomy (percutaneous or endoscopic)</b>
<b>PRBC</b>	<b>Packed Red Blood Cells*</b>	<b>GASTJEJ</b>	<b>Percutaneous (endoscopic) gastrojejunostomy</b>
<b>FFP</b>	<b>Fresh Frozen Plasma*</b>	<b>ICP</b>	<b>Intracranial pressure monitor insertion*</b>
<i>MASS</i>	<i>Massive Transfusion</i>	<b>INTUB</b>	<b>Insertion of Oral intubation Tube*</b>
<b>PLAT</b>	<b>Platelets*</b>	<b>IVC</b>	<b>IVC filter</b>
<i>CRANI</i>	<i>Craniotomy</i>	<i>MRIBRAIN</i>	<i>MRI Brain</i>
<b>OXYMON</b>	<b>Cerebral oxygen monitoring*</b>	<i>MRISPINE</i>	<i>MRI Spine</i>
<b>CHEST</b>	<b>Chest Tube insertion/Thoracostomy*</b>	<i>MRIOther</i>	<i>MRI Other</i>
<b>CLRD</b>	<b>Closed reduction of Fracture</b>	<i>NONE</i>	<i>None</i>
<b>CPR</b>	<b>CPR</b>	<i>NEEDLE</i>	<i>Needle decompression</i>
<b>CTABD</b>	<b>CT Scan Abdomen*</b>	<b>OCM</b>	<b>Open cardiac massage</b>
<b>CTCHEST</b>	<b>CT Scan Chest*</b>	<i>OTHER</i>	<i>Other unspecified procedure</i>
<i>CTFACE</i>	<i>CT Scan Face*</i>	<b>REBOA</b>	<b>Resuscitative Endovascular Balloon Occlusion of Aorta</b>
<b>CTHEAD</b>	<b>CT Scan Head*</b>	<b>SUPCYST</b>	<b>Suprapubic cystostomy</b>
<b>CTPELVIS</b>	<b>CT Scan Pelvis*</b>	<i>THORC</i>	<i>Open Thoracotomy</i>
<b>CTSPINE.C</b>	<b>CT Scan Spine – Cervical*</b>	<b>TRACH</b>	<b>Tracheostomy</b>
<b>CTSPINE.T</b>	<b>CT Scan Spine – Thoracic*</b>	<b>TRACT</b>	<b>Skeletal and halo traction</b>
<b>CTSPINE.L</b>	<b>CT Scan Spine – Lumbar*</b>	<b>URSTNT</b>	<b>Ureteric catheterization (Ureteric stent)</b>
<b>DEBRD</b>	<b>Soft tissue/ bony debridement*</b>	<b>VENT</b>	<b>Mechanical Ventilation (continuous)*</b>
<b>DOPPLER</b>	<b>Doppler of Extremities*</b>	<b>VENTRIC</b>	<b>Ventriculostomy</b>
<b>ENDO</b>	<b>Endoscopy (includes gastroscopy, sigmoidoscopy, colonoscopy)</b>	<b>XRAY</b>	<b>Plain radiography (whole body, whole skeleton, infant whole body)</b>

**Bolded element value indicates NTDS required data element**

*Italicized element value indicates ORDD required data element*

See additional information

## **Additional Information**

- Operative and/or essential procedures are defined as procedures performed in the Operating Room, Emergency Department, and/or Intensive Care Unit that were essential to the diagnoses, stabilization, or treatment of the patient's specific injuries or their complications at your hospital.
- Include only procedures performed at your hospital.
- At a minimum, the procedures listed should be captured. The hospital may choose to capture additional procedures for internal use. Procedures included in the Procedures List that are designated with an asterisk have the potential to be performed multiple times during one episode of hospitalization. In this case, capture only the first event. If there is no asterisk, capture each event even if there is more than one.
- FAST is defined as a rapid bedside ultrasound examination 'Focused Assessment with Sonography for Trauma'
- XRAY – If facility chooses to collect additional x-ray procedures, report as "OTHER – Other unspecified procedure"
- See page 17 of ORDD for additional clarifications

## **Data Source Hierarchy Guide**

- 1 Operative Reports
- 2 ED and ICU Records
- 3 Trauma Flow Sheet
- 4 Anesthesia Record
- 5 Billing Sheet/Medical Records Coding Summary Sheet
- 6 Hospital Discharge Summary

## INJURY DIAGNOSES DESCRIPTION

---

### Element intent

Provides additional details of injuries, utilized for validation processes

### Description

*Injury Diagnoses Description* is a free text element of the patient's description for all injuries identified at your ED/hospital for this injury event that match the corresponding ICD-10 assigned. Diagnoses must be confirmed by a physician at your facility.

### Element Values

- Relevant Value for Data Element

### Additional Information

- Provide detailed information of injury
  - Example: Right femur fx, comminuted and displaced
  - Scalp laceration, 7 cm
  - SAH with coma > 6 hours
- Can be utilized to generate Abbreviated Injury Score and Injury Severity Score
- The maximum number of diagnoses that may be reported for an individual patient is 50
- In DI system this would be completed in the narrative

### Common Null Values

- Accepted

### Data Source Hierarchy Guide

- 1 Autopsy Report
- 2 Operative Report
- 3 Discharge Summary
- 4 Trauma Flow Sheet
- 5 Radiology Results
- 6 Billing Sheet/Medical Records Coding Summary Sheet
- 7 ED and ICU Records



# ISS BODY REGION

---

## Element Intent

Utilized for reporting and regional validation processes

## Description

*ISS Body Region* is the Injury Severity Score assigned by body region codes that reflects the patient's injury(ies) diagnosed at your ED/hospital for this injury event.

## Element Values

- 1 Head or Neck
- 2 Face
- 3 Chest
- 4 Abdominal or Pelvic Contents
- 5 Extremities or Pelvic Girdle
- 6 External

## Additional Information

- Field value #1, *Head or Neck*, includes injury to the brain, skull, cervical spine and/or cervical spine fractures
- Field value #2, *Face*, includes those areas involving the mouth, ears, nose and/or facial bones
- Field value #3, *Chest*, includes all lesions to internal organs within the chest, diaphragm, rib cage and/or thoracic spine
- Field value #4, *Abdominal or Pelvic Contents*, includes all lesions to internal organs within the abdomen and lumbar spine
- Field value #5, *Extremities or Pelvic Girdle*, includes sprains, dislocations, fractures and amputations ***except for the spinal column, skull and rib cage***
- Field value #6, *External*, includes injuries such as lacerations, contusions, abrasions and burns independent of their location on the body surface

## Common Null Values

- Accepted

## Data Source Hierarchy Guide

- 1 Autopsy Report
- 2 Operative Report
- 3 Discharge Summary
- 4 Trauma Flow Sheet
- 5 Radiology Results
- 6 Billing Sheet/Medical Records Coding Summary Sheet
- 7 ED and ICU Records

## LENGTH OF STAY

---

### Element Intent

Utilized for reporting

### Description

*Length of Stay documents* the total number of days that the patient occupied a bed while in your hospital.

### Element Values

- Relevant value for data element

### Additional Information

- This field is calculated from data in the “Hospital Arrival Date” and “Discharge Date” fields, automatically.
- Recorded in full day increments with any partial calendar day counted as a full calendar day.

### Data Source Hierarchy Guide

- 1 Registration Form
- 2 Discharge Form

# INJURY MECHANISM DEFINITION

## APPENDIX E

Mechanism	Definition
ANIMAL	Animal Injury (Including but not limited to, bite, struck by) (rider of animal – see OV)
ASSAULT	Assault by person (blunt mechanism only)
BIKE	Any accident involving a bicyclist
BITING	Human bite only
BURN	Burn – Chemical, Thermal, Electrical, or Other
CRUSH	Crushing mechanism (when force or pressure is put onto the body)
CUT	Cut (includes cut by machine or chainsaw)
DROWN	Drowning/ Submersion
EXPOSURE	Exposure effects of heat and cold only Primary injury – environmental, not treatment related or sequela (AIS2015)
FALL.SL	Any fall from standing (feet on ground), may include subsequent strike against object
FALL.MINOR	Any fall <10 feet that is not a same level fall
FALL.MAJOR	Any fall >= 10 feet.
FALL.NFS	Use only if no details stated about the fall, unwitnessed fall
GSW	Gunshot wound
INHAL	Smoke or chemical inhalation type injuries
MACHINE	Injury caused by machinery
MCC	Motorcycle related injuries (includes 2 wheel motor scooter and electric bike)
MVC	Cars, trucks, vans, SUV's on roads or parking lots etc.
OV	All off road and other vehicles not included elsewhere. (ATVs, snowmobiles, riding lawnmowers, 4-wheelers, golf carts, rider of animal, Dirt bike etc.)
OVEREX	Overexertion injuries (ie. heavy lifting). To be used for strains and sprains from improper lifting or repetitive motion. Do not include stress fractures here.
PED	Person walking (or using their typical mode of mobility) struck by motor vehicle. (If a person uses a wheelchair, mobility scooter, or other such conveyance to get around, they are still considered a pedestrian though they are not walking)
STAB	Stabbing/ Pierce/ Impalement
STRUCK	Struck by or against a person or object (not intentionally by someone)
SUFF	Suffocation, Hanging, or Asphyxiation
WATERCRAFT	Injury involving any boat (including jet skis), to include anything pulled behind watercraft (water skis, inner tubes, etc.)
UNK	Unknown (Found Down)

## INJURY MECHANISM REFERENCES

### Height Categories

**House Story Height:** The average height of 1, 2, and 3 story houses are respectively 10, 20, and 30 feet. A good rule of thumb when calculating the height of a house is to allocate approximately 10 feet per floor plus the height of the roof.

(www.weekendbuilds.com)

**Commercial building height:** Average commercial building single story roof equals 12 feet

**Office/ skyscraper building height:** Building story height varies widely from just under the 14-foot average to well above it.

That's because several factors affect the story's-to-feet standard for a building: building material, insulation type, and different height allowances for different levels. (theskydeck.com)

**Step height:** Standard should be between 7 or 7 ¾ inches at the most, and no less than 4 inches (www.stairsupplies.com)

## ICD-10-PCS CODING REFERENCE

<b>DIAGNOSTIC AND THERAPEUTIC IMAGING</b>		
Computerized Tomography of Head	BW28	
Computerized Tomography of Chest	BW24	If wanting to combine CHEST/ABD/PELV- BW25
Computerized Tomography of Abdomen	BW20	If wanting to combine ABD/PELV- BW21
Computerized Tomography of Pelvis	BW2G	
Computerized Tomography of C-Spine	BR20	
Computerized Tomography of T-Spine	BR27	
Computerized Tomography of L-Spine	BR29	
Computerized Tomography Angiography	See Computerized Tomography Codes	
Doppler Ultrasound of extremities	RLE- B54B	
	LLE- B54C	
	BLE- B54D	
	RUE- B54M	
	LUE- B54N	
	BUE- B54P	
Diagnostic ultrasound (Includes FAST)	BW40ZZZ (abdomen), BW4GZZZ (pelvic), B24CZZZ (pericardium), BB4CZZZ (mediastinum)	
Angioembolization	Head, neck, upper arm artery- 03L	
	Aorta, abdominal, leg arteries- 04L	
	Head, neck, upper arm vein 05L	
	Leg veins- 06L	
Angiography	Upper arteries- B30	
	Lower arteries- B40	
IVC Filter	06H03DZ	
REBOA	Heart & Great Vessels- 02L	
	Lower Arteries- 04L	
Plain Radiography of Adult whole body	BW0KZZZ	
Plain Radiography of whole skeleton	BW0LZZZ	
Plain Radiography of infant whole body	BW0MZZZ	
<b>CARDIOVASCULAR</b>		
Open Cardiac Massage	02QA0ZZ	
CPR	5A12012 (manual) 5A1221J (mechanical)	

<b>CNS</b>		
Insertion of ICP Monitor	4A103BD	
Ventriculostomy	009630Z	
Cerebral Oxygen Monitor	4A103RD	
<b>GENITOURINARY</b>		
Ureteric Catheterization (Ureteric Stent)	0T7	
Suprapubic Cystostomy	0T1B0ZD	
<b>MUSCULOSKETETAL</b>		
Soft Tissue/Bony Debridement	Excisional Tissue Debridement- OJB	
	Excisional Debridement- Upper Bone- OPB	
	Excisional Debridement- Lower Bone- OQB	
	Non Excisional Tissue Debridement- OJD	
	Non Excisional Debridement- Upper Bone- OPD	
	None Excisional Debridement- Lower Bone- OQD	
Closed Reduction of Fractures	Upper Bones- OPS	
	Lower Bones- OQS	
Skeletal and Halo Traction	2W6	
Fasciotomy	0J8	
<b>TRANSFUSION</b>		
Transfusion of red blood cells (first 24 hours)	302_3N1	
Transfusion of platelets (first 24 hours)	302_3R1	
Transfusion of plasma (first 24 hours)	302_3K1	
<b>RESPIRATORY</b>		
Insertion of endotracheal tube	0BH17EZ	
Continuous mechanical ventilation	<24 HR- 5A1935Z	
	24-96HR- 5A1945Z	
	>96HR- 5A1955Z	
Chest Tube/ Thoracostomy	R- 0W9930Z	
	L- 0W9B30Z	
Bronchoscopy	Inspection- 0BJ	
	Drainage- 0B9	
	Excision- 0BB	
Thoracotomy	R- 0W99	
	L- 0W9B	
Tracheostomy	0B110F4	

<b>GASTROINTESTINAL</b>		
Endoscopy (includes gastroscopy, sigmoidoscopy, colonoscopy)	Gastroscopy- 0DJ68ZZ	
	Sigmoidoscopy- 0DJD8ZZ	
	Colonoscopy- 0DJD8ZZ	
Gastrostomy/jejunostomy (percutaneous or endoscopic)	Bypass- 0D1	
	Drainage- 0D9	
Percutaneous (endoscopic) gastrojejunoscopy	0DJ	

## CHANGE LOG

### 2026 CHANGES

Scene Delay	Retired
Cause Code – Sport	Retired
Cause Code – OV	Added Dirt bike to description