

---

# Ohio Medicaid

Pharmacy Benefit Management Program

---



## Unified Preferred Drug List

Medicaid Fee-for-Service  
and Managed Care Plans

Effective July 1, 2025

---

## Helpful Links

---

### Prior Authorization (PA)

[Prior Authorization \(PA\) Information | medicaid.ohio.gov](#)

- General Prior Authorization Requirements
- PA and Step Therapy Frequently Asked Questions (FAQ)

### Unified Preferred Drug List (UPDL)

[Ohio Unified Preferred Drug List | medicaid.ohio.gov](#)

- Unified Preferred Drug List (UPDL)

---

## General Information

---

- The Statewide UPDL is not an all-inclusive list of drugs covered by the Ohio Department of Medicaid (ODM). Non UPDL drugs receive PA in accordance with the Gainwell SPBM medical necessity policy as posted on the Gainwell SPBM website.
  - Medications that are new to the market will be non-preferred, PA required, until reviewed by the ODM Pharmacy and Therapeutics (P&T) Committee.
  - The UPDL document is organized by therapeutic class. Brand name drugs are listed in CAPITAL letters; generic drug names are listed in lower case letters. In most cases, when a generic of a brand-name drug is available, the generic drug will be preferred and appear on the UPDL while the brand name will be non-preferred but not appear on the UPDL. The [Drug Search tool](#) is a handy reference to check the status of a drug. Some generic drugs may require a specific labeler, or the brand to be dispensed.
  - ODM will only cover drugs that are part of the Medicaid Drug Rebate Program, with limited exceptions. This document may not reflect the most current rebate status of a drug (i.e., a drug may be listed on the document but is non-rebateable and therefore non-payable).
  - Some therapeutic categories are deemed 'legacy' categories. These categories are denoted with an "\*" and LEGACY CATEGORY listed next to their title on the table on contents and their place within the criteria document. Legacy is defined as: Patients who have a claim for a non-preferred drug in the previous 120 days will be automatically approved to continue the drug. Patients who have taken the drug previously, but do not have claims history (e.g., new to Medicaid), will need to submit a prior authorization to continue coverage.
  - ALL authorizations must be prescribed in accordance with FDA approved labeling or listed on a CMS-supported compendia.
  - For ALL authorizations, there must be a trial and failure of preferred strengths prior to authorization of non-preferred strengths (if available).
-

- 
- For ALL non-preferred authorizations, there must be documentation of medical necessity beyond convenience for why the patient cannot be changed to a preferred drug form (i.e., allergies, drug-drug interactions, contraindications, or intolerances). Must have had an inadequate clinical response of preferred individual components for any combination non-preferred product.
  - For any nonsolid oral dosage formulation, there must be documentation of medical necessity for why patient cannot be changed to a solid oral dosage formulation (if available).
  - For non-preferred extended-release formulations, there must be documentation of an inadequate clinical response with its immediate release formulation (if available).
  - For non-preferred brand names that have preferred generics, there must be documentation of an inadequate clinical response or allergy to two or more generic labelers (if available).
  - For ALL subsequent authorizations, there must be documentation of patient's clinical response to treatment and ongoing safety monitoring unless otherwise stated.
  - Some therapeutic categories have sub-sections to divide the medications by their mechanism of action, route of administration, or duration of action. References to 'sub-section' in the Clinical Criteria shall be defined as the separate groupings that appear in that category's drug placement columns.
  - Some therapeutic categories may have quantity limits on specific drugs. For a list of the quantity limits on specific drugs, please reference the Quantity Limit Document found here: [Quantity Limits Document | spbm.medicaid.ohio.gov](https://spbm.medicaid.ohio.gov/QuantityLimitsDocument)
- 

---

## Terminology/Abbreviations:

---

**AR** (Age Restriction) – An edit allowing claims for members within a defined age range to be covered without PA

**BvG** (Brand Preferred Over the Generic) – The brand name drug is preferred over the generic equivalent

**PA** (Clinical Prior Authorization) – PA is required before the drug will be covered

**ST** (Step Therapy) – Drug requires a trial with one or more preferred drugs before being covered

---

## UPDL Format

- With a few exceptions, the clinical criteria have a cumulative top-to-bottom format.

Example Category		
PREFERRED AGENTS	NON-PREFERRED AGENTS	PA CRITERIA
Example Drug	Example Drug	<p><b><u>LENGTH OF AUTHORIZATIONS:</u></b> X days or Initial: X days; Subsequent: X days (if different)</p> <p><b><u>LEGACY*:</u></b> Patients who have a claim for a non-preferred drug in the previous 120 days will be automatically approved to continue the drug. Patients who have taken the drug previously, but do not have claims history (e.g. new to Medicaid), will need to submit a prior authorization in order to continue coverage.</p> <p><b><u>CLINICAL PA CRITERIA (if applicable):</u></b></p> <p><b><u>“DRUG” CRITERIA (if applicable):</u></b></p> <p><b><u>STEP THERAPY CRITERIA:</u></b></p> <ul style="list-style-type: none"> <li>• Must have had an inadequate clinical response of at least <u>X days</u> with at least <u>X preferred</u> drugs</li> </ul> <p><b><u>NON-PREFERRED CRITERIA:</u></b></p> <ul style="list-style-type: none"> <li>• Must have had an inadequate clinical response of at least <u>X days</u> with <u>X preferred</u> drugs</li> </ul> <p><b><u>ADDITIONAL “DRUG” CRITERIA (if applicable):</u></b></p> <p><b><u>ADDITIONAL INFORMATION (if applicable):</u></b></p> <p><b><u>SUBSEQUENT AUTHORIZATION CRITERIA:</u></b></p> <ul style="list-style-type: none"> <li>• Must provide documentation of patient’s response to treatment from baseline and/or attestation of clinical stabilization</li> </ul> <p><b>AR</b> – a PA is required for patients X years and older OR younger than X years</p>

---

## Interpretation of UPDL Format

---

- The UPDL criteria is designed to have a cumulative approach from top-to-bottom. The following scenarios will aid in illustrating this point:

### **Scenario 1: Clinical PA drug**

- All Authorizations
- Clinical PA Criteria

### **Scenario 2: Clinical PA drug with drug-specific criteria**

- All Authorizations
- Drug-Specific Criteria

### **Scenario 3: Step-therapy drug**

- All Authorizations
- Clinical PA Criteria (if applicable)
- Step Therapy Criteria

### **Scenario 4: Non-preferred drug**

- All Authorizations
- Clinical PA Criteria (if applicable)
- Step Therapy Criteria (if applicable)
- Non-Preferred Criteria

### **Scenario 5: Non-preferred drug with drug-specific criteria**

- All Authorizations
- Clinical PA Criteria (if applicable)
- Step Therapy Criteria (if applicable)
- Non-Preferred Criteria
- Additional Drug-Specific Criteria

## Table of Contents

<b>Example Category.....</b>	<b>4</b>
<b>Analgesic Agents: Gout .....</b>	<b>10</b>
<b>Analgesic Agents: NSAIDS .....</b>	<b>11</b>
<b>Analgesic Agents: Opioids .....</b>	<b>12</b>
<b>Blood Formation, Coagulation, and Thrombosis Agents: Colony Stimulating Factors .....</b>	<b>17</b>
<b>Blood Formation, Coagulation, and Thrombosis Agents: Hematopoietic Agents .....</b>	<b>18</b>
<b>Blood Formation, Coagulation, and Thrombosis Agents: Hemophilia A, von Willebrand Disease, and Factor XIII Deficiency* LEGACY CATEGORY.....</b>	<b>19</b>
<b>Blood Formation, Coagulation, and Thrombosis Agents: Hemophilia B* LEGACY CATEGORY .....</b>	<b>20</b>
<b>Blood Formation, Coagulation, and Thrombosis Agents: Heparin-Related Preparations .....</b>	<b>21</b>
<b>Blood Formation, Coagulation, and Thrombosis Agents: Oral Anticoagulants .....</b>	<b>22</b>
<b>Blood Formation, Coagulation, and Thrombosis Agents: Oral Antiplatelet .....</b>	<b>23</b>
<b>Cardiovascular Agents: Angina, Hypertension and Heart Failure .....</b>	<b>24</b>
<b>Cardiovascular Agents: Antiarrhythmics .....</b>	<b>27</b>
<b>Cardiovascular Agents: Lipotropics.....</b>	<b>28</b>
<b>Cardiovascular Agents: Pulmonary Arterial Hypertension* LEGACY CATEGORY.....</b>	<b>30</b>
<b>Central Nervous System (CNS) Agents: Alzheimer's Agents* LEGACY CATEGORY .....</b>	<b>31</b>
<b>Central Nervous System (CNS) Agents: Anti-Migraine Agents, Acute.....</b>	<b>32</b>
<b>Central Nervous System (CNS) Agents: Anti-Migraine Agents, Cluster Headache .....</b>	<b>33</b>
<b>Central Nervous System (CNS) Agents: Anti-Migraine Agents, Prophylaxis .....</b>	<b>34</b>
<b>Central Nervous System (CNS) Agents: Anticonvulsants* LEGACY CATEGORY .....</b>	<b>35</b>
<b>Central Nervous System (CNS) Agents: Anticonvulsants Rescue .....</b>	<b>37</b>
<b>Central Nervous System (CNS) Agents: Antidepressants* LEGACY CATEGORY .....</b>	<b>38</b>
<b>Central Nervous System (CNS) Agents: Attention Deficit Hyperactivity Disorder Agents .....</b>	<b>40</b>
<b>Central Nervous System (CNS) Agents: Atypical Antipsychotics* LEGACY CATEGORY .....</b>	<b>41</b>

Central Nervous System (CNS) Agents: Fibromyalgia Agents.....	43
Central Nervous System (CNS) Agents: Medication Assisted Treatment of Opioid Addiction.....	44
Central Nervous System (CNS) Agents: Movement Disorders.....	45
Central Nervous System (CNS) Agents: Multiple Sclerosis* LEGACY CATEGORY .....	46
Central Nervous System (CNS) Agents: Narcolepsy .....	47
Central Nervous System (CNS) Agents: Neuropathic Pain .....	48
Central Nervous System (CNS) Agents: Parkinson's Agents .....	49
Central Nervous System (CNS) Agents: Restless Legs Syndrome .....	50
Central Nervous System (CNS) Agents: Sedative-Hypnotics, Non-Barbiturate.....	51
Central Nervous System (CNS) Agents: Skeletal Muscle Relaxants, Non-Benzodiazepine.....	52
Central Nervous System (CNS) Agents: Smoking Deterrents.....	53
Dermatologic Agents: Oral Acne Products .....	54
Dermatologic Agents: Topical Acne Products .....	55
Endocrine Agents: Androgens.....	56
Endocrine Agents: Diabetes – Hypoglycemia Treatments .....	57
Endocrine Agents: Diabetes – Insulin .....	58
Endocrine Agents: Diabetes – Non-Insulin .....	60
Endocrine Agents: Endometriosis.....	62
Endocrine Agents: Estrogenic Agents .....	63
Endocrine Agents: Growth Hormone.....	64
Endocrine Agents: Osteoporosis – Bone Ossification Enhancers .....	65
Endocrine Agents: Progestin Agents.....	66
Endocrine Agents: Uterine Fibroids .....	67
Gastrointestinal Agents: Anti-Emetics.....	68
Gastrointestinal Agents: Bowel Preparations .....	69

<b>Gastrointestinal Agents: Crohn's Disease .....</b>	<b>70</b>
<b>Gastrointestinal Agents: Hepatic Encephalopathy .....</b>	<b>71</b>
<b>Gastrointestinal Agents: Irritable Bowel Syndrome (IBS) with Diarrhea .....</b>	<b>72</b>
<b>Gastrointestinal Agents: Pancreatic Enzymes .....</b>	<b>73</b>
<b>Gastrointestinal Agents: Proton Pump Inhibitors .....</b>	<b>74</b>
<b>Gastrointestinal Agents: Ulcerative Colitis.....</b>	<b>75</b>
<b>Gastrointestinal Agents: Unspecified GI.....</b>	<b>76</b>
<b>Genitourinary Agents: Benign Prostatic Hyperplasia.....</b>	<b>77</b>
<b>Genitourinary Agents: Electrolyte Depleter Agents.....</b>	<b>78</b>
<b>Genitourinary Agents: Urinary Antispasmodics .....</b>	<b>79</b>
<b>Hyperkalemia Agents: Potassium Binders.....</b>	<b>80</b>
<b>Immunomodulator Agents: Systemic Inflammatory Disease .....</b>	<b>81</b>
<b>Infectious Disease Agents: Antibiotics – Cephalosporins .....</b>	<b>83</b>
<b>Infectious Disease Agents: Antibiotics – Inhaled .....</b>	<b>84</b>
<b>Infectious Disease Agents: Antibiotics – Macrolides .....</b>	<b>85</b>
<b>Infectious Disease Agents: Antibiotics – Quinolones.....</b>	<b>86</b>
<b>Infectious Disease Agents: Antibiotics – Tetracyclines .....</b>	<b>87</b>
<b>Infectious Disease Agents: Antifungals .....</b>	<b>88</b>
<b>Infectious Disease Agents: Antivirals – Hepatitis C Agents .....</b>	<b>89</b>
<b>Infectious Disease Agents: Antivirals – Herpes .....</b>	<b>90</b>
<b>Infectious Disease Agents: Antivirals – HIV* LEGACY CATEGORY .....</b>	<b>91</b>
<b>Ophthalmic Agents: Antibiotic and Antibiotic-Steroid Combination Drops and Ointments.....</b>	<b>93</b>
<b>Ophthalmic Agents: Antihistamines &amp; Mast Cell Stabilizers .....</b>	<b>94</b>
<b>Ophthalmic Agents: Dry Eye Treatments .....</b>	<b>95</b>
<b>Ophthalmic Agents: Glaucoma Agents.....</b>	<b>96</b>



<b>Ophthalmic Agents: NSAIDs</b> .....	97
<b>Ophthalmic Agents: Ophthalmic Steroids</b> .....	98
<b>Otic Agents: Antibacterial and Antibacterial/Steroid Combinations</b> .....	99
<b>Respiratory Agents: Antihistamines – Second Generation</b> .....	100
<b>Respiratory Agents: Cystic Fibrosis</b> .....	101
<b>Respiratory Agents: Epinephrine</b> .....	102
<b>Respiratory Agents: Hereditary Angioedema</b> .....	103
<b>Respiratory Agents: Inhaled Agents</b> .....	104
<b>Respiratory Agents: Leukotriene Receptor Modifiers &amp; Inhibitors</b> .....	106
<b>Respiratory Agents: Monoclonal Antibodies-Anti-IL/Anti-IgE</b> .....	107
<b>Respiratory Agents: Nasal Preparations</b> .....	108
<b>Respiratory Agents: Pulmonary Fibrosis</b> .....	109
<b>Topical Agents: Antifungals</b> .....	110
<b>Topical Agents: Antiparasitics</b> .....	111
<b>Topical Agents: Corticosteroids</b> .....	112
<b>Topical Agents: Immunomodulators</b> .....	113

### Analgesic Agents: Gout

PREFERRED AGENTS	NON-PREFERRED AGENTS	PA CRITERIA
allopurinol 100, 300mg colchicine tab probenecid probenecid/colchicine	allopurinol 200mg febuxostat MITIGARE <sup>BvG</sup>	<p><b><u>LENGTH OF AUTHORIZATIONS:</u></b> 365 days</p> <p><b><u>NON-PREFERRED CRITERIA:</u></b></p> <ul style="list-style-type: none"> <li>Must have had an inadequate clinical response of at least <u>30 days</u> with at least <u>one preferred</u> drug in this UPDL category and indicated for diagnosis</li> </ul> <p><b><u>ADDITIONAL COLCHICINE CAPSULE (MITIGARE) CRITERIA:</u></b></p> <ul style="list-style-type: none"> <li>Must have had an inadequate clinical response of <u>30 days</u> with colchicine tablets</li> </ul>

### Analgesic Agents: NSAIDs

PREFERRED AGENTS	NON-PREFERRED AGENTS	PA CRITERIA
celecoxib diclofenac sodium DR, ER, gel 1% etodolac IR, ER flurbiprofen ibuprofen indomethacin IR, ER cap ketorolac mefenamic acid meloxicam tab nabumetone naproxen IR naproxen susp <sup>AR</sup> oxaprozin piroxicam sulindac	diclofenac/misoprostol diclofenac patch 1.3%; soln 1.5%, 2% diclofenac potassium ELYXYB fenoprofen ibuprofen/famotidine indomethacin supp, susp ketoprofen IR, ER meclofenamate meloxicam cap naproxen EC, ER naproxen/esomeprazole RELAFEN DS	<p><b><u>LENGTH OF AUTHORIZATIONS:</u></b> 365 days</p> <p><b><u>NON-PREFERRED CRITERIA:</u></b></p> <ul style="list-style-type: none"> <li>Must have had an inadequate clinical response of at least <u>30 days</u> with at least <u>two preferred</u> drugs in this UPDL category and indicated for diagnosis</li> </ul> <p><b>AR</b> – naproxen susp: a PA is required for patients 12 years old and older</p>

## Analgesic Agents: Opioids

PREFERRED AGENTS	NON-PREFERRED AGENTS	PA CRITERIA
SHORT-ACTING		<p><b>**Ohio law requires prescribers to request and review an OARRS report before initially prescribing or personally furnishing any controlled substance, such as an opioid analgesic or a benzodiazepine, and gabapentin**</b></p> <p><b><u>LENGTH OF AUTHORIZATIONS:</u></b> Initial short-acting and long-acting requests may only be authorized for up to 90 days. For reauthorization, up to 180 days.</p> <p><b><u>BUPRENORPHINE TOPICAL (BUTRANS) CRITERIA:</u></b></p> <ul style="list-style-type: none"><li>For doses greater than 5 mcg/hour must provide documentation of an inadequate clinical response with at least one opioid formulation taken for at least 30 of the last 60 days</li></ul> <p><b><u>MORPHINE SULFATE ER (MS CONTIN) CRITERIA:</u></b></p> <ul style="list-style-type: none"><li>Unless receiving for cancer pain, palliative care, or end-of-life/hospice care, must provide documentation of an inadequate clinical response with at least one opioid formulation taken for at least 30 of the last 60 days</li><li>Must also meet LONG-ACTING OPIOID CRITERIA</li></ul> <p><b><u>NON-PREFERRED CRITERIA:</u></b></p> <ul style="list-style-type: none"><li>Must have had an inadequate clinical response of at least <u>7 days</u> of at least <u>two</u> preferred drugs with different active ingredients of the same duration of action (SHORT-ACTING or LONG-ACTING)</li><li>Must also meet applicable SHORT-ACTING or LONG-ACTING OPIOID CRITERIA</li></ul> <p><b><u>ADDITIONAL SHORT-ACTING OPIOIDS CRITERIA:</u></b></p> <ul style="list-style-type: none"><li>The system defines an “initial request” as having no opioid</li></ul>
APAP/codeine <sup>AR</sup> but/APAP/caff/cod <sup>AR</sup> 50/325/40/30mg but/ASA/caff/cod <sup>AR</sup> butorphanol codeine <sup>AR</sup> hydrocodone/APAP hydromorphone IR morphine IR oxycodone IR cap, soln, tab oxycodone/APAP tramadol IR <sup>AR</sup> 50mg tramadol/APAP <sup>AR</sup>	APAP/cafeine/ dihydrocodeine but/APAP/caff/cod <sup>AR</sup> 50/300/40/30mg hydrocodone/APAP 5, 7.5, 10-300mg hydrocodone/ibuprofen levorphanol meperidine oxymorphone IR pentazocine/naloxone PROLATE ROXYBOND SEGLENTIS <sup>AR</sup> tramadol IR <sup>AR</sup> soln, 25, 75, 100mg tab	
LONG-ACTING		
BUTRANS <sup>BvG PA</sup> morphine ER tab <sup>PA</sup>	BELBUCA buprenorphine TD patch weekly fentanyl hydrocodone bitartrate ER 12HR cap hydrocodone bitartrate ER 24HR tab hydromorphone ER methadone morphine ER 24HR cap oxycodone ER oxymorphone ER tramadol ER <sup>AR</sup>	

claims in the previous 90 days

- **Initial short-acting requests** can be authorized up to 90 days
  - Length of authorization is dependent on indication, previous patient utilization, and requested length of therapy (could be more restrictive)
  - To exceed acute opioid limits, documentation of the following must be provided:
    - Diagnosis code which must be for somatic type pain
    - Prescriber attestation that the benefits and risks of opioid therapy have been discussed with patient
  - Exemptions to the additional criteria:
    - Patients receiving short-acting opioids for active cancer treatment, palliative care, and end-of-life/hospice care, sickle cell, severe burn, traumatic crushing of tissue, amputation, major orthopedic surgery
    - Prescriber attestation that patient is opioid tolerant (i.e., new to Medicaid or was on higher dose in hospital)
- **Subsequent short-acting requests** can be authorized up to 180 days
  - Documentation of the following must be provided:
    - Current treatment plan
    - Demonstrated adherence to treatment plan through progress notes, including pain and function scores, random urine screening results reviewed, concerns addressed, and no serious adverse outcomes observed
  - Exemptions to the additional criteria:
    - Patients receiving short-acting opioids for cancer pain, palliative care, or end-of-life/hospice care
    - Patients residing in LTC facilities are

exempted from urine drug screening requirements

- **Dose escalation requests** can be authorized up to 180 days
  - Documentation of the following must be provided:
    - Prescriber attestation that dose escalation is likely to result in improved function or pain control
    - Requests for a cumulative daily dose > 80 MED must be prescribed by or in consultation with a pain specialist, specialist in the area of the body affected by pain, or anesthesiologist

*Patients with initial prescriptions for opioid therapy, defined as no rx claims for opioids in the last 90 days, will be limited to 30 MED per day and a maximum of 7 days per prescription. Prior authorization will be required to exceed these limits.*

**ADDITIONAL LONG-ACTING OPIOIDS CRITERIA:**

- The system defines an “initial long-acting request” as having no opioid claims in the previous 90 days
- **Initial long-acting requests** can be authorized up to 90 days
  - Documentation of the following must be provided:
    - Request is a daily dose equivalent of  $\leq 80$  MED
    - Inadequate clinical response to both non-opioid pharmacologic and non-pharmacologic treatments
    - Current use of opioids for  $\geq 30$  of the last 60 days
    - Treatment plan including risk assessment, substance abuse history, concurrent therapies, and requirements for random urine screenings (baseline urine drug tests must be submitted)

- Pain and function scores at each visit
  - Opioid contract required to be in place and submitted with PA form
- Exemptions to the additional criteria:
  - Patients receiving long-acting opioids for cancer pain, palliative care, or end-of-life/hospice care
  - Patients residing in LTC facilities are exempted from urine drug screening and opioid contract requirements
- **Subsequent long-acting requests** can be authorized up to 180 days
  - Documentation of the following must be provided:
    - Current treatment plan
    - Demonstrated adherence to treatment plan through progress notes, including pain and function scores, random urine screening results reviewed, concerns addressed, and no serious adverse outcomes observed
  - Exemptions to the additional criteria:
    - Patients receiving long-acting opioids for cancer pain, palliative care, or end-of-life/hospice care
    - Patients residing in LTC facilities are exempted from urine drug screening and opioid contract requirements
- **Dose escalation requests** can be authorized up to 180 days
  - Documentation of the following must be provided:
    - Prescriber attestation that dose escalation is likely to result in improved function or pain control
    - Requests for a cumulative daily dose > 80

MED must be prescribed by or in consultation with a pain specialist, specialist in the area of the body affected by pain, or anesthesiologist

**ADDITIONAL TRANSMUCOSAL FENTANYL CRITERIA:**

- Must be prescribed by an oncologist, pain specialist, or hospice/palliative prescriber
- Must be concurrently taking a long-acting opioid at a therapeutic dose of any of the following for at least 7 days without adequate pain relief:

≥ 60 mg oral morphine/day	≥ 8 mg oral hydromorphone/day
≥ 25 mcg/hr transdermal fentanyl	≥ 25 mg oral oxymorphone/day
≥ 30 mg oral oxycodone/day	Equianalgesic dose of another opioid

**BUPRENORPHINE BUCCAL FILM (BELBUCA) CRITERIA:**

- Must meet ADDITIONAL LONG-ACTING OPIOID Criteria

**AR** – All codeine and tramadol containing products: a PA is required for patients younger than 12 years old



Blood Formation, Coagulation, and Thrombosis Agents: Colony Stimulating Factors		
PREFERRED AGENTS	NON-PREFERRED AGENTS	PA CRITERIA
FULPHILA <sup>PA</sup> NEUPOGEN <sup>PA</sup> NIVESTYM <sup>PA</sup> NYVEPRIA <sup>PA</sup>	FYLNETRA GRANIX LEUKINE NEULASTA RELEUKO ROLVEDON STIMUFEND UDENYCA ZARXIO ZIEXTENZO	<p><b><u>LENGTH OF AUTHORIZATIONS:</u></b> 30 days or duration of chemotherapy regimen</p> <p><b><u>CLINICAL PA CRITERIA:</u></b></p> <ul style="list-style-type: none"> <li>Must provide documentation of diagnosis, patient's weight (for weight-based dosed medications only), and duration of treatment</li> </ul> <p><b><u>NON-PREFERRED CRITERIA:</u></b></p> <ul style="list-style-type: none"> <li>Must have had an inadequate clinical response of at least <u>14 days</u> with at least <u>one preferred</u> drug in this UPDL category and indicated for diagnosis</li> </ul>

Blood Formation, Coagulation, and Thrombosis Agents: Hematopoietic Agents		
PREFERRED AGENTS	NON-PREFERRED AGENTS	PA CRITERIA
EPOGEN <sup>PA</sup> RETACRIT <sup>PA</sup>	ARANESP MIRCERA PROCRIT	<p><b><u>LENGTH OF AUTHORIZATIONS:</u></b> 180 days; except 365 days for patients with chronic renal failure</p> <p><b><u>CLINICAL PA CRITERIA:</u></b></p> <ul style="list-style-type: none"> <li>• Must provide documentation of baseline hemoglobin level</li> </ul> <p><b><u>NON-PREFERRED CRITERIA:</u></b></p> <ul style="list-style-type: none"> <li>• Must have had an inadequate clinical response of at least <u>14 days</u> with at least <u>one preferred</u> drug in this UPDL category and indicated for diagnosis</li> </ul> <p><b><u>SUBSEQUENT AUTHORIZATION CRITERIA:</u></b></p> <ul style="list-style-type: none"> <li>• Provide current hemoglobin lab result</li> </ul>

**Blood Formation, Coagulation, and Thrombosis Agents: Hemophilia A, von Willebrand Disease, and Factor XIII Deficiency\* LEGACY CATEGORY**

PREFERRED AGENTS	NON-PREFERRED AGENTS	PA CRITERIA
ADVATE <sup>PA</sup> ADYNOVATE <sup>PA</sup> AFSTYLA <sup>PA</sup> ALPHANATE <sup>PA</sup> ALTUVIIIIO <sup>PA</sup> CORIFACT <sup>PA</sup> ELOCTATE <sup>PA</sup> ESPEROCT <sup>PA</sup> FEIBA <sup>PA</sup> HEMLIBRA <sup>PA</sup> HEMOFIL M <sup>PA</sup> HUMATE-P <sup>PA</sup> JIVI <sup>PA</sup> KOATE <sup>PA</sup> KOGENATE FS <sup>PA</sup> KOVALTRY <sup>PA</sup> NOVOEIGHT <sup>PA</sup> NOVOSEVEN RT <sup>PA</sup> NUWIQ KIT <sup>PA</sup> RECOMBINATE <sup>PA</sup> WILATE <sup>PA</sup> XYNTHA <sup>PA</sup>	HYMPAVZI NUWIQ INJ OBIZUR SEVENFACT VONVENDI	<p><b><u>LENGTH OF AUTHORIZATIONS:</u></b> 365 Days</p> <p><b><u>CLINICAL PA CRITERIA:</u></b></p> <ul style="list-style-type: none"> <li>Must provide documentation of patient's body weight (for weight-based dosed medications only)</li> </ul> <p><b><u>NON-PREFERRED CRITERIA:</u></b></p> <ul style="list-style-type: none"> <li>Must have had an inadequate clinical response of at least <u>14 days</u> with at least <u>one preferred</u> drug in this UPDL category and indicated for diagnosis</li> </ul> <p><b><u>ADDITIONAL EXTENDED HALF-LIFE FACTOR CRITERIA</u></b></p> <ul style="list-style-type: none"> <li>Must provide attestation that the patient is not a suitable candidate for treatment with a shorter-acting half-life drug</li> </ul> <p><b><u>ADDITIONAL HYMPAVZI (MARSTACIMAB-HNCQ) CRITERIA</u></b></p> <ul style="list-style-type: none"> <li>Must have had an inadequate clinical response of at least <u>30 days</u> with HEMLIBRA</li> </ul>

## Blood Formation, Coagulation, and Thrombosis Agents: Hemophilia B\* LEGACY CATEGORY

PREFERRED AGENTS	NON-PREFERRED AGENTS	PA CRITERIA
ALPHANINE SD <sup>PA</sup> ALPROLIX <sup>PA</sup> BENEFIX <sup>PA</sup> FEIBA <sup>PA</sup> IDELVION <sup>PA</sup> IXINITY <sup>PA</sup> NOVOSEVEN RT <sup>PA</sup> PROFILNINE <sup>PA</sup> REBINYN <sup>PA</sup> RIXUBIS <sup>PA</sup>	HYMPAVZI SEVENFACT	<p><b><u>LENGTH OF AUTHORIZATIONS:</u></b> 365 Days</p> <p><b><u>CLINICAL PA CRITERIA:</u></b></p> <ul style="list-style-type: none"> <li>Must provide documentation of patient's body weight (for weight-based dosed medications only)</li> </ul> <p><b><u>NON-PREFERRED CRITERIA:</u></b></p> <ul style="list-style-type: none"> <li>Must have had an inadequate clinical response of at least <u>14 days</u> with at least <u>one preferred</u> drug in this UPDL category and indicated for diagnosis</li> </ul> <p><b><u>ADDITIONAL EXTENDED HALF-LIFE FACTOR CRITERIA</u></b></p> <ul style="list-style-type: none"> <li>Must provide attestation that the patient is not a suitable candidate for treatment with a shorter-acting half-life drug</li> </ul>

Blood Formation, Coagulation, and Thrombosis Agents: Heparin-Related Preparations		
PREFERRED AGENTS	NON-PREFERRED AGENTS	PA CRITERIA
enoxaparin	fondaparinux FRAGMIN	<p><b><u>LENGTH OF AUTHORIZATIONS:</u></b> 35 days; except 365 days for patients with cancer, pregnancy, or unable to be converted to an oral anticoagulant</p> <p><b><u>NON-PREFERRED CRITERIA:</u></b></p> <ul style="list-style-type: none"> <li>• Must have had an inadequate clinical response of at least <u>14 days</u> with at least <u>one preferred</u> drug in this UPDL category and indicated for diagnosis</li> </ul>

Blood Formation, Coagulation, and Thrombosis Agents: Oral Anticoagulants		
PREFERRED AGENTS	NON-PREFERRED AGENTS	PA CRITERIA
ELIQUIS PRADAXA CAP <sup>BvG</sup> , PELLET PAK <sup>AR</sup> warfarin XARELTO <sup>BvG</sup> SUSP <sup>AR</sup> , TAB	dabigatran cap rivaroxaban tab SAVAYSA	<p><b><u>LENGTH OF AUTHORIZATION:</u></b> 365 days</p> <p><b><u>NON-PREFERRED CRITERIA:</u></b></p> <ul style="list-style-type: none"> <li>Must have had an inadequate clinical response of at least <u>14 days</u> with at least <u>two preferred</u> drugs in this UPDL category and indicated for diagnosis</li> </ul> <p><b>AR</b> – PRADAXA PELLET PAK, XARELTO SUSP: a PA is required for patients 12 years and older</p>

Blood Formation, Coagulation, and Thrombosis Agents: Oral Antiplatelet		
PREFERRED AGENTS	NON-PREFERRED AGENTS	PA CRITERIA
aspirin IR, ER aspirin/dipyridamole ER BRILINTA <sup>BvG</sup> clopidogrel prasugrel	ticagrelor	<p><b><u>LENGTH OF AUTHORIZATION:</u></b> 365 days</p> <p><b><u>NON-PREFERRED CRITERIA:</u></b></p> <ul style="list-style-type: none"> <li>Must have had an inadequate clinical response of at least <u>30 days</u> with at least <u>one preferred</u> drug in this UPDL category and indicated for diagnosis</li> </ul>

Cardiovascular Agents: Angina, Hypertension and Heart Failure		
PREFERRED AGENTS	NON-PREFERRED AGENTS	PA CRITERIA
ACE INHIBITORS/DIURETICS/COMBINATIONS		LENGTH OF AUTHORIZATIONS: 365 days except nimodipine: 21 days  <u>PROPRANOLOL ORAL SOLN (HEMANGEOL) CRITERIA:</u> <ul style="list-style-type: none"><li>Must provide documentation of the patient’s weight</li></ul> <u>NON-PREFERRED CRITERIA:</u> <ul style="list-style-type: none"><li>Must have had an inadequate clinical response of at least <u>30 days</u> of at least <u>two preferred</u> drugs within the same sub-section classification in this UPDL category and indicated for diagnosis</li></ul> <u>ADDITIONAL APROCITENTAN (TRYVIO) CRITERIA:</u> <ul style="list-style-type: none"><li>Must have had an inadequate clinical response of at least <u>30 days</u> of at least <u>four</u> different classes of antihypertensive medications <u>concurrently</u> without adequate blood pressure control</li></ul> <u>ADDITIONAL FINERENONE (KERENDIA) CRITERIA:</u> <ul style="list-style-type: none"><li>Must be on a maximally tolerated dose of an angiotensin-converting enzyme inhibitor or angiotensin receptor blocker <b>AND</b></li><li>Must provide documentation of an inadequate clinical response to a SGLT2 Inhibitor <b>OR</b> provide documentation of medical necessity beyond convenience for why the patient cannot try a SGLT2 inhibitor (i.e., chronic kidney disease diagnosis)</li></ul> <u>ADDITIONAL MAVACAMTEN (CAMZYOS) CRITERIA:</u> <ul style="list-style-type: none"><li>Must be prescribed by or in consultation with a cardiologist</li><li>Must provide documentation of NYHA Class II-III symptoms and left ventricular ejection fraction ≥55%</li></ul> <u>ADDITIONAL SOTAGLIFLOZIN (INPEFA) CRITERIA:</u> <ul style="list-style-type: none"><li>Must provide documentation of an inadequate clinical response to at least <u>two</u> SGLT2 Inhibitors (refer to Endocrine Agents: Diabetes – Non-Insulin class for complete list)</li></ul>
amlodipine/benazepril benazepril benazepril/HCTZ captopril captopril/HCTZ enalapril soln, tab enalapril/HCTZ fosinopril fosinopril/HCTZ lisinopril lisinopril/HCTZ moexipril quinapril quinapril/HCTZ ramipril trandolapril trandolapril/verapamil	QBRELIS	
ARBs/DIURETICS/COMBINATIONS		
amlodipine/olmesartan amlodipine/valsartan amlodipine/valsartan/HCTZ candesartan candesartan/HCTZ irbesartan irbesartan/HCTZ losartan losartan/HCTZ olmesartan olmesartan/amlodipine/HCTZ olmesartan/HCTZ telmisartan telmisartan/amlodipine	EDARBI EDARBYCLOR valsartan soln	



telmisartan/HCTZ		<b>ADDITIONAL VERICIGUAT (VERQUVO) CRITERIA:</b> <ul style="list-style-type: none"><li>• Must provide documentation of ejection fraction</li><li>• Must have been hospitalized for the treatment of heart failure in the previous 180 days or needs treatment with an outpatient intravenous diuretic in the previous 90 days</li><li>• Must be treated with an agent from <b>ALL</b> the following unless contraindicated:<ul style="list-style-type: none"><li>○ Angiotensin-converting enzyme inhibitor, angiotensin II receptor blocker, <b>OR</b> an angiotensin receptor neprilysin inhibitor</li><li>○ Beta-blocker</li><li>○ Aldosterone antagonist and/or SGLT2 inhibitor as appropriate for renal function</li></ul></li></ul> <b>AR</b> – SOTYLIZE SOLN: a PA is required for patients 6 years and older
valsartan tab		
valsartan/HCTZ		
<b>BETA BLOCKERS/COMBINATIONS</b>		
acebutolol	bisoprolol 2.5mg	
atenolol	carvedilol ER	
atenolol/chlorthalidone	INNOPRAN XL	
betaxolol	KAPSPARGO	
bisoprolol 5, 10mg	labetalol 400mg	
bisoprolol/HCTZ	SOTYLIZE <sup>AR</sup>	
carvedilol IR		
HEMANGEOL <sup>PA</sup>		
labetalol 100, 200, 300mg		
metoprolol succ		
metoprolol tart		
metoprolol/HCTZ		
nadolol		
nebivolol		
propranolol IR, ER		
sotalol		
timolol		
<b>CALCIUM CHANNEL BLOCKERS</b>		
amlodipine	diltiazem 24HR ER tabs	
cartia XT	isradipine	
diltiazem IR	KATERZIA	
diltiazem 12HR ER cap	nimodipine	
diltiazem 24HR ER cap	nisoldipine	
felodipine ER	NORLIQVA	
levamlodipine	NYMALIZE	
nicardipine	verapamil ER (gen of VERELAN PM)	
nifedipine IR, ER		
verapamil IR, ER, SR		
<b>OTHER</b>		
amiloride	aliskiren	
amiloride/HCTZ	ASPRUZYO SPRINKLE	
clonidine IR, patch	CAMZYOS	

doxazosin	clonidine ER (gen of NEXICLON XR)	
ENTRESTO TAB <sup>BvG</sup>	CORLANOR SOLN	
eplerenone	ENTRESTO SPRINKLE CAP	
guanfacine IR, ER	INPEFA	
hydralazine	ivabradine tab (gen of CORLANOR)	
methyldopa	KERENDIA	
minoxidil	spironolactone susp	
ranolazine	TRYVIO	
spironolactone tab	VERQUVO	
spironolactone/HCTZ		
terazosin		

Cardiovascular Agents: Antiarrhythmics		
PREFERRED AGENTS	NON-PREFERRED AGENTS	PA CRITERIA
amiodarone disopyramide dofetilide flecainide mexiletine MULTAQ NORPACE CR propafenone IR, ER	quinidine IR, ER	<p><b><u>LENGTH OF AUTHORIZATIONS:</u></b> 365 Days</p> <p><b><u>NON-PREFERRED CRITERIA:</u></b></p> <ul style="list-style-type: none"> <li>Must have had an inadequate clinical response of at least <u>30 days</u> with at least <u>one preferred</u> drug in this UPDL category and indicated for diagnosis</li> </ul>

Cardiovascular Agents: Lipotropics								
PREFERRED AGENTS	NON-PREFERRED AGENTS	PA CRITERIA						
BILE ACID SEQUESTRANTS		<b>LENGTH OF AUTHORIZATIONS:</b> See below <table><tr><td>JUXTAPID (Initial)</td><td>180 days</td></tr><tr><td>icosapent ethyl cap, LOVAZA, ACL inhibitors (Initial)</td><td>84 days</td></tr><tr><td>All others (Initial and Subsequent)</td><td>365 days</td></tr></table>	JUXTAPID (Initial)	180 days	icosapent ethyl cap, LOVAZA, ACL inhibitors (Initial)	84 days	All others (Initial and Subsequent)	365 days
JUXTAPID (Initial)	180 days							
icosapent ethyl cap, LOVAZA, ACL inhibitors (Initial)	84 days							
All others (Initial and Subsequent)	365 days							
cholestyramine light, regular colesevelam tab colestipol tab prevalite	colesevelam packet colestipol granules							
FIBRIC ACID DERIVATIVES		<b>CLINICAL PA CRITERIA:</b> <ul style="list-style-type: none"><li>Must provide baseline labs <b>AND</b> have adherence to <u>90 days</u> of preferred lipid lowering medications</li><li>Must have had an inadequate clinical response of at least <u>90 days</u> <b>AND</b> unable to reach goal LDL-C (see below) despite treatment with maximally tolerated or high-potency statin (or a clinical reason that these drugs cannot be utilized)</li><li>Must have had an inadequate clinical response of at least 90 days AND unable to reach goal LDL-C (see below) despite treatment with ezetimibe OR documentation that LDL is &gt;25% above goal despite current statin therapy</li></ul> <b>NON-PREFERRED CRITERIA:</b> <ul style="list-style-type: none"><li>Must have had an inadequate clinical response of at least <u>30 days</u> (or <u>90 days</u> for fibrates) with at least <u>one preferred</u> drug within the same sub-section classification in this UPDL category and indicated for diagnosis</li></ul> <b>ADDITIONAL LOVASTATIN ER (ALTOPREV), PITAVASTATIN (LIVALO), FLUVASTATIN (LESCOL) CRITERIA:</b> <ul style="list-style-type: none"><li>Must have had an inadequate clinical response of at least <u>30 days</u> with <u>two preferred</u> drugs in the same drug class</li></ul> <b>ADDITIONAL COLESEVELAM (WELCHOL) CRITERIA:</b> <ul style="list-style-type: none"><li>Must provide documentation of a Type 2 Diabetes diagnosis</li></ul> <b>ADDITIONAL ICOSAPENT ETHYL CRITERIA:</b>						
fenofibrate 48, 54, 145, 160mg tab gemfibrozil	fenofibrate IR, DR cap fenofibrate 40, 120mg tab fenofibric acid							
PCSK9 INHIBITORS								
PRALUENT <sup>PA</sup> REPATHA <sup>PA</sup>								
STATINS/COMBINATIONS								
atorvastatin ezetimibe/simvastatin lovastatin pravastatin rosuvastatin simvastatin	ALTOPREV amlodipine/atorvastatin ATORVALIQ EZALLOR fluvastatin IR, ER pitavastatin ZYPITAMAG							
OTHER								
ezetimibe niacin IR, ER OTC omega-3-acid ethyl esters	icosapent ethyl cap JUXTAPID NEXLETOL NEXLIZET niacin ER tab							

- Must provide documentation of baseline labs indicating triglyceride levels  $\geq 500\text{mg/dL}$  after an inadequate clinical response to fibrates, niacin, and diet/exercise

#### **ADDITIONAL LOMITAPIDE (JUXTAPID) & ATP CITRATE LYASE (ACL) INHIBITOR**

##### **CRITERIA:**

- Must provide documentation of baseline labs **AND** have documented adherence to 90 days of prescribed lipid lowering medications
- Must have had inadequate clinical response of at least 90 days **AND** unable to reach goal LDL-C with high-potency statin, ezetimibe and PCSK9 inhibitor (or a clinical reason that these drugs cannot be utilized)

##### **ADDITIONAL INFORMATION:**

- High potency statins: atorvastatin (LIPITOR) 40-80mg & rosuvastatin (CRESTOR) 20-40mg
- LDL goals for Familial Hypercholesterolemia (includes Heterozygous & Homozygous FH): LDL  $\leq 100\text{mg/dL}$  for adults or LDL  $\leq 110\text{mg/dL}$  for those < 18 years of age
- LDL goals for Clinical Atherosclerotic Cardiovascular Disease (ASCVD) not at very high risk: LDL  $\leq 70\text{mg/dL}$
- LDL goals for Clinical Atherosclerotic Cardiovascular Disease (ASCVD) at very high risk: LDL  $\leq 55\text{mg/dL}$
- Must provide documentation of multiple major ASCVD events or 1 major ASCVD event and multiple high-risk conditions if citing goal LDL  $\leq 55\text{mg/dL}$

Cardiovascular Agents: Pulmonary Arterial Hypertension* LEGACY CATEGORY		
PREFERRED AGENTS	NON-PREFERRED AGENTS	PA CRITERIA
ENDOTHELIN RECEPTOR ANTAGONISTS		<b>LENGTH OF AUTHORIZATIONS:</b> 365 Days  <b>CLINICAL PA CRITERIA:</b> <ul style="list-style-type: none"><li>Must provide documentation of NYHA Functional Class symptoms for Pulmonary Hypertension experienced by patient</li></ul> <b>NON-PREFERRED CRITERIA:</b> <ul style="list-style-type: none"><li>Must have had an inadequate clinical response of at least <u>30 days</u> with at least <u>two preferred</u> drugs in this UPDL category and indicated for diagnosis, if available, <u>one</u> of which must be a phosphodiesterase-5 inhibitor</li></ul> <b>ADDITIONAL INFORMATION:</b> <ul style="list-style-type: none"><li>Patients who have class III or IV symptoms defined by the NYHA Functional Class for Pulmonary Hypertension may be authorized for inhalation or intravenous agents</li></ul> <b>AR</b> – sildenafil susp: a PA is required for patients 18 years and older <b>AR</b> – TADLIQ: a PA is required for patients younger than 18 years
ambrisentan <sup>PA</sup> bosentan <sup>PA</sup>	OPSUMIT TRACLEER SUSP	
PDE5 INHIBITORS		
sildenafil <sup>PA</sup> sildenafil susp <sup>AR PA</sup> tadalafil <sup>PA</sup> TADLIQ <sup>AR PA</sup>		
PROSTAGLANDINS		
epoprostenol	ORENITRAM treprostinil TYVASO VENTAVIS	
OTHER		
	ADEMPAS OPSYNVI UPTRAVI WINREVAIR	

Central Nervous System (CNS) Agents: Alzheimer's Agents* LEGACY CATEGORY		
PREFERRED AGENTS	NON-PREFERRED AGENTS	PA CRITERIA
donepezil <sup>AR</sup> galantamine IR tab, ER cap <sup>AR</sup> memantine IR, ER tab <sup>AR</sup> rivastigmine cap <sup>AR</sup> rivastigmine patch <sup>AR</sup>	ADLARITY <sup>AR</sup> galantamine soln <sup>AR</sup> memantine/donepezil cap <sup>AR</sup> 14-10, 21-10, 28-10mg memantine soln <sup>AR</sup> NAMZARIC <sup>AR</sup>	<b><u>LENGTH OF AUTHORIZATIONS:</u></b> 365 Days  <b><u>NON-PREFERRED CRITERIA:</u></b> <ul style="list-style-type: none"> <li>Must have had an inadequate clinical response of at least <u>30 days</u> with at least <u>two preferred</u> drugs in this UPDL category and indicated for diagnosis</li> </ul> <b>AR</b> – All drugs: a PA is required for patients younger than 40 years

Central Nervous System (CNS) Agents: Anti-Migraine Agents, Acute		
PREFERRED AGENTS	NON-PREFERRED AGENTS	PA CRITERIA
CGRP INHIBITORS		<b>LENGTH OF AUTHORIZATIONS:</b> 180 Days  <b>STEP THERAPY CRITERIA:</b> <ul style="list-style-type: none"><li>Must have had an inadequate clinical response of at least <u>14 days</u> with at least <u>two preferred</u> drugs in this UPDL category <b>OR</b> documentation why patient is unable to take product not requiring step therapy</li></ul> <b>NON-PREFERRED CRITERIA:</b> <ul style="list-style-type: none"><li>Must have had an inadequate clinical response of at least <u>14 days</u> with at least <u>one</u> preferred drug and <u>one</u> step therapy drug in this UPDL category and indicated for diagnosis, if available</li></ul> <b>ADDITIONAL INFORMATION:</b> <ul style="list-style-type: none"><li>NURTEC has a maximum quantity of <b>8</b> tablets per month for acute migraines</li></ul>
NURTEC ODT <sup>ST</sup> UBRELVY <sup>ST</sup>	ZAVZPRET	
TRIPTANS/COMBINATIONS		
IMITREX NASAL SPRAY naratriptan rizatriptan sumatriptan inj, nasal spray, tab	almotriptan eletriptan frovatriptan sumatriptan/naproxen TOSYMRA zolmitriptan	
OTHER		
	dihydroergotamine MIGERGOT REYVOW	



Central Nervous System (CNS) Agents: Anti-Migraine Agents, Cluster Headache		
PREFERRED AGENTS	NON-PREFERRED AGENTS	PA CRITERIA
verapamil IR, ER	EMGALITY 100mg/ml	<p><b><u>LENGTH OF AUTHORIZATIONS:</u></b> 180 days</p> <p><b><u>NON-PREFERRED CRITERIA:</u></b></p> <ul style="list-style-type: none"> <li>• Must have had an inadequate clinical response of at least <u>60 days</u> to at least <u>one preferred</u> drug in this UPDL category and indicated for diagnosis</li> </ul> <p><b><u>ADDITIONAL INFORMATION:</u></b></p> <ul style="list-style-type: none"> <li>• An inadequate clinical response to verapamil is defined as a titration to at least 480mg daily or maximally tolerated dose based on blood pressure or heart rate and maintained for at least 60 days</li> </ul>

Central Nervous System (CNS) Agents: Anti-Migraine Agents, Prophylaxis		
PREFERRED AGENTS	NON-PREFERRED AGENTS	PA CRITERIA
AIMOVIG <sup>ST</sup> AJOVY <sup>ST</sup> EMGALITY 120mg/ml <sup>ST</sup> Cardiovascular Agents: Beta-Blockers CNS Agents: Anticonvulsants CNS Agents: Serotonin-Norepinephrine Reuptake Inhibitors CNS Agents: Tricyclic Antidepressants	NURTEC ODT QULIPTA	<p><b><u>LENGTH OF AUTHORIZATIONS:</u></b> Initial: 180 days; Subsequent: 365 days</p> <p><b><u>STEP THERAPY CRITERIA:</u></b></p> <ul style="list-style-type: none"> <li>Must have had an inadequate clinical response of at least <u>30 days</u> with at least <u>two preferred</u> controller migraine drugs. <ul style="list-style-type: none"> <li>For patients already established on a serotonergic medication, only <u>one</u> preferred controller migraine drugs will be required</li> </ul> </li> <li>Must include objective documentation of severity, frequency, type of migraine, and number of headache days per month</li> <li>Controller migraine drug classes include beta-blockers, anticonvulsants, serotonin-norepinephrine reuptake inhibitors, or tricyclic antidepressants</li> </ul> <p><b><u>ERENUMAB (AIMOVIG) CRITERIA:</u></b></p> <ul style="list-style-type: none"> <li>Must have had an inadequate clinical response of at least <u>60 days</u> with the 70mg dose to request a dose increase</li> </ul> <p><b><u>FREMANEZUMAB (AJOVY) CRITERIA:</u></b></p> <ul style="list-style-type: none"> <li>Must have demonstrated efficacy for at least <u>90 days</u> before quarterly administration will be authorized</li> </ul> <p><b><u>NON-PREFERRED CRITERIA:</u></b></p> <ul style="list-style-type: none"> <li>Must have had an inadequate clinical response of at least <u>30 days</u> with at least <u>three preferred</u> controller migraine drugs <b>AND</b> <u>one step therapy</u> drug in this UPDL category</li> </ul> <p><b><u>ADDITIONAL INFORMATION:</u></b></p> <ul style="list-style-type: none"> <li>NURTEC has a maximum quantity of <b>16</b> tablets per month for migraine prophylaxis</li> </ul> <p><b><u>SUBSEQUENT AUTHORIZATION CRITERIA:</u></b></p> <ul style="list-style-type: none"> <li>Must provide documentation of patient's clinical response to treatment (Objective documentation of severity, frequency, and number of headache days per month).</li> </ul>

**Central Nervous System (CNS) Agents: Anticonvulsants\* LEGACY CATEGORY**

PREFERRED AGENTS	NON-PREFERRED AGENTS	PA CRITERIA
BANZEL TAB <sup>BvG</sup> BRIVIACT SOLN <sup>AR</sup> , TAB carbamazepine IR, ER clobazam clonazepam DIACOMIT <sup>PA</sup> divalproex DR, ER EPIDIOLEX <sup>PA</sup> EPRONTIA <sup>AR</sup> ethosuximide FYCOMPA <sup>ST</sup> gabapentin lacosamide lamotrigine chew, IR, ODT levetiracetam IR tab, soln oxcarbazepine IR tab phenobarbital phenytoin IR, ER pregabalin IR primidone topiramate IR TRILEPTAL SUSP <sup>BvG</sup> valproic acid zonisamide cap	APTIOM CELONTIN <sup>BvG</sup> clonazepam ODT ELEPSIA XR felbamate FINTEPLA lamotrigine ER levetiracetam ER tab MOTPOLY XR oxcarbazepine susp OXTELLAR XR <sup>BvG</sup> QUDEXY XR <sup>BvG</sup> rufinamide tab, soln SPRITAM <sup>BvG</sup> SYMPAZAN tiagabine topiramate sprinkle cap TROKENDI XR <sup>BvG</sup> vigabatrin vigabatrin powder <sup>AR</sup> VIGAFYDE <sup>AR</sup> XCOPRI ZONISADE SUSP ZTALMY	<p><b><u>LENGTH OF AUTHORIZATIONS:</u></b> 365 days except EPIDIOLEX and DIACOMIT – Initial: 180 days</p> <p><b><u>STEP THERAPY CRITERIA:</u></b></p> <ul style="list-style-type: none"> <li>Must have had an inadequate clinical response of at least <u>30 days</u> with at least <u>one preferred</u> drug in this UPDL category</li> </ul> <p><b><u>CANNABIDIOL (EPIDIOLEX) CRITERIA</u></b></p> <ul style="list-style-type: none"> <li>Must have had an inadequate clinical response of at least <u>30 days</u> with any <u>two</u> of the following anticonvulsants: clobazam, levetiracetam, valproic acid, lamotrigine, topiramate, rufinamide, or felbamate within the past <u>365 days</u> (members who meet this criterion will not require a PA)</li> </ul> <p><b><u>STIRIPENTOL (DIACOMIT) CRITERIA</u></b></p> <ul style="list-style-type: none"> <li>Exempt from Legacy rules</li> <li>Must be prescribed by or in consultation with a neurologist</li> <li>Must be concomitantly taking clobazam (ONFI)</li> <li>Must provide documentation of addressed comorbidities and baseline hematologic testing (CBC)                             <ul style="list-style-type: none"> <li>Patients with phenylketonuria (PKU) must provide evidence of total daily amount of phenylalanine</li> <li>Prescribers must include management plans for patients with neutrophil counts &lt;1,500 cells/mm<sup>3</sup> or platelet count &lt;150,000/μL</li> </ul> </li> <li>Must provide documentation of patient's weight                             <ul style="list-style-type: none"> <li>Maximum daily dose does not exceed: 50 mg/kg/day or 3,000mg/day</li> </ul> </li> </ul> <p><b><u>NON-PREFERRED CRITERIA:</u></b></p> <ul style="list-style-type: none"> <li>Must have had an inadequate clinical response of at least <u>30 days</u> with at least <u>two preferred</u> drugs in this UPDL category and indicated for diagnosis</li> </ul>

- Prescriptions submitted from a prescriber who is credentialed as a neurology specialty with Ohio Medicaid AND for drugs that are used only for seizures, there must have been an inadequate clinical response of at least 30 days with one preferred drug. This provision applies only to the standard tablet/capsule dosage form.

**AR** – BRIVIACT SOLN: a PA is required for patients 12 years and older

**AR** – EPRONTIA SOLN: a PA is required for patients 12 years and older

**AR** – vigabatrin powder: a PA is required for patients 2 years and older

**AR** – VIGAFYDE SOLN: a PA is required for patients 2 years and older

Central Nervous System (CNS) Agents: Anticonvulsants Rescue		
PREFERRED AGENTS	NON-PREFERRED AGENTS	PA CRITERIA
diazepam gel LIBERVANT <sup>AR</sup> NAYZILAM <sup>AR</sup> VALTOCO <sup>AR</sup>		All products are covered without a PA  <b>AR</b> – LIBERVANT: a PA is required for patients 5 years and older <b>AR</b> – NAYZILAM: a PA is required for patients younger than 12 years old <b>AR</b> – VALTOCO: a PA is required for patients younger than 2 years old

Central Nervous System (CNS) Agents: Antidepressants* LEGACY CATEGORY		
PREFERRED AGENTS	NON-PREFERRED AGENTS	PA CRITERIA
NDRIs		<b>LENGTH OF AUTHORIZATIONS:</b> 365 Days except 14 days with no renewal for ZURZUVAE
bupropion bupropion SR (gen of WELLBUTRIN SR) bupropion XL (gen of WELLBUTRIN XL)	APLENZIN bupropion XL (gen of FORFIVO XL)	
SNRIs		<b>PSYCHIATRIST EXEMPTION:</b> <ul style="list-style-type: none"><li>Prescribers (as identified below) are exempt from prior authorization of any non-preferred antidepressant, or step therapy of any preferred drug, in the standard tablet/capsule dosage forms. Other dosage forms may still require prior authorization. The exemption will be processed by the claims system when the pharmacy has submitted the prescriber on the claim using the individual national provider identifier (NPI) for the prescriber. <b>Prescribers are defined as:</b> Physicians with a specialty in psychiatry, nurse practitioners certified in psychiatric mental health, or clinical nurse specialists certified in psychiatric mental health, who are credentialed with the Ohio Department of Medicaid.</li></ul>
desvenlafaxine succ ER (gen of PRISTIQ) duloxetine 20, 30, 60mg venlafaxine IR tab, ER cap	desvenlafaxine ER (gen of KHEDEZLA) DRIZALMA SPRINKLE duloxetine 40mg FETZIMA venlafaxine ER tab	
SSRIs		<b>CLINICAL PA CRITERIA:</b> <ul style="list-style-type: none"><li>Must have a diagnosis of moderate to severe Post-Partum Depression (PPD) no earlier than the 3<sup>rd</sup> trimester OR within 12 months of pregnancy delivery</li></ul>
citalopram tab, soln escitalopram fluoxetine IR 10, 20, 40mg fluoxetine soln fluvoxamine IR paroxetine IR tab, soln sertraline tab	citalopram cap fluoxetine IR 60mg, DR fluvoxamine ER paroxetine ER tab sertraline cap	
OTHER		<b>STEP THERAPY CRITERIA:</b> <ul style="list-style-type: none"><li>Must have had an inadequate clinical response of at least <u>30 days</u> with at least <u>two preferred</u> drugs in this UPDL category</li></ul> <b>NON-PREFERRED CRITERIA:</b> <ul style="list-style-type: none"><li>Must have had an inadequate clinical response of at least <u>30 days</u> with at least <u>two preferred</u> drugs in this UPDL category and indicated for diagnosis</li></ul> <b>ADDITIONAL DEXTROMETHORPHAN/BUPROPION (AUVELITY) CRITERIA:</b> <ul style="list-style-type: none"><li>Must have an inadequate clinical response of at least <u>30 days</u> with <b>ALL</b> of the following:<ul style="list-style-type: none"><li><u>ONE</u> norepinephrine/dopamine reuptake inhibitor (NDRI)</li></ul></li></ul>
mirtazapine nefazodone tranylcypromine trazodone 50, 100, 150mg vilazodone VRAYLAR <sup>ST</sup> ZURZUVAE <sup>PA</sup>	AUVELITY CAPLYTA clomipramine EMSAM MARPLAN phenelzine REXULTI trazodone 300mg TRINTELLIX	

- |  |  |  |
|--|--|--|
|  |  | <ul style="list-style-type: none"><li>○ <u>ONE</u> serotonin and norepinephrine reuptake inhibitor (SNRI)</li><li>○ <u>TWO</u> selective serotonin reuptake inhibitors (SSRIs) (<u>ONE</u> of which must be either vilazodone (VIBRYD) OR vortioxetine (TRINTELLIX))</li></ul> |
|--|--|--|

Central Nervous System (CNS) Agents: Attention Deficit Hyperactivity Disorder Agents		
PREFERRED AGENTS	NON-PREFERRED AGENTS	PA CRITERIA
<b>NON-STIMULANTS</b>		<b>LENGTH OF AUTHORIZATIONS:</b> 365 days
atomoxetine cap <sup>AR</sup> clonidine ER guanfacine ER ONYDA XR SUSP <sup>AR</sup> QELBREE <sup>ST</sup>		<b>STEP THERAPY CRITERIA:</b> <ul style="list-style-type: none"> <li>Must have had an inadequate clinical response of at least <u>30 days</u> with atomoxetine <b>OR</b> at least <u>one preferred</u> ADHD agent.</li> </ul>
<b>STIMULANTS</b>		<b>NON-PREFERRED CRITERIA:</b>
amphetamine/dextroamphetamine IR, ER <sup>AR</sup> CONCERTA dexmethylphenidate tab <sup>AR</sup> dexmethylphenidate ER (gen of FOCALIN XR) <sup>AR</sup> dextroamphetamine IR tab, ER cap <sup>AR</sup> DYANAVEL XR FOCALIN XR <sup>AR</sup> methylphenidate ER cap (gen of METADATE CD, RITALIN LA) methylphenidate ER tab (gen of CONCERTA, METHYLIN ER, RITALIN SR) methylphenidate soln <sup>AR</sup> methylphenidate tab PROCENTRA <sup>BvG AR</sup> QUILLICHEW ER <sup>AR</sup> QUILLIVANT XR <sup>AR</sup> RITALIN LA VYVANSE CAP <sup>BvG</sup>	ADZENYS XR ODT amphetamine tab AZSTARYS <sup>AR</sup> COTEMPLA XR ODT DAYTRANA <sup>BvG</sup> dextroamphetamine soln <sup>AR</sup> EVEKEO ODT JORNAY PM lisdexamfetamine cap methamphetamine methylphenidate chewable tab <sup>AR</sup> methylphenidate ER cap, tab (gen of APTENSIO XR, RELEXXII) MYDAYIS <sup>BvG</sup> VYVANSE CHEWABLE TAB <sup>BvG</sup> XELSTRYM <sup>AR</sup>	<b>NON-PREFERRED CRITERIA:</b> <ul style="list-style-type: none"> <li>Must have had an inadequate clinical response of at least <u>30 days</u> with at least <u>two preferred</u> drugs in this UPDL category and indicated for diagnosis</li> </ul> <b>ADDITIONAL INFORMATION:</b> <ul style="list-style-type: none"> <li>Requests for non-preferred immediate-release formulations must have all required trials with preferred immediate-release drugs, and requests for non-preferred extended-release formulations must have all required trials with preferred extended-release drugs</li> <li>For patients established on drugs that change from preferred to non-preferred on January 1, a prior authorization is <b>NOT</b> required until <b>after</b> June 30<sup>th</sup> of that year.</li> </ul> <p><b>AR</b> –amphetamine/dextroamphetamine, dextroamphetamine IR: a PA is required for patients younger than 3 years</p> <p><b>AR</b> –amphetamine/dextroamphetamine XR, atomoxetine, dextroamphetamine ER, dexmethylphenidate &amp; XELSTRYM: a PA is required for patients younger than 6 years</p> <p><b>AR</b> – dextroamphetamine soln: a PA is required for patients 12 years and older</p> <p><b>AR</b> – methylphenidate soln/susp/chewable tab: a PA is required for patients 12 years and older</p> <p><b>AR</b> – ONYDA XR SUSP: a PA is required for patients 12 years and older</p>



Central Nervous System (CNS) Agents: Atypical Antipsychotics* LEGACY CATEGORY		
PREFERRED AGENTS	NON-PREFERRED AGENTS	PA CRITERIA
ABILIFY ASIMTUFI, MAINTENA aripiprazole ARISTADA ARISTADA INITIO asenapine <sup>ST</sup> clozapine FANAPT <sup>ST</sup> GEODON INVEGA HAFYERA ER <sup>PA</sup> INVEGA SUSTENNA INVEGA TRINZA lurasidone olanzapine paliperidone tab PERSERIS quetiapine IR, ER RISPERDAL CONSTA <sup>BvG</sup> risperidone RYKINDO UZEDY VRAYLAR <sup>ST</sup> ziprasidone	ABILIFY MYCITE aripiprazole ODT, soln CAPLYTA clozapine ODT COBENFY EQUETRO ERZOFRI fluoxetine/olanzapine LYBALVI NUPLAZID OPIPZA REXULTI risperidone microspheres SECUADO VERSACLOZ ZYPREXA RELPREVV	<p><b><u>LENGTH OF AUTHORIZATIONS:</u></b> 365 Days</p> <p><b><u>PSYCHIATRIST EXEMPTION:</u></b></p> <ul style="list-style-type: none"> <li>Prescribers (as identified below) are exempt from prior authorization of any non-preferred second-generation antipsychotic, or step therapy of any preferred drug, in the standard tablet/capsule and long-acting injectable dosage forms. Other dosage forms may still require prior authorization. The exemption will be processed by the claims system when the pharmacy has submitted the prescriber on the claim using the individual national provider identifier (NPI) for the prescriber. <b>Prescribers are defined as:</b> Physicians with a specialty in psychiatry, nurse practitioners certified in psychiatric mental health, or clinical nurse specialists certified in psychiatric mental health, who are credentialed with the Ohio Department of Medicaid.</li> </ul> <p><b><u>PALIPERIDONE PALMITATE (INVEGA HAFYERA) CRITERIA:</u></b></p> <ul style="list-style-type: none"> <li>Must have had 4 months of treatment with INVEGA SUSTENNA or 3 months with INVEGA TRINZA</li> </ul> <p><b><u>STEP THERAPY CRITERIA:</u></b></p> <ul style="list-style-type: none"> <li>Must have had an inadequate clinical response of at least <u>30 days</u> with at least <u>one preferred</u> drug in this UPDL category</li> </ul> <p><b><u>NON-PREFERRED CRITERIA:</u></b></p> <ul style="list-style-type: none"> <li>Must have had an inadequate clinical response of at least <u>30 days</u> with at least <u>two preferred</u> drugs in this UPDL category and indicated for diagnosis</li> </ul> <p><b><u>ADDITIONAL ARIPIPRAZOLE (ABILIFY MYCITE) CRITERIA:</u></b></p> <ul style="list-style-type: none"> <li>Must be prescribed by or in consultation with a psychiatrist following an aripiprazole serum blood level draw indicating need for further investigation of adherence</li> </ul>

**ADDITIONAL OLANZAPINE/SAMIDORPHAN (LYBALVI) CRITERIA:**

- Must provide documentation that patient is not using opioids or undergoing acute opioid withdrawal

**ADDITIONAL PIMAVANSERIN (NUPLAZID) CRITERIA:**

- For Parkinson-related Hallucinations & Delusions **ALL** of the following must be met:
  - Psychotic symptoms are severe and frequent enough to warrant treatment with an antipsychotic **AND** are not related to dementia or delirium
  - The patient's other Parkinson's Disease drugs have been reduced or adjusted and psychotic symptoms persist **OR** patient is unable to tolerate adjustment of these other drugs
  - Must have been inadequate clinical response or contraindication to at least 30 days of either quetiapine or clozapine
- An exemption to the criteria will be authorized for prescribers with a neurology specialty to a patient with a history of the related condition

**ADDITIONAL INFORMATION:**

- Long-acting injectable antipsychotics may be billed by the pharmacy if they are not dispensed directly to the patient. If not administered by the pharmacist, the drug must be released only to the administering provider or administering provider's staff, following all regulations for a Prescription Pick-Up Station as described by the Ohio Board of Pharmacy

Central Nervous System (CNS) Agents: Fibromyalgia Agents		
PREFERRED AGENTS	NON-PREFERRED AGENTS	PA CRITERIA
pregabalin IR SAVELLA		All products are covered without a PA

## Central Nervous System (CNS) Agents: Medication Assisted Treatment of Opioid Addiction

PREFERRED AGENTS	NON-PREFERRED AGENTS	PA CRITERIA
BRIXADI buprenorphine/naloxone clonidine IR, ER SUBLOCADE SUBOXONE VIVITROL ZUBSOLV	buprenorphine LUCEMYRA <sup>BvG</sup>	<p><b><u>LENGTH OF AUTHORIZATIONS:</u></b> 180 days except 14 days for LUCEMYRA</p> <p><b><u>ADDITIONAL LOFEXIDINE (LUCEMYRA) CRITERIA</u></b></p> <ul style="list-style-type: none"> <li>May be authorized if <b>ALL</b> of the following criteria are met:                             <ul style="list-style-type: none"> <li>Must provide medical justification supporting why an opioid taper (such as with buprenorphine or methadone) cannot be used</li> <li>Must have had an inadequate clinical response or contraindication to clonidine</li> </ul> </li> <li>Must provide documentation that the drug was initiated in an inpatient setting to be exempt from the above criteria</li> </ul> <p><b><u>BUPRENORPHINE SAFETY EDITS AND DRUG UTILIZATION REVIEW CRITERIA:</u></b></p> <ul style="list-style-type: none"> <li>Prescribing for buprenorphine products must follow the requirements of Ohio Administrative Code rule 4731-33-03 <i>Office based treatment for opioid addiction</i>.</li> <li>In favor of eliminating prior authorization for all forms of oral short acting buprenorphine- containing products, ODM and the Managed Care Plans will implement safety edits and a retrospective drug utilization review process for all brand and generic forms of oral short acting buprenorphine-containing products. Safety edits are in place for dosages over 24mg of buprenorphine equivalents/day.</li> <li>buprenorphine sublingual tablets (generic SUBUTEX) will be restricted to pregnancy, breastfeeding, or allergy/contraindication to preferred products</li> </ul> <p><b><u>ADDITIONAL INFORMATION</u></b></p> <ul style="list-style-type: none"> <li>VIVITROL, SUBLOCADE, and BRIXADI may be billed by the pharmacy if it is not dispensed directly to the patient. If not administered by the pharmacist, the drug must be released only to the administering provider or administering provider's staff, following all regulations for a Prescription Pick-Up Station as described by the Ohio Board of Pharmacy.</li> </ul>

Central Nervous System (CNS) Agents: Movement Disorders		
PREFERRED AGENTS	NON-PREFERRED AGENTS	PA CRITERIA
AUSTEDO IR, XR <sup>PA ST</sup> INGREZZA <sup>PA ST</sup> tetrabenazine		<p><b><u>LENGTH OF AUTHORIZATIONS:</u></b> 365 Days</p> <p><b><u>CLINICAL PA CRITERIA:</u></b></p> <ul style="list-style-type: none"> <li>• Must be prescribed by or in consultation with a neurologist or psychiatrist</li> </ul> <p><b><u>STEP THERAPY CRITERIA:</u></b></p> <ul style="list-style-type: none"> <li>• Must have an inadequate clinical response of at least <u>90 days</u> to a maximally tolerated dose of tetrabenazine for Huntington's Disease only</li> </ul>

Central Nervous System (CNS) Agents: Multiple Sclerosis* LEGACY CATEGORY		
PREFERRED AGENTS	NON-PREFERRED AGENTS	PA CRITERIA
AVONEX BETASERON COPAXONE <sup>BvG</sup> dalfampridine dimethyl fumarate fingolimod GILENYA KESIMPTA REBIF teriflunomide	BAFIERTAM glatiramer glatopa MAVENCLAD MAYZENT OCREVUS PLEGRIDY PONVORY TASCENSO ODT VUMERITY ZEPOSIA	<p><b><u>LENGTH OF AUTHORIZATIONS:</u></b> 365 Days</p> <p><b><u>NON-PREFERRED CRITERIA:</u></b></p> <ul style="list-style-type: none"> <li>Must have had an inadequate clinical response of at least <u>30 days</u> with at least <u>one preferred</u> drug in this UPDL category and indicated for diagnosis</li> </ul> <p><b><u>ADDITIONAL OCRELIZUMAB (OCREVUS) CRITERIA:</u></b></p> <ul style="list-style-type: none"> <li>Must provide documentation of diagnosis of primary progressive multiple sclerosis <b>OR</b> must have had an inadequate clinical response of at least <u>30 days</u> with at least <u>one preferred</u> drug in this UPDL category</li> </ul> <p><b><u>ADDITIONAL SIPONIMOD (MAYZENT) CRITERIA:</u></b></p> <ul style="list-style-type: none"> <li>Must provide documentation of CYP2C9 genotype</li> </ul>

Central Nervous System (CNS) Agents: Narcolepsy		
PREFERRED AGENTS	NON-PREFERRED AGENTS	PA CRITERIA
amphetamine/ dextroamphetamine IR/ER <sup>AR</sup> armodafinil dextroamphetamine ER <sup>AR</sup> methylphenidate ER methylphenidate tab modafinil	SUNOSI WAKIX XYREM <sup>BvG</sup> XYWAV	<p><b><u>LENGTH OF AUTHORIZATIONS:</u></b> 365 days</p> <p><b><u>NON-PREFERRED CRITERIA:</u></b></p> <ul style="list-style-type: none"> <li>Must have had an inadequate clinical response with at least <u>two preferred</u> drugs - either at least <u>30 days</u> of armodafinil or modafinil; <b>OR</b> at least <u>7 days</u> of a preferred amphetamine or methylphenidate drug in this UPDL category and indicated for diagnosis</li> </ul> <p><b><u>ADDITIONAL OXYBATE SALTS (XYWAV) CRITERIA:</u></b></p> <ul style="list-style-type: none"> <li>Must have documented adherence to sodium restricted diet</li> </ul> <p><b>AR</b> –amphetamine/dextroamphetamine: a PA is required for patients younger than 3 years  <b>AR</b> –amphetamine/dextroamphetamine XR, dextroamphetamine ER: a PA is required for patients younger than 6 years</p>

Central Nervous System (CNS) Agents: Neuropathic Pain		
PREFERRED AGENTS	NON-PREFERRED AGENTS	PA CRITERIA
DIBENZAZEPINES		<b>LENGTH OF AUTHORIZATIONS:</b> 365 Days  <b>STEP THERAPY CRITERIA:</b> <ul style="list-style-type: none"><li>Must have had an inadequate clinical response of at least <u>30 days</u> with generic lidocaine patch</li></ul> <b>NON-PREFERRED CRITERIA:</b> <ul style="list-style-type: none"><li>Must have had an inadequate clinical response of at least <u>30 days</u> with at least <u>two preferred</u> drugs within the same sub-section classification in this UPDL category and indicated for diagnosis</li></ul>
carbamazepine IR, ER oxcarbazepine tab TRILEPTAL SUSP <sup>BvG</sup>	oxcarbazepine susp	
GAPAPENTINOIDS		
gabapentin IR GRALISE <sup>BvG</sup> HORIZANT	gabapentin ER	
TRICYCLIC ANTIDEPRESSANTS		
amitriptyline desipramine doxepin 10, 25, 50, 75, 100, 150mg doxepin soln imipramine nortriptyline		
OTHER		
duloxetine 20, 30, 60mg lidocaine patch pregabalin IR ZTLIDO <sup>ST</sup>	duloxetine 40mg pregabalin ER	



Central Nervous System (CNS) Agents: Parkinson's Agents		
PREFERRED AGENTS	NON-PREFERRED AGENTS	PA CRITERIA
COMT INHIBITORS		<b>LENGTH OF AUTHORIZATIONS:</b> 365 Days  <b>NON-PREFERRED CRITERIA:</b> <ul style="list-style-type: none"><li>Must have had an inadequate clinical response of at least <u>30 days</u> with at least <u>one preferred</u> drug within the same sub-section classification in this UPDL category and indicated for diagnosis</li></ul> <b>ADDITIONAL APOMORPHINE (APOKYN/KYNMOBI), LEVODOPA INHALATION (INBRIJA), &amp; ISTRADEFYLLINE (NOURIANZ) CRITERIA:</b> <ul style="list-style-type: none"><li>Must have had inadequate clinical response to at least <u>30 days</u> with one other drug for the treatment of “off episodes” (COMT inhibitor, dopamine agonist, or MAO-B inhibitor)</li></ul>
entacapone	ONGENTYS tolcapone	
DOPAMINE AGONISTS		
pramipexole IR ropinirole IR, ER	apomorphine KYNMOBI NEUPRO pramipexole ER	
MAO-B INHIBITORS		
selegiline	rasagiline XADAGO ZELAPAR	
OTHER		
amantadine cap, tab carbidopa carbidopa/levodopa IR, ER	amantadine soln carbidopa/levodopa dispersible carbidopa/levodopa/entacapone CREXONT GOCOVRI INBRIJA NOURIANZ OSMOLEX ER RYTARY VYALEV	

Central Nervous System (CNS) Agents: Restless Legs Syndrome		
PREFERRED AGENTS	NON-PREFERRED AGENTS	PA CRITERIA
HORIZANT pramipexole IR ropinirole IR, ER	NEUPRO	<p><b><u>LENGTH OF AUTHORIZATIONS:</u></b> 365 Days</p> <p><b><u>NON-PREFERRED CRITERIA:</u></b></p> <ul style="list-style-type: none"> <li>Must have had an inadequate clinical response of at least <u>30 days</u> with at least <u>one preferred</u> drug in this UPDL category and indicated for diagnosis</li> </ul>

Central Nervous System (CNS) Agents: Sedative-Hypnotics, Non-Barbiturate		
PREFERRED AGENTS	NON-PREFERRED AGENTS	PA CRITERIA
BELSOMRA estazolam eszopiclone ramelteon temazepam triazolam zaleplon zolpidem tab ER, IR	DAYVIGO doxepin 3, 6mg EDLUAR flurazepam quazepam QUVIVIQ zolpidem cap, SL ZOLPIMIST	<p><b><u>LENGTH OF AUTHORIZATIONS:</u></b> 180 Days</p> <p><b><u>NON-PREFERRED CRITERIA:</u></b></p> <ul style="list-style-type: none"> <li>Must have had an inadequate clinical response of at least <u>7 days</u> with at least <u>two preferred</u> drugs in this UPDL category and indicated for diagnosis</li> </ul> <p><b><u>ADDITIONAL INFORMATION</u></b></p> <ul style="list-style-type: none"> <li>Non-controlled medications may be authorized if the prescriber indicates the patient has a history of addiction</li> </ul>

Central Nervous System (CNS) Agents: Skeletal Muscle Relaxants, Non-Benzodiazepine		
PREFERRED AGENTS	NON-PREFERRED AGENTS	PA CRITERIA
baclofen susp <sup>AR</sup> , tab chlorzoxazone 500mg cyclobenzaprine IR dantrolene metaxalone 800mg methocarbamol 500, 750mg orphenadrine tizanidine	baclofen soln carisoprodol chlorzoxazone 250, 375, 750mg cyclobenzaprine ER FLEQSUVY <sup>AR</sup> LYVISPAH metaxalone 400mg methocarbamol 1000mg orphenadrine/ASA/caffeine	<p><b><u>LENGTH OF AUTHORIZATIONS:</u></b> 365 Days</p> <p><b><u>NON-PREFERRED CRITERIA:</u></b></p> <ul style="list-style-type: none"> <li>Must have had an inadequate clinical response of at least <u>30 days</u> with at least <u>two preferred</u> drugs in this UPDL category and indicated for diagnosis</li> </ul> <p><b><u>ADDITIONAL CARISOPRODOL (SOMA) CRITERIA:</u></b></p> <ul style="list-style-type: none"> <li>Must provide medical justification that no other muscle relaxant or agent to treat fibromyalgia, or any musculoskeletal condition would serve the clinical needs of the patient</li> </ul> <p><b>AR</b> – FLEQSUVY (baclofen susp): a PA is required for patients 12 years and older</p>

Central Nervous System (CNS) Agents: Smoking Deterrents		
PREFERRED AGENTS	NON-PREFERRED AGENTS	PA CRITERIA
bupropion SR CHANTIX nicotine varenicline		All products are covered without a PA

Dermatologic Agents: Oral Acne Products		
PREFERRED AGENTS	NON-PREFERRED AGENTS	PA CRITERIA
amnesteem <sup>PA</sup> claravis <sup>PA</sup> isotretinoin <sup>PA</sup> zenatane <sup>PA</sup>	ABSORICA ABSORICA LD	<p><b><u>LENGTH OF AUTHORIZATIONS:</u></b> 150 days</p> <p><b><u>CLINICAL PA CRITERIA:</u></b></p> <ul style="list-style-type: none"> <li>Must have had an inadequate clinical response of at least <u>90 days</u> with at least <u>one preferred</u> topical <b>AND</b> <u>one preferred</u> oral antibiotic for acne</li> </ul> <p><b><u>NON-PREFERRED CRITERIA:</u></b></p> <ul style="list-style-type: none"> <li>Must have had an inadequate clinical response of at least <u>90 days</u> with at least <u>two preferred</u> drugs in this UPDL category and indicated for diagnosis</li> </ul> <p><b><u>ADDITIONAL INFORMATION</u></b></p> <ul style="list-style-type: none"> <li>Authorization length will be for no more than 150 days at a time then must take 56 days off</li> </ul>

Dermatologic Agents: Topical Acne Products		
PREFERRED AGENTS	NON-PREFERRED AGENTS	PA CRITERIA
NON-RETINOIDS		<b>LENGTH OF AUTHORIZATIONS:</b> 365 Days  <b>NON-PREFERRED CRITERIA:</b> <ul style="list-style-type: none"><li>Must have had an inadequate clinical response with at least <u>two preferred</u> drugs within the same sub-section classification in this UPDL category. Trials must be 30 days for preferred non-retinoids and 90 days for preferred retinoids.</li></ul> <b>ADDITIONAL CLINDAMYCIN/ADAPALENE/BENZOYL PEROXIDE (CABTREO) CRITERIA</b> <ul style="list-style-type: none"><li>Must provide documentation for patient’s inability to use the individual drugs in this UPDL category</li></ul> <b>ADDITIONAL INFORMATION</b> <ul style="list-style-type: none"><li>All retinoids - May be authorized with a diagnosis of skin cancer</li><li>tazarotene (TAZORAC) - May be authorized with a diagnosis of psoriasis</li></ul> <b>AR</b> - All topical retinoids: a PA is required for patients 24 years and older
azelaic acid gel benzoyl peroxide clindamycin gel, lot, soln, swabs clind/benz perox 1-5%, 1.2-2.5%, 1.2-5% erythromycin erythromycin/benzoyl peroxide ONEXTON GEL <sup>BvG</sup> sodium sulfacetamide gel, liq	CLINDACIN KIT clindamycin foam clindamycin/benz perox 1.2-3.75% dapson gel FINACEA FOAM NEUAC sodium sulfacetamide/sulfur sodium sulfacetamide pads WINLEVI ZMA CLEAR SUSP	
RETINOIDS/COMBINATIONS		
adapalene gel <sup>AR</sup> 0.1%, 0.3% adapalene/benzoyl peroxide <sup>AR</sup> ALTRENO <sup>AR</sup> RETIN-A MICRO <sup>AR BvG</sup> 0.04%, 0.1% tretinoin <sup>AR</sup> cream, gel	adapalene cream <sup>AR</sup> ARAZLO <sup>AR</sup> CABTREO GEL <sup>AR</sup> clindamycin/tretinoin <sup>AR</sup> RETIN-A MICRO <sup>AR</sup> 0.06%, 0.08% <sup>BvG</sup> tazarotene <sup>AR</sup> cream, foam, gel 0.1% tretinoin micro <sup>AR</sup> 0.04%, 0.1%	

## Endocrine Agents: Androgens

PREFERRED AGENTS	NON-PREFERRED AGENTS	PA CRITERIA
depo-testosterone <sup>AR PA</sup> testosterone cypionate <sup>AR PA</sup> testosterone gel 1% packet <sup>AR PA</sup> testosterone gel 1.62% pump <sup>AR PA</sup>	AVEED <sup>AR</sup> AZMIRO <sup>AR</sup> JATENZO <sup>AR</sup> methyltestosterone <sup>AR</sup> NATESTO <sup>AR</sup> TESTOPEL <sup>AR</sup> testosterone gel 1% pump <sup>AR</sup> testosterone gel 1.62% packet <sup>AR</sup> testosterone gel 2% <sup>AR</sup> testosterone soln 30mg/ACT <sup>AR</sup> TLANDO <sup>AR</sup> XYOSTED <sup>AR</sup>	<p><b><u>LENGTH OF AUTHORIZATIONS:</u></b> 365 Days</p> <p><b><u>CLINICAL PA CRITERIA:</u></b></p> <ul style="list-style-type: none"> <li>Must provide documentation of baseline lab work to support the need for testosterone supplementation. If baseline testosterone level is within normal limits, provide clinical justification for why replacement therapy is required.</li> </ul> <p><b><u>NON-PREFERRED CRITERIA:</u></b></p> <ul style="list-style-type: none"> <li>Must have had an inadequate clinical response of at least <u>90 days</u> with <b><u>ALL preferred</u></b> drugs in this UPDL category and indicated for diagnosis</li> </ul> <p><b><u>ADDITIONAL TESTOSTERONE ENANTHATE (XYOSTED) CRITERIA:</u></b></p> <ul style="list-style-type: none"> <li>Must have a trial and failure of a preferred testosterone cypionate injectable product <b>OR</b></li> <li>Must provide a clinical rationale why testosterone cypionate injectable product is not appropriate</li> </ul> <p><b><u>SUBSEQUENT AUTHORIZATION CRITERIA:</u></b></p> <ul style="list-style-type: none"> <li>Must provide documentation of patient's clinical response to treatment and ongoing safety monitoring (i.e., testosterone and hematocrit)</li> </ul> <p><b>AR:</b> All drugs: a PA is required for patients younger than 18 years</p>



Endocrine Agents: Diabetes – Hypoglycemia Treatments		
PREFERRED AGENTS	NON-PREFERRED AGENTS	PA CRITERIA
BAQSIMI glucagon emerg kit [labeler 00548] GVOKE ZEGALOGUE	glucagon emerg kit [labeler 00378, 63323]	<p><b><u>LENGTH OF AUTHORIZATIONS:</u></b> 365 Days</p> <p><b><u>NON-PREFERRED CRITERIA:</u></b></p> <ul style="list-style-type: none"> <li>Must have had an inadequate clinical response of at least <u>one preferred</u> drug in this UPDL category and indicated for diagnosis <b>OR</b> the inability of the member and/or caregiver to administer a preferred glucagon product in a timely fashion</li> </ul> <p><b><u>SUBSEQUENT AUTHORIZATION CRITERIA:</u></b></p> <ul style="list-style-type: none"> <li>Renewal will be allowed for expired/unused products <b>WITHOUT</b> documentation of patient's clinical response to treatment</li> </ul>

## Endocrine Agents: Diabetes – Insulin

PREFERRED AGENTS	NON-PREFERRED AGENTS	PA CRITERIA
RAPID-ACTING		<b>LENGTH OF AUTHORIZATIONS:</b> 365 Days  <b>STEP THERAPY CRITERIA:</b> <ul style="list-style-type: none"><li>Must have had an inadequate clinical response (defined as the inability to reach target A1C) after at least <u>120 days</u> with at least <u>one preferred</u> drug having a similar duration of action in this UPDL category</li></ul> <b>NON-PREFERRED CRITERIA:</b> <ul style="list-style-type: none"><li>Must have had an inadequate clinical response (defined as the inability to reach target A1C) after at least <u>120 days</u> with at least <u>two preferred</u> drugs having a similar duration of action in this UPDL category and indicated for diagnosis, if available</li></ul> <b>ADDITIONAL TEMPO PEN CRITERIA</b> <ul style="list-style-type: none"><li>Must have had an inadequate clinical response or documentation of medical necessity beyond convenience for why the patient cannot use the corresponding FlexPens or Kwikpens</li></ul> <b>ADDITIONAL INHALED INSULIN (AFREZZA) CRITERIA:</b> <ul style="list-style-type: none"><li>Must provide documentation of spirometry testing prior to initiation with a predicted FEV1 ≥70% - Will not be authorized for patients with asthma or COPD</li><li>Must provide documentation of being nicotine-free for at least 180 days</li></ul> <b>ADDITIONAL INFORMATION</b> <ul style="list-style-type: none"><li>An inadequate clinical response is defined as the inability to reach A1C goal after at least 120 days of current regimen with documented adherence and appropriate dose escalation.<ul style="list-style-type: none"><li>Must include a patient specific A1C goal if less than 7%</li><li>Must include current A1C (within last 6 months)</li></ul></li><li>Requests may be authorized for patients with a condition that is difficult to control (i.e., prone to ketoacidosis, hypoglycemia)</li></ul>
APIDRA HUMALOG U-100 KWIKPEN, VIAL insulin aspart insulin lispro	ADMELOG AFREZZA FIASP HUMALOG U-100 TEMPO PEN HUMALOG U-200 LYUMJEV NOVOLOG U-100	
SHORT-ACTING		
HUMULIN R U-500	HUMULIN R U-100 NOVOLIN R U-100	
INTERMEDIATE-ACTING		
	HUMULIN N U-100 NOVOLIN N U-100	
LONG-ACTING		
LANTUS <sup>BvG</sup> LEVEMIR TOUJEO <sup>BvG</sup> TRESIBA <sup>BvG ST</sup>	BASAGLAR insulin degludec insulin glargine REZVOGLAR SEMGLEE <sup>BvG</sup>	
MIXED INSULIN		
HUMALOG 50-50 HUMALOG 75-25 HUMULIN 70-30 insulin aspart pro/insulin aspart	NOVOLIN 70-30 NOVOLOG 70-30	

		<b><u>SUBSEQUENT AUTHORIZATION CRITERIA:</u></b>
--	--	--

- |  |  |  |
|--|--|--|
|  |  | <ul style="list-style-type: none"><li>• Must provide documentation of patient's clinical response to treatment and ongoing safety monitoring<ul style="list-style-type: none"><li>○ Must include a patient specific A1C goal if less than 7%</li><li>○ Must include current A1C (within last 6 months)</li></ul></li></ul> |
|--|--|--|

Endocrine Agents: Diabetes – Non-Insulin		
PREFERRED AGENTS	NON-PREFERRED AGENTS	PA CRITERIA
DPP4 INHIBITORS/COMBINATIONS		LENGTH OF AUTHORIZATIONS: 365 Days  NON-PREFERRED CRITERIA: <ul style="list-style-type: none"><li>Must have had an inadequate clinical response of at least <u>120 days</u> with at least <u>three preferred</u> drugs in this UPDL category and indicated for diagnosis, if available</li></ul> ADDITIONAL SITAGLIPTIN (ZITUVIO) CRITERIA <ul style="list-style-type: none"><li>Must have had a trial of at least 120 days with JANUVIA <b>OR</b> must provide documentation of medical necessity for patient’s inability to use JANUVIA</li></ul> ADDITIONAL INFORMATION <ul style="list-style-type: none"><li>An inadequate clinical response is defined as the inability to reach A1C goal after at least 120 days of current regimen, with use of two or more drugs concomitantly per ADA guidelines, documented adherence, and appropriate dose escalation (must achieve maximum recommended dose or document that maximum recommended dose is not tolerated or is clinically inappropriate).<ul style="list-style-type: none"><li>Must include a patient specific A1C goal if less than 7%</li><li>Must include current A1C (within last 6 months)</li></ul></li><li>For non-preferred drugs that have preferred drugs in the same drug class: must provide documentation that there was at least <u>one</u> inadequate clinical response with a drug in same drug class</li></ul> SUBSEQUENT AUTHORIZATION CRITERIA: <ul style="list-style-type: none"><li>Must provide documentation of patient’s clinical response to treatment and ongoing safety monitoring<ul style="list-style-type: none"><li>Must include a patient specific A1C goal if less than 7%</li><li>Must include current A1C (within last 6 months)</li></ul></li><li>Must meet all initial clinical criteria for subsequent authorizations.</li></ul>
JANUMET JANUMET XR JANUVIA JENTADUETO KOMBIGLYZE XR <sup>BvG</sup> ONGLYZA <sup>BvG</sup> TRADJENTA	alogliptin alogliptin/metformin alogliptin/pioglitazone JENTADUETO XR saxagliptin saxagliptin/metformin sitagliptin/metformin (gen of ZITUVIMET) ZITUVIMET XR ZITUVIO <sup>BvG</sup>	
GLP-1 RECEPTOR AGONISTS/COMBINATIONS		
BYETTA TRULICITY VICTOZA <sup>BvG</sup>	BYDUREON BCISE liraglutide MOUNJARO OZEMPIC RYBELSUS SOLIQUA XULTOPHY	
METFORMIN		
metformin ER (gen of GLUCOPHAGE XR) metformin IR 500, 850, 1000mg	metformin ER (gen of FORTAMET, GLUMETZA) metformin IR 625, 750mg metformin soln	
SGLT2 INHIBITORS/COMBINATIONS		
FARXIGA <sup>BvG</sup> JARDIANCE SYNJARDY XIGDUO XR <sup>BvG</sup>	dapagliflozin dapagliflozin/metformin ER GLYXAMBI INVOKAMET INVOKANA QTERN SEGLUROMET STEGLATRO STEGLUJAN SYNJARDY XR	

	TRIJARDY XR	
SULFONYLUREAS/COMBINATIONS		
glimepiride 1, 2, 4mg	glimepiride 3mg	
glipizide IR, ER	glimepiride/pioglitazone	
glipizide/metformin		
glyburide		
glyburide/metformin		
OTHER		
acarbose	SYMLINPEN	
miglitol		
nateglinide		
pioglitazone		
pioglitazone/metformin		
repaglinide		

## Endocrine Agents: Endometriosis

PREFERRED AGENTS	NON-PREFERRED AGENTS	PA CRITERIA
danazol <sup>ST</sup> DEPO-SUBQ PROVERA 104 <sup>ST</sup> LUPRON DEPOT <sup>ST</sup> 3.75, 11.25mg MYFEMBREE <sup>ST</sup> ORILISSA <sup>ST</sup>	SYNAREL	<p><b><u>LENGTH OF AUTHORIZATIONS:</u></b> 365 Days</p> <p><b><u>STEP THERAPY CRITERIA:</u></b></p> <ul style="list-style-type: none"> <li>Must have had an inadequate clinical response of at least <u>84 days</u> with at least <u>one preferred</u> NSAID and <u>one</u> oral contraceptive</li> </ul> <p><b><u>NON-PREFERRED CRITERIA:</u></b></p> <ul style="list-style-type: none"> <li>Must have had an inadequate clinical response of at least <u>84 days</u> with at least <u>one preferred</u> step-therapy drug in this UPDL category and indicated for diagnosis</li> </ul> <p><b><u>ADDITIONAL INFORMATION:</u></b></p> <ul style="list-style-type: none"> <li>A total lifetime duration of therapy of 730 days between ORILISSA and MYFEMBREE or 365 days for LUPRON DEPOT will be authorized</li> </ul>

Endocrine Agents: Estrogenic Agents		
PREFERRED AGENTS	NON-PREFERRED AGENTS	PA CRITERIA
ORAL		<b>LENGTH OF AUTHORIZATIONS:</b> 365 Days  <b>NON-PREFERRED CRITERIA:</b> <ul style="list-style-type: none"><li>Must have had an inadequate clinical response of at least <u>30 days</u> with at least <u>two preferred</u> drugs in this UPDL category within the same sub-section classification and indicated for diagnosis</li></ul>
ANGELIQ estradiol tab ethinyl estradiol/norethindrone PREMARIN TAB PREMPHASE PREMPRO	DUAVEE estradiol/norethindrone MENEST	
TOPICAL		
DIVIGEL <sup>BvG</sup> ELESTRIN estradiol cream	estradiol gel 0.06% (gen of ESTROGEL)	
TRANSDERMAL		
CLIMARA COMBIPATCH dotti estradiol patch lyllana MINIVELLE VIVELLE -DOT	EVAMIST MENOSTAR	
VAGINAL		
ESTRING PREMARIN CREAM	estradiol 10mcg vag tab FEMRING	

Endocrine Agents: Growth Hormone		
PREFERRED AGENTS	NON-PREFERRED AGENTS	PA CRITERIA
DAILY-DOSING		<b>LENGTH OF AUTHORIZATIONS:</b> Initial: 180 days; Subsequent: 365 days  <b>CLINICAL PA CRITERIA:</b> <b>Pediatric Approvals (under 18 years of age):</b> <ul style="list-style-type: none"><li>• Must be treated and followed by a pediatric endocrinologist, nephrologist, clinical geneticist, endocrinologist, or gastroenterologist (or as appropriate for diagnosis)</li><li>• Must provide documentation to justify criteria being met, including height, weight, bone age (children), date and results of most current x-ray, stimulus test results, IGF-1 levels, and a growth chart (children)</li><li>• Must not be used in combination with another somatropin agent</li></ul> <b>Adult Approvals (18 years of age or older):</b> <ul style="list-style-type: none"><li>• Must be treated and followed by an endocrinologist</li><li>• Must provide documentation of growth hormone deficiency by means of a negative response to an appropriate stimulation test (clonidine test is not acceptable for adults)</li></ul> <b>STEP THERAPY CRITERIA:</b> <ul style="list-style-type: none"><li>• Must have had an inadequate clinical response of at least <u>90 days</u> with at least <u>one preferred</u> daily-dosed growth hormone formulation</li></ul> <b>NON-PREFERRED CRITERIA:</b> <ul style="list-style-type: none"><li>• Must have had an inadequate clinical response of at least <u>90 days</u> with at least <u>one preferred</u> drug within the same sub-section classification in this UPDL category and indicated for diagnosis</li></ul> <b>SUBSEQUENT AUTHORIZATION CRITERIA:</b> <ul style="list-style-type: none"><li>• Must provide documentation of patient’s clinical response to treatment and ongoing safety monitoring (i.e., height, weight gain, improved body composition)</li><li>• For adults: must provide documentation by endocrinologist that discontinuing agent would have a detrimental effect on body composition or other metabolic parameters</li></ul>
GENOTROPIN <sup>PA</sup> NORDITROPIN <sup>PA</sup>	HUMATROPE NUTROPIN OMNITROPE SEROSTIM ZOMACTON	
WEEKLY-DOSING		
SKYTROFA <sup>PA ST</sup>	NGENLA SOGROYA	



Endocrine Agents: Osteoporosis – Bone Ossification Enhancers		
PREFERRED AGENTS	NON-PREFERRED AGENTS	PA CRITERIA
<b>BISPHOSPHONATES</b>		<b>LENGTH OF AUTHORIZATIONS:</b> 365 Days
alendronate tab ibandronate	alendronate soln BINOSTO FOSAMAX PLUS D risedronate zoledronic acid	<b>CLINICAL PA CRITERIA:</b> <ul style="list-style-type: none"> <li>• Must have had an inadequate clinical response of at least <u>365 days</u> with <u>one</u> bisphosphonate</li> <li>• A total lifetime duration of therapy of 730 days will be authorized between any parathyroid analog</li> </ul>
<b>OTHER BONE RESORPTION SUPPRESSION AND RELATED AGENTS</b>		
calcitonin-salmon FORTEO <sup>BvG PA</sup> raloxifene	EVENITY PROLIA teriparatide TYMLOS	<b>NON-PREFERRED CRITERIA:</b> <ul style="list-style-type: none"> <li>• Must have had an inadequate clinical response of at least <u>365 days</u> with at least <u>one preferred</u> drug within the same sub-section classification in this UPDL category and indicated for diagnosis</li> </ul> <b>ADDITIONAL “OTHER BONE RESORPTION SUPPRESSION AND RELATED AGENTS” CRITERIA:</b> <ul style="list-style-type: none"> <li>• Must have had an inadequate clinical response of at least <u>365 days</u> with <u>one</u> bisphosphonate</li> <li>• A total lifetime duration of therapy of 730 days will be authorized between any parathyroid analog</li> <li>• A total lifetime duration of therapy of 365 days will be authorized for EVENITY</li> </ul> <b>ADDITIONAL INFORMATION</b> <ul style="list-style-type: none"> <li>• Patients should only be on ONE of the therapeutic classes (bisphosphonates, calcitonin-salmon)</li> </ul>

Endocrine Agents: Progestin Agents		
PREFERRED AGENTS	NON-PREFERRED AGENTS	PA CRITERIA
medroxyprogesterone acetate tab megestrol norethindrone acetate progesterone progesterone in oil		All products are covered without a PA

Endocrine Agents: Uterine Fibroids		
PREFERRED AGENTS	NON-PREFERRED AGENTS	PA CRITERIA
LUPRON DEPOT <sup>PA</sup> 3.75, 11.25mg MYFEMBREE <sup>PA</sup> ORIAHNN <sup>PA</sup>		<p><b><u>LENGTH OF AUTHORIZATIONS:</u></b> Up to 180 Days</p> <p><b><u>CLINICAL PA CRITERIA:</u></b></p> <ul style="list-style-type: none"> <li>• Must have had an inadequate clinical response of at least 90 days with at least one oral contraceptive</li> </ul> <p><b><u>ADDITIONAL INFORMATION:</u></b></p> <ul style="list-style-type: none"> <li>• A total lifetime duration of therapy of 730 days between MYFEMBREE and ORIAHNN or 365 days for LUPRON DEPOT will be authorized</li> </ul>

Gastrointestinal Agents: Anti-Emetics		
PREFERRED AGENTS	NON-PREFERRED AGENTS	PA CRITERIA
5-HT3 ANTAGONISTS		<b>LENGTH OF AUTHORIZATIONS:</b> 365 Days  <b>CLINICAL PA CRITERIA:</b> <ul style="list-style-type: none"><li>dronabinol is only covered for nausea and vomiting associated with chemotherapy in adult patients who failed at least <u>3 days</u> with at least <u>one preferred</u> drug in this UPDL category.</li></ul> <b>NON-PREFERRED CRITERIA:</b> <ul style="list-style-type: none"><li>Must have had an inadequate clinical response of at least <u>3 days</u> with at least <u>one preferred</u> drug in this UPDL category within the same sub-section classification and indicated for diagnosis</li></ul>
granisetron tab ondansetron 4, 8mg	ondansetron 16mg SANCUSO	
ANTICHOLINERGICS		
scopolamine		
ANTI-HISTAMINES and ANTI-HISTAMINE COMBINATIONS		
dimenhydrinate diphenhydramine doxylamine/pyridoxine meclizine trimethobenzamide	BONJESTA	
PHENOTHIAZINES		
prochlorperazine promethazine		
SUBSTANCE P/NEUROKININ 1 (NK-1) ANTAGONISTS		
aprepitant 40mg, tripac EMEND 125mg SUSP	aprepitant 80, 125mg	
OTHER		
dronabinol <sup>PA</sup> metoclopramide		

Gastrointestinal Agents: Bowel Preparations		
PREFERRED AGENTS	NON-PREFERRED AGENTS	PA CRITERIA
CLENPIQ GAVILYTE -C GAVILYTE -G GAVILYTE -N GOLYTELY MOVIPREP sod sulf-potass sulf-mag sulf soln	PLENVU SUFLAVE SUTAB	<p><b><u>LENGTH OF AUTHORIZATIONS:</u></b> 365 Days</p> <p><b><u>NON-PREFERRED CRITERIA:</u></b></p> <ul style="list-style-type: none"> <li>• Must have had an inadequate clinical response with at least <u>one preferred</u> drug in this UPDL category and indicated for diagnosis</li> </ul>

Gastrointestinal Agents: Crohn's Disease		
PREFERRED AGENTS	NON-PREFERRED AGENTS	PA CRITERIA
azathioprine 50mg budesonide ER mercaptopurine tab methotrexate sulfasalazine IR, DR	azathioprine 75, 100mg	<p><b><u>LENGTH OF AUTHORIZATIONS:</u></b> 365 days</p> <p><b><u>NON-PREFERRED CRITERIA:</u></b></p> <ul style="list-style-type: none"> <li>Must have had an inadequate clinical response of at least <u>30 days</u> with at least <u>two preferred</u> drugs in this UPDL category and indicated for diagnosis</li> </ul>

Gastrointestinal Agents: Hepatic Encephalopathy		
PREFERRED AGENTS	NON-PREFERRED AGENTS	PA CRITERIA
lactulose XIFAXAN <sup>ST</sup>		<p><b><u>LENGTH OF AUTHORIZATIONS:</u></b> 365 Days</p> <p><b><u>STEP THERAPY CRITERIA:</u></b></p> <ul style="list-style-type: none"> <li>• Must have had an inadequate clinical response of at least <u>14 days</u> with at least <u>one preferred</u> drug in this UPDL category</li> </ul>

Gastrointestinal Agents: Irritable Bowel Syndrome (IBS) with Diarrhea		
PREFERRED AGENTS	NON-PREFERRED AGENTS	PA CRITERIA
diphenoxylate/atropine loperamide XIFAXAN <sup>ST</sup>	alosetron VIBERZI	<p><b><u>LENGTH OF AUTHORIZATIONS:</u></b> 365 Days</p> <p><b><u>STEP THERAPY CRITERIA:</u></b></p> <ul style="list-style-type: none"> <li>• Must have had an inadequate clinical response of at least <u>14 days</u> with at least <u>one preferred</u> drug in this UPDL category</li> </ul> <p><b><u>NON-PREFERRED CRITERIA:</u></b></p> <ul style="list-style-type: none"> <li>• Must have had an inadequate clinical response of at least <u>14 days</u> with at least <u>one preferred</u> drug and one <u>step therapy</u> drug in this UPDL category and indicated for diagnosis</li> </ul>



### Gastrointestinal Agents: Pancreatic Enzymes

PREFERRED AGENTS	NON-PREFERRED AGENTS	PA CRITERIA
<p>CREON</p> <p>PERTZYE <sup>ST</sup></p> <p>ZENPEP</p>	<p>VIOKACE</p>	<p><b><u>LENGTH OF AUTHORIZATIONS:</u></b> 365 Days</p> <p><b><u>STEP THERAPY CRITERIA:</u></b></p> <ul style="list-style-type: none"> <li>• For a diagnosis of Cystic Fibrosis, no trials required</li> <li>• For all other diagnoses, must have had an inadequate clinical response of at least <u>14 days</u> with at least <u>one preferred</u> drug in this UPDL category</li> </ul> <p><b><u>NON-PREFERRED CRITERIA:</u></b></p> <ul style="list-style-type: none"> <li>• Must have had an inadequate clinical response of at least <u>14 days</u> with at least <u>two preferred</u> drugs in this UPDL category and indicated for diagnosis</li> </ul>

Gastrointestinal Agents: Proton Pump Inhibitors		
PREFERRED AGENTS	NON-PREFERRED AGENTS	PA CRITERIA
esomeprazole lansoprazole cap NEXIUM GRANULES <sup>BvG</sup> omeprazole pantoprazole tab PROTONIX PAK <sup>AR BvG</sup> rabeprazole	DEXILANT <sup>BvG</sup> esomeprazole granules KONVOMEK lansoprazole ODT omeprazole/sodium bicarbonate pantoprazole packet <sup>AR</sup> PRILOSEC SUSP	<p><b><u>LENGTH OF AUTHORIZATIONS:</u></b> 180 days, except as listed under additional criteria</p> <p><b><u>NON-PREFERRED CRITERIA:</u></b></p> <ul style="list-style-type: none"> <li>Must have had an inadequate clinical response of at least <u>30 days</u> with at least <u>two preferred</u> drugs in this UPDL category and indicated for diagnosis</li> </ul> <p><b><u>ADDITIONAL CRITERIA FOR PPI DOSES GREATER THAN ONCE DAILY</u></b></p> <ul style="list-style-type: none"> <li>Must have had an inadequate clinical response of at least <u>30 days</u> of once daily dosing with the requested drug <b>OR</b></li> <li>For H. Pylori diagnosis: Must provide documentation of diagnosis               <ul style="list-style-type: none"> <li>Authorization length: 30 days</li> </ul> </li> <li>For any of the following diagnoses: carcinoma of GI tract, COPD, Crest Syndrome, dyspepsia, esophageal varices, gastritis, gastroparesis, scleroderma, symptomatic uncomplicated Barret's Esophagus, systemic mastocytosis, or Zollinger Ellison Syndrome: Must provide documentation of diagnosis <b>AND</b> must have failed once-daily dosing of the requested drug               <ul style="list-style-type: none"> <li>Authorization length: 365 days</li> </ul> </li> </ul> <p><b><u>ADDITIONAL INFORMATION</u></b></p> <ul style="list-style-type: none"> <li>Request may be authorized If the drug was initiated in the hospital for the treatment of a condition such as a GI bleed or the presence of a gastrostomy and/or jejunostomy (G, GJ, J-tube)</li> </ul> <p><b>AR – PROTONIX PAK/pantoprazole packet:</b> a PA is required for patients 6 years and older</p>

Gastrointestinal Agents: Ulcerative Colitis		
PREFERRED AGENTS	NON-PREFERRED AGENTS	PA CRITERIA
<b>ORAL</b>		<b>LENGTH OF AUTHORIZATIONS:</b> 365 Days; except UCERIS FOAM – 90 days
balsalazide disodium budesonide ER tab mesalamine DR cap, tab 1.2gm mesalamine ER cap 0.375gm, 500mg PENTASA 250mg sulfasalazine IR, DR	DIPENTUM mesalamine DR tab 800mg VELSIPITY ZEPOSIA	<b>NON-PREFERRED CRITERIA:</b> <ul style="list-style-type: none"> <li>Must have had an inadequate clinical response of at least <u>30 days</u> with at least <u>two preferred</u> drugs in this UPDL category within the same sub-section classification and indicated for diagnosis, if available</li> </ul>
<b>RECTAL</b>		<b>ADDITIONAL BUDESONIDE (UCERIS) CRITERIA:</b>
mesalamine enema, supp	mesalamine enema kit SF ROWASA UCERIS FOAM <sup>BvG</sup>	<ul style="list-style-type: none"> <li>Must have had a documented side effect, allergy, or treatment failure of at least <u>30 days</u> with mesalamine enema or suppository</li> </ul>
		<b>ADDITIONAL OZANIMOD (ZEPOSIA) AND ETRASIMOD (VELSIPITY) CRITERIA:</b> <ul style="list-style-type: none"> <li>Must have had a documented side effect, allergy, or treatment failure of at least <u>90 days</u> with at least <u>one preferred Systemic Immunomodulator indicated for Ulcerative Colitis</u> (refer to Immunomodulator Agents: Systemic Inflammatory Disease class for complete list)</li> </ul>

## Gastrointestinal Agents: Unspecified GI

PREFERRED AGENTS	NON-PREFERRED AGENTS	PA CRITERIA
bisacodyl dicyclomine diphenoxylate/atropine lactulose LINZESS loperamide lubiprostone <sup>ST</sup> MOVANTIK <sup>ST</sup> polyethylene glycol oral powder bottle senna TRULANCE <sup>ST</sup> XIFAXAN <sup>ST</sup>	AEMCOLO AMITIZA GATTEX IBSRELA MYTESI polyethylene glycol oral powder packet prucalopride RELISTOR SYMPROIC	<p><b><u>LENGTH OF AUTHORIZATIONS:</u></b> 365 days except 3 days for AEMCOLO</p> <p><b><u>STEP THERAPY CRITERIA:</u></b></p> <ul style="list-style-type: none"> <li>Must have had an inadequate clinical response to at least <u>14 days</u> with at least <u>two preferred</u> drugs in this UPDL category, if indicated for diagnosis</li> </ul> <p><b><u>NON-PREFERRED CRITERIA:</u></b></p> <ul style="list-style-type: none"> <li>Must have had an inadequate clinical response of at least <u>14 days</u> with <u>one step therapy</u> drug this UPDL category and indicated for diagnosis</li> </ul> <p><b><u>ADDITIONAL METHYLNALTREXONE (RELISTOR) AND NALDEMEDINE (SYMPROIC) CRITERIA:</u></b></p> <ul style="list-style-type: none"> <li>Must have a history of chronic pain requiring continuous opioid therapy for ≥84 days</li> </ul> <p><b><u>ADDITIONAL RIFAMYCIN DELAYED-RELEASE (AEMCOLO) CRITERIA:</u></b></p> <ul style="list-style-type: none"> <li>Must have the inability to take, or failure of <b>ALL</b> of the following: azithromycin, ciprofloxacin, levofloxacin, ofloxacin, or rifaximin</li> </ul> <p><b><u>SUBSEQUENT AUTHORIZATION CRITERIA:</u></b></p> <ul style="list-style-type: none"> <li>Must provide documentation of patient's clinical response to treatment and ongoing safety monitoring (i.e., decreased frequency of specialized nutrition support or improvement in symptoms)</li> </ul>

Genitourinary Agents: Benign Prostatic Hyperplasia		
PREFERRED AGENTS	NON-PREFERRED AGENTS	PA CRITERIA
ALPHA BLOCKERS		<b>LENGTH OF AUTHORIZATIONS:</b> 365 Days  <b>TADALAFIL (CIALIS) CRITERIA:</b> <ul style="list-style-type: none"><li>Must have had an inadequate clinical response of at least <u>30 days</u> with at least <u>one</u> alpha-1 adrenergic blocker. If prostate volume of &gt; 30cc on imaging, a prostate specific antigen (PSA) &gt; 1.5ng/dL, or palpable prostate enlargement on digital rectal exam (DRE), then a trial of at least <u>90 days</u> of finasteride is required.</li></ul> <b>NON-PREFERRED CRITERIA:</b> <ul style="list-style-type: none"><li>Must have had an inadequate clinical response of at least <u>60 days</u> with at least <u>two preferred</u> drugs, with at least <u>one preferred</u> within the same sub-section classification and indicated for diagnosis, if available</li></ul>
alfuzosin doxazosin prazosin silodosin tamsulosin terazosin	CARDURA XL	
5-ALPHA-REDUCTASE (5AR) INHIBITORS		
dutasteride finasteride 5mg		
ALPHA BLOCKER/5AR/PDE5 INHIBITOR COMBINATIONS		
	dutasteride/tamsulosin	
PHOSPHODIESTERASE 5 (PDE5) INHIBITORS		
tadalafil <sup>PA</sup> 2.5, 5mg		

Genitourinary Agents: Electrolyte Depleter Agents		
PREFERRED AGENTS	NON-PREFERRED AGENTS	PA CRITERIA
CALCIUM BASED		LENGTH OF AUTHORIZATIONS: 365 Days
calcium acetate, carbonate		
IRON BASED		STEP THERAPY CRITERIA: <ul style="list-style-type: none"><li>Must have had an inadequate clinical response of at least <u>7 days</u> with at least <u>one preferred</u> drug in this UPDL category</li></ul>
VELPHORO <sup>ST</sup>	ferric citrate tab	
OTHER		
sevelamer	FOSRENOL POWDER lanthanum carbonate XPHOZAH	NON-PREFERRED CRITERIA: <ul style="list-style-type: none"><li>Must have had an inadequate clinical response of at least <u>14 days</u> with at least <u>one preferred</u> step therapy drug in this UPDL category and indicated for diagnosis, if available</li></ul>

Genitourinary Agents: Urinary Antispasmodics		
PREFERRED AGENTS	NON-PREFERRED AGENTS	PA CRITERIA
ANTIMUSCARINICS		<u>LENGTH OF AUTHORIZATIONS:</u> 365 Days  <u>NON-PREFERRED CRITERIA:</u> <ul style="list-style-type: none"><li>Must have had an inadequate clinical response of at least <u>30 days</u> with at least <u>two preferred</u> drugs in this UPDL category and indicated for diagnosis, one of which must be within the same sub-section classification, if available</li></ul>  <b>AR – MYRBETRIQ GRANULES:</b> a PA is required for patients younger than 3 years old AND 5 years and older <b>AR – VESICARE LS:</b> a PA is required for patients younger than 2 years old AND 5 years and older
fesoterodine oxybutynin IR, ER OXYTROL solifenacin trospium IR, ER	darifenacin tolterodine IR, ER VESICARE LS <sup>AR</sup>	
BETA-3 AGONISTS		
MYRBETRIQ TAB <sup>BvG</sup>	GEMTESA mirabegron tab MYRBETRIQ GRANULES <sup>AR</sup>	

**AR – MYRBETRIQ GRANULES:** a PA is required for patients younger than 3 years old AND 5 years and older

**AR – VESICARE LS:** a PA is required for patients younger than 2 years old AND 5 years and older

Hyperkalemia Agents: Potassium Binders		
PREFERRED AGENTS	NON-PREFERRED AGENTS	PA CRITERIA
LOKELMA	kionex susp sodium polystyrene sulfonate VELTASSA	<p><b><u>LENGTH OF AUTHORIZATIONS:</u></b> 365 Days</p> <p><b><u>NON-PREFERRED CRITERIA:</u></b></p> <ul style="list-style-type: none"> <li>• Must have had an inadequate clinical response of at least <u>30</u> days with at least <u>one preferred</u> drug in this UPDL category and indicated for diagnosis</li> </ul>



Immunomodulator Agents: Systemic Inflammatory Disease		
PREFERRED AGENTS	NON-PREFERRED AGENTS	PA CRITERIA
INTERLEUKIN ANTAGONISTS		<b>LENGTH OF AUTHORIZATIONS:</b> Initial: 90 days; Subsequent: 365 days  <b>CLINICAL PA CRITERIA:</b> <ul style="list-style-type: none"><li>Authorization of dosing regimens (loading/maintenance) will be based upon diagnosis. Document the requested loading and maintenance dosing on PA form, if applicable</li><li>Must not have a current, active infection</li><li>Must provide evidence of negative TB test prior to initiation of biologic therapy, if required by labeling</li></ul> <b>STEP THERAPY CRITERIA:</b> <ul style="list-style-type: none"><li>Must have had an inadequate clinical response of at least <u>90 days</u> with at least <u>one preferred</u> TNF inhibitor indicated for diagnosis in this UPDL category</li></ul> <b>NON-PREFERRED CRITERIA:</b> <ul style="list-style-type: none"><li>Must have had an inadequate clinical response of at least <u>90 days</u> with at least <u>two preferred</u> drugs in this UPDL category that are not biosimilars of the same reference product and indicated for diagnosis<ul style="list-style-type: none"><li>For non-preferred immunomodulators: must provide documentation of inadequate clinical response to its preferred reference product or biosimilar, in this UPDL category and indicated for the diagnosis, if available</li></ul></li></ul> <b>ADDITIONAL ALOPECIA AREATA CRITERIA:</b> <ul style="list-style-type: none"><li>Must be prescribed by or in consultation with a specialist (i.e., dermatologist, rheumatologist)</li><li>Must provide documentation of an inadequate clinical response of at least 90 days with a topical steroid</li></ul>
ADBRY <sup>PA</sup> DUPIXENT <sup>PA</sup> EBGLYSS <sup>PA</sup> KINERET <sup>PA</sup> SKYRIZI INJ <sup>PA</sup> TALTZ <sup>PA ST</sup> TREMFA <sup>PA</sup> TYENNE <sup>PA</sup> (Bio of ACTEMRA)	ACTEMRA BIMZELX COSENTYX ILUMYA KEVZARA NEMLUVIO OMVOH SILIQ SKYRIZI IV SOLN STELARA	
JAK INHIBITORS		
RINVOQ <sup>PA</sup> XELJANZ IR <sup>PA</sup>	CIBINQO LITFULO OLUMIANT XELJANZ SOLN, XR	
TNF INHIBITORS		
adalimumab-adaz <sup>PA</sup> (gen of HYRIMOZ) adalimumab-fkjp <sup>PA</sup> (gen of HULIO) AMJEVITA <sup>PA</sup> 10/0.1ml (Bio of HUMIRA) ENBREL <sup>PA</sup> HUMIRA <sup>PA</sup> INFLECTRA <sup>PA</sup> (Bio of REMICADE) SIMLANDI <sup>BvG PA</sup> (Bio of HUMIRA)	ABRILADA (Bio of HUMIRA) adalimumab-aacf (gen of IDACIO) adalimumab-aaty (gen of YUFLYMA) adalimumab-adbm (gen of CYLTEZO) adalimumab-ryvk (gen of SIMLANDI) AMJEVITA 10/0.2ml (Bio of HUMIRA) AVSOLA (Bio of REMICADE) CIMZIA HADLIMA (Bio of HUMIRA) HYRIMOZ (Bio of HUMIRA) infliximab (gen of REMICADE) RENFLEXIS (Bio of REMICADE) SIMPONI YUSIMRY (Bio of HUMIRA)	

	ZYMFENTRA	<b><u>ADDITIONAL ATOPIC DERMATITIS CRITERIA:</u></b> <ul style="list-style-type: none"> <li>• Must have at least 10% body surface area (BSA) involvement with an inadequate clinical response of at least <u>90 days</u> with <u>two</u> of the following: topical corticosteroids or topical calcineurin inhibitors [e.g., ELIDEL] unless atopic dermatitis is severe and involves &gt;25% BSA</li> </ul> <b><u>ADDITIONAL PRURIGO NODULARIS CRITERIA:</u></b> <ul style="list-style-type: none"> <li>• Must be prescribed by or in consultation with a specialist (i.e., dermatologist, rheumatologist)</li> <li>• Must provide documentation of an inadequate clinical response of at least 90 days with a topical steroid</li> </ul>
	<b>OTHER</b>	
OTEZLA <sup>PA</sup>	ENTYVIO ORENCIA SOTYKTU	

Infectious Disease Agents: Antibiotics – Cephalosporins		
PREFERRED AGENTS	NON-PREFERRED AGENTS	PA CRITERIA
cefaclor IR cefaclor susp <sup>AR</sup> cefadroxil cefdinir cefprozil cefprozil susp <sup>AR</sup> cefuroxime cephalixin cap 250, 500mg cephalixin susp <sup>AR</sup>	cefaclor ER cefixime cap cefixime susp <sup>AR</sup> cefpodoxime cephalixin cap 750mg, tab	<p><b><u>LENGTH OF AUTHORIZATIONS:</u></b> Based on indication</p> <p><b><u>NON-PREFERRED CRITERIA:</u></b></p> <ul style="list-style-type: none"> <li>Must have had an inadequate clinical response of at least <u>3 days</u> with at least <u>one preferred</u> drug in this UPDL category and indicated for diagnosis</li> </ul> <p><b><u>ADDITIONAL INFORMATION</u></b></p> <ul style="list-style-type: none"> <li>Requests may be authorized if the patient is completing a course of therapy that was started in the hospital or other similar location or was started before Medicaid eligibility, only the remaining course will be authorized</li> </ul> <p><b><u>SUBSEQUENT AUTHORIZATION CRITERIA:</u></b></p> <ul style="list-style-type: none"> <li>Must provide documentation of patient's clinical response to treatment, ongoing safety monitoring, <b>AND</b> medical necessity for continued use</li> </ul> <p><b>AR</b> – cefaclor susp: a PA is required for patients 12 years and older  <b>AR</b> – cefixime susp: a PA is required for patients 12 years and older  <b>AR</b> – cefprozil susp: a PA is required for patients 12 years and older  <b>AR</b> – cephalixin susp: a PA is required for patients 12 years and older</p>

### Infectious Disease Agents: Antibiotics – Inhaled

PREFERRED AGENTS	NON-PREFERRED AGENTS	PA CRITERIA
tobramycin 300mg/5ml neb soln <sup>PA</sup> tobramycin inj	ARIKAYCE CAYSTON TOBI PODHALER tobramycin 300mg/4ml neb soln	<p><b><u>LENGTH OF AUTHORIZATIONS:</u></b> Initial: 180 days; Subsequent: 365 days</p> <p><b><u>CLINICAL PA CRITERIA:</u></b></p> <ul style="list-style-type: none"> <li>• Must provide documentation of cultures demonstrating drug is prescribed in alignment with approved indication</li> </ul> <p><b><u>NON-PREFERRED CRITERIA:</u></b></p> <ul style="list-style-type: none"> <li>• Must have had an inadequate clinical response of at least <u>28 days</u> with at least <u>one preferred</u> drug in this UPDL category and indicated for diagnosis</li> </ul> <p><b><u>SUBSEQUENT AUTHORIZATION CRITERIA:</u></b></p> <ul style="list-style-type: none"> <li>• Must provide documentation of patient’s clinical response to treatment and ongoing safety monitoring (i.e., culture conversion, symptom improvement)</li> </ul>

Infectious Disease Agents: Antibiotics – Macrolides		
PREFERRED AGENTS	NON-PREFERRED AGENTS	PA CRITERIA
azithromycin clarithromycin IR, susp <sup>AR</sup>	clarithromycin ER erythromycin IR, ER	<p><b><u>LENGTH OF AUTHORIZATIONS:</u></b> Based on indication</p> <p><b><u>NON-PREFERRED CRITERIA:</u></b></p> <ul style="list-style-type: none"> <li>Must have had an inadequate clinical response of at least <u>3 days</u> with at least <u>one preferred</u> drug in this UPDL category and indicated for diagnosis</li> </ul> <p><b><u>ADDITIONAL INFORMATION</u></b></p> <ul style="list-style-type: none"> <li>Requests may be authorized if the patient is completing a course of therapy that was started in the hospital or other similar location or was started before Medicaid eligibility, only the remaining course will be authorized</li> </ul> <p><b><u>SUBSEQUENT AUTHORIZATION CRITERIA:</u></b></p> <ul style="list-style-type: none"> <li>Must provide documentation of patient's clinical response to treatment, ongoing safety monitoring, <b>AND</b> medical necessity for continued use</li> </ul> <p><b>AR</b> – clarithromycin susp: a PA is required for patients 12 years and older</p>

Infectious Disease Agents: Antibiotics – Quinolones		
PREFERRED AGENTS	NON-PREFERRED AGENTS	PA CRITERIA
CIPRO ORAL SUSP <sup>AR</sup> ciprofloxacin ciprofloxacin susp <sup>AR</sup> levofloxacin soln <sup>AR</sup> , tab moxifloxacin	BAXDELA ofloxacin	<p><b><u>LENGTH OF AUTHORIZATIONS:</u></b> Based on indication</p> <p><b><u>NON-PREFERRED CRITERIA:</u></b></p> <ul style="list-style-type: none"> <li>Must have had an inadequate clinical response of at least <u>3 days</u> with at least <u>one preferred</u> drug in this UPDL category and indicated for diagnosis</li> </ul> <p><b><u>ADDITIONAL INFORMATION</u></b></p> <ul style="list-style-type: none"> <li>Requests may be authorized if the patient is completing a course of therapy that was started in the hospital or other similar location or was started before Medicaid eligibility, only the remaining course will be authorized</li> </ul> <p><b><u>SUBSEQUENT AUTHORIZATION CRITERIA:</u></b></p> <ul style="list-style-type: none"> <li>Must provide documentation of patient's clinical response to treatment, ongoing safety monitoring, <b>AND</b> medical necessity for continued use</li> </ul> <p><b>AR</b> – ciprofloxacin susp: a PA is required for patients 12 years and older  <b>AR</b> – CIPRO ORAL SUSP: a PA is required for patients 12 years and older  <b>AR</b> – levofloxacin oral soln: a PA is required for patients 12 years and older</p>

### Infectious Disease Agents: Antibiotics – Tetracyclines

PREFERRED AGENTS	NON-PREFERRED AGENTS	PA CRITERIA
doxycycline 20, 50, 100mg doxycycline susp <sup>AR</sup> minocycline IR tetracycline	demeclocycline doxycycline 75, 150mg doxycycline DR minocycline ER MINOLIRA NUZYRA	<p><b><u>LENGTH OF AUTHORIZATIONS:</u></b> Based on indication for acute infections or 365 days for acne</p> <p><b><u>NON-PREFERRED CRITERIA:</u></b></p> <ul style="list-style-type: none"> <li>Must have had an inadequate clinical response of at least <u>3 days</u> with at least <u>one preferred</u> drug for acute infections <b>OR</b> at least <u>90 days</u> with at least <u>one preferred oral</u> drug for acne in this UPDL category and indicated for diagnosis</li> </ul> <p><b><u>ADDITIONAL INFORMATION</u></b></p> <ul style="list-style-type: none"> <li>Requests may be authorized if the patient is completing a course of therapy that was started in the hospital or other similar location or was started before Medicaid eligibility, only the remaining course will be authorized</li> </ul> <p><b><u>SUBSEQUENT AUTHORIZATION CRITERIA:</u></b></p> <ul style="list-style-type: none"> <li>Must provide documentation of patient's clinical response to treatment, ongoing safety monitoring, <b>AND</b> medical necessity for continued use</li> </ul> <p><b>AR</b> – doxycycline susp: a PA is required for patients 12 years and older</p>

Infectious Disease Agents: Antifungals		
PREFERRED AGENTS	NON-PREFERRED AGENTS	PA CRITERIA
clotrimazole fluconazole griseofulvin itraconazole cap ketoconazole nystatin terbinafine voriconazole susp <sup>AR</sup> , tab	BREXAFEMME CRESEMBA flucytosine itraconazole soln NOXAFIL PAK ORAVIG posaconazole TOLSURA VIVJOA	<p><b><u>LENGTH OF AUTHORIZATIONS:</u></b> Based on indication</p> <p><b><u>NON-PREFERRED CRITERIA:</u></b></p> <ul style="list-style-type: none"> <li>Must have had an inadequate clinical response of at least <u>3 days</u> with at least <u>two preferred</u> drugs in this UPDL category and indicated for diagnosis</li> </ul> <p><b><u>ADDITIONAL OTESECONAZOLE (VIVJOA) CRITERIA:</u></b></p> <ul style="list-style-type: none"> <li>Must provide documentation of at least three symptomatic episodes of vulvovaginal candidiasis in the past 12 months</li> <li>Must provide documentation of non-reproductive potential (i.e., post-menopausal)</li> <li>Must have had an inadequate clinical response of at least <u>180-day</u> maintenance course with oral fluconazole shown by documentation of more than <u>one</u> breakthrough infection</li> </ul> <p><b><u>ADDITIONAL INFORMATION:</u></b></p> <ul style="list-style-type: none"> <li>posaconazole can be approved for aspergillosis treatment and prophylaxis without trials of preferred agents</li> <li>Requests may be authorized if the patient is completing a course of therapy that was started in the hospital or other similar location or was started before Medicaid eligibility, only the remaining course will be authorized</li> </ul> <p><b><u>SUBSEQUENT AUTHORIZATION CRITERIA:</u></b></p> <ul style="list-style-type: none"> <li>Must provide documentation of patient's clinical response to treatment, ongoing safety monitoring, <b>AND</b> medical necessity for continued use</li> </ul> <p><b>AR</b> – voriconazole susp: a PA is required for patients 12 years and older</p>



Infectious Disease Agents: Antivirals – Hepatitis C Agents		
PREFERRED AGENTS	NON-PREFERRED AGENTS	PA CRITERIA
MAVYRET <sup>PA</sup> PEGASYS <sup>PA</sup> ribavirin <sup>PA</sup> sofosbuvir/velpatasvir <sup>PA</sup>	HARVONI ledipasvir/sofosbuvir SOVALDI VOSEVI ZEPATIER	<p><b><u>LENGTH OF AUTHORIZATIONS:</u></b> Dependent upon authorized course</p> <p><b><u>CLINICAL PA CRITERIA:</u></b></p> <ul style="list-style-type: none"> <li>Only regimens recommended by the American Association for the Study of Liver Diseases (AASLD) will be authorized</li> <li>Please see the <a href="#">Hepatitis C Direct Acting Antiviral Prior Authorization Form</a> for criteria</li> </ul> <p><b><u>NON-PREFERRED CRITERIA:</u></b></p> <ul style="list-style-type: none"> <li>Must have had an inadequate clinical response defined as not achieving sustained virologic response (SVR) with guideline-recommended preferred drugs in this UPDL category and indicated for diagnosis</li> </ul>

Infectious Disease Agents: Antivirals – Herpes		
PREFERRED AGENTS	NON-PREFERRED AGENTS	PA CRITERIA
acyclovir valacyclovir	famciclovir SITAVIG	<p><b><u>LENGTH OF AUTHORIZATIONS:</u></b> For the duration of the prescription (up to 180 days)</p> <p><b><u>NON-PREFERRED CRITERIA:</u></b></p> <ul style="list-style-type: none"> <li>• Must have had an inadequate clinical response of at least <u>3 days</u> with at least <u>one preferred</u> drug in this UPDL category and indicated for diagnosis</li> </ul>

Infectious Disease Agents: Antivirals – HIV* LEGACY CATEGORY		
PREFERRED AGENTS	NON-PREFERRED AGENTS	PA CRITERIA
INTEGRASE STRAND TRANSFER INHIBITORS		<b>LENGTH OF AUTHORIZATIONS:</b> 365 Days  <b>ABACAVIR/DOLUTEGRAVIR/LAMIVUDINE (TRIUMEQ PD) CRITERIA:</b> <ul style="list-style-type: none"><li>Must provide documentation of patient’s weight (only authorized for those 6 – 25 kg)</li></ul> <b>NON-PREFERRED CRITERIA:</b> <ul style="list-style-type: none"><li>Must have had an inadequate clinical response of at least <u>30 days</u> with at least <u>one preferred</u> drug in this UPDL category and indicated for diagnosis. If applicable, the request must address the inability to use the individual components.</li></ul> <b>AR</b> – ISENTRESS CHEWABLE TABLET: a PA is required for patients 12 years and older <b>AR</b> – lamivudine soln: a PA is required for patients 3 years and older <b>AR</b> – nevirapine soln: a PA is required for patients 3 years and older
APRETUDE ISENTRESS ISENTRESS CHEW TAB <sup>AR</sup> TIVICAY TIVICAY PD	VOCABRIA	
NUCLEOSIDE REVERSE TRANSCRIPTASE INHIBITORS		
abacavir emtricitabine entecavir lamivudine soln <sup>AR</sup> tenofovir dis fum 300mg VIREAD 150, 200mg TAB, POWDER zidovudine	abacavir soln EMTRIVA SOLN lamivudine tab VIREAD 250, 300mg TAB	
NON-NUCLEOSIDE REVERSE TRANSCRIPTASE INHIBITORS		
efavirenz nevirapine soln <sup>AR</sup> PIFELTRO	EDURANT etravirine nevirapine IR, ER tab	
PROTEASE INHIBITORS		
atazanavir darunavir 600, 800mg tab EVOTAZ PREZCOBIX REYATAZ POWDER ritonavir tab	APTIVUS fosamprenavir NORVIR POWDER PREZISTA SUSP, 75, 150mg TAB VIRACEPT	
OTHER SINGLE INGREDIENT PRODUCTS		
RUKOBIA	FUZEON SELZENTRY <sup>BvG</sup> SUNLENCA TYBOST	
COMBINATION PRODUCTS		

abacavir/lamivudine	CIMDUO	
BIKTARVY	lamivudine/zidovudine	
CABENUVA	STRIBILD	
COMPLERA	SYMFI <sup>BvG</sup>	
DELSTRIGO	SYMFI LO <sup>BvG</sup>	
DESCOVY		
DOVATO		
efavirenz/emtricitabine/tenofovir		
emtricitabine/tenofovir dis fum		
GENVOYA		
JULUCA		
lopinavir/ritonavir		
ODEFSEY		
SYMTUZA		
TRIUMEQ		
TRIUMEQ PD <sup>PA</sup>		

## Ophthalmic Agents: Antibiotic and Antibiotic-Steroid Combination Drops and Ointments

PREFERRED AGENTS	NON-PREFERRED AGENTS	PA CRITERIA
bacitracin-polymyxin CILOXAN ciprofloxacin erythromycin gentamicin moxifloxacin neo/poly/bacitracin neo/poly/bacitracin/hydrocortisone neo/poly/dexamethasone neo/poly/gramicidin ofloxacin polymyxin/trimethoprim sulfacetamide sodium soln 10% sulfacetamide/prednisolone TOBRADEX OINT tobramycin tobramycin/dexameth 0.3/0.1% TOBREX OINT	AZASITE bacitracin BESIVANCE gatifloxacin moxifloxacin (gen of MOXEZA) neo/poly/hydrocortisone sulfacetamide sodium oint 10% TOBRADEX ST ZYLET	<p><b><u>LENGTH OF AUTHORIZATIONS:</u></b> 30 days</p> <p><b><u>NON-PREFERRED CRITERIA:</u></b></p> <ul style="list-style-type: none"> <li>Must have had an inadequate clinical response of at least <u>3 days</u> with at least <u>two preferred</u> drugs in this UPDL category and indicated for diagnosis</li> </ul> <p><b><u>ADDITIONAL INFORMATION</u></b></p> <ul style="list-style-type: none"> <li>Requests may be authorized if the patient is completing a course of therapy that was started in the hospital or other similar location or was started before Medicaid eligibility, only the remaining course will be authorized</li> </ul>

Ophthalmic Agents: Antihistamines & Mast Cell Stabilizers		
PREFERRED AGENTS	NON-PREFERRED AGENTS	PA CRITERIA
azelastine BEPREVE <sup>BvG</sup> cromolyn ketotifen olopatadine	alomide bepotastine epinastine ZERVIA TE	<p><b><u>LENGTH OF AUTHORIZATIONS:</u></b> 365 Days</p> <p><b><u>NON-PREFERRED CRITERIA:</u></b></p> <ul style="list-style-type: none"> <li>Must have had an inadequate clinical response of at least <u>7 days</u> with at least <u>two preferred</u> drugs in this UPDL category and indicated for diagnosis</li> </ul>

## Ophthalmic Agents: Dry Eye Treatments

PREFERRED AGENTS	NON-PREFERRED AGENTS	PA CRITERIA
RESTASIS TRAYS <sup>BvG</sup> XIIDRA <sup>ST</sup>	CEQUA cyclosporine EYSUVIS MIEBO RESTASIS MULTI-DOSE TYRVAYA VEVYE	<p><b><u>LENGTH OF AUTHORIZATIONS:</u></b> 14 days for EYSUVIS; 365 days for all other drugs</p> <p><b><u>STEP THERAPY CRITERIA:</u></b></p> <ul style="list-style-type: none"> <li>Must have had an inadequate clinical response of at least <u>14 days</u> with at least <u>one</u> preferred drug in this UPDL category in the previous 120 days</li> </ul> <p><b><u>NON-PREFERRED CRITERIA:</u></b></p> <ul style="list-style-type: none"> <li>Must have had an inadequate clinical response of at least <u>14 days</u> with at least <u>two preferred</u> drugs in this UPDL category and indicated for diagnosis</li> </ul>

Ophthalmic Agents: Glaucoma Agents		
PREFERRED AGENTS	NON-PREFERRED AGENTS	PA CRITERIA
ALPHA-2 AGONISTS		<b>LENGTH OF AUTHORIZATIONS:</b> 365 Days  <b>STEP THERAPY CRITERIA:</b> <ul style="list-style-type: none"><li>Must have had an inadequate clinical response of at least <u>30 days</u> with at least <u>one preferred</u> drug in the same sub-section classification in this UPDL category and indicated for diagnosis, if available</li></ul> <b>NON-PREFERRED CRITERIA:</b> <ul style="list-style-type: none"><li>Must have had an inadequate clinical response of at least <u>30 days</u> with at least <u>two preferred</u> drugs within the same sub-section classification in this UPDL category and indicated for diagnosis</li></ul>
ALPHAGAN P 0.1% <sup>BvG</sup> ALPHAGAN P 0.15% <sup>BvG</sup> brimonidine 0.2%	apraclonidine brimonidine 0.1%, 0.15% IOPIDINE	
BETA BLOCKERS		
betaxolol carteolol levobunolol timolol gel, soln	BETIMOL 0.25% BETOPTIC S timolol hemihydrate soln 0.5% timolol maleate once daily, PF	
CARBONIC ANHYDRASE INHIBITORS		
AZOPT <sup>BvG ST</sup> dorzolamide	brinzolamide	
PROSTAGLANDINS		
latanoprost TRAVATAN Z <sup>BvG ST</sup>	bimatoprost IYUZEH LUMIGAN tafluprost travoprost VYZULTA XELPROS	
OTHER		
COMBIGAN <sup>BvG ST</sup> dorzolamide/timolol RHOPRESSA ROCKLATAN SIMBRINZA	brimonidine/timolol	



### Ophthalmic Agents: NSAIDs

PREFERRED AGENTS	NON-PREFERRED AGENTS	PA CRITERIA
diclofenac flurbiprofen ketorolac NEVANAC	ACUVAIL bromfenac ILEVRO	<p><b><u>LENGTH OF AUTHORIZATIONS:</u></b> 30 days</p> <p><b><u>NON-PREFERRED CRITERIA:</u></b></p> <ul style="list-style-type: none"> <li>Must have had an inadequate clinical response of at least <u>3 days</u> with at least <u>two preferred</u> drugs in this UPDL category and indicated for diagnosis</li> </ul>

Ophthalmic Agents: Ophthalmic Steroids		
PREFERRED AGENTS	NON-PREFERRED AGENTS	PA CRITERIA
ALREX <sup>BvG</sup> dexamethasone sodium phosphate difluprednate DUREZOL FLAREX fluorometholone FML FORTE LOTEMAX <sup>BvG</sup> MAXIDEX PRED FORTE PRED MILD prednisolone acetate prednisolone sodium phosphate	INVELTYS LOTEMAX SM loteprednol	<b><u>LENGTH OF AUTHORIZATIONS:</u></b> 30 days  <b><u>NON-PREFERRED CRITERIA:</u></b> <ul style="list-style-type: none"> <li>Must have had an inadequate clinical response of at least <u>7 days</u> with at least <u>two preferred</u> drugs in this UPDL category and indicated for diagnosis</li> </ul>

Otic Agents: Antibacterial and Antibacterial/Steroid Combinations		
PREFERRED AGENTS	NON-PREFERRED AGENTS	PA CRITERIA
CIPRO HC ciprofloxacin/dexamethasone CORTISPORIN-TC neomycin/poly B/hydrocortisone ofloxacin	ciprofloxacin ciprofloxacin/fluocinolone	<b>LENGTH OF AUTHORIZATIONS:</b> 30 days  <b>NON-PREFERRED CRITERIA:</b> <ul style="list-style-type: none"> <li>Must have had an inadequate clinical response of at least <u>3 days</u> with at least <u>two preferred</u> drugs in this UPDL category and indicated for diagnosis</li> </ul>

Respiratory Agents: Antihistamines – Second Generation		
PREFERRED AGENTS	NON-PREFERRED AGENTS	PA CRITERIA
cetirizine cap, syr, tab cetirizine/pseudoephedrine desloratadine fexofenadine levocetirizine loratadine rapid dissolve loratadine syr, tab loratadine/pseudoephedrine	cetirizine chewable <sup>AR</sup> CLARINEX-D loratadine chewable <sup>AR</sup> fexofenadine/pseudoephedrine	<b>LENGTH OF AUTHORIZATIONS:</b> 365 Days  <b>NON-PREFERRED CRITERIA:</b> <ul style="list-style-type: none"> <li>Must have had an inadequate clinical response of at least <u>7 days</u> with at least <u>two preferred</u> drugs in this UPDL category and indicated for diagnosis</li> </ul> <b>AR</b> – cetirizine chewable, loratadine chewable: a PA is required for patients 6 years and older

## Respiratory Agents: Cystic Fibrosis

PREFERRED AGENTS	NON-PREFERRED AGENTS	PA CRITERIA
ALYFTREK <sup>PA</sup> KALYDECO <sup>PA</sup> ORKAMBI <sup>PA</sup> PULMOZYME <sup>PA</sup> SYMDEKO <sup>PA</sup> TRIKAFTA <sup>PA</sup> PAK <sup>AR</sup> , TAB	BRONCHITOL	<p><b><u>LENGTH OF AUTHORIZATIONS:</u></b> Initial: 90 days; Subsequent: 365 days</p> <p><b><u>CLINICAL PA CRITERIA:</u></b></p> <ul style="list-style-type: none"> <li>• Must be prescribed by or in consultation with a pulmonologist or infectious disease specialist</li> <li>• Must provide documentation of the specific Cystic Fibrosis Transmembrane Conductance Regular (CFTR) genetic mutation</li> </ul> <p><b><u>NON-PREFERRED CRITERIA:</u></b></p> <ul style="list-style-type: none"> <li>• Must have had an inadequate clinical response of at least <u>30 days</u> with at least <u>one preferred</u> drug in this UPDL category and indicated for diagnosis</li> </ul> <p><b><u>ADDITIONAL BRONCHITOL CRITERIA:</u></b></p> <ul style="list-style-type: none"> <li>• Must be used as an add-on maintenance therapy</li> <li>• Must provide documentation of a completed BRONCHITOL Tolerance Test</li> </ul> <p><b>AR – TRIKAFTA PAK:</b> a PA is required for patients 6 years and older</p>

## Respiratory Agents: Epinephrine

PREFERRED AGENTS	NON-PREFERRED AGENTS	PA CRITERIA
epinephrine (labeler 49502) EPIPEN EPIPEN JR	AUVI-Q epinephrine (labeler 00093, 00115) NEFFY	<p><b><u>LENGTH OF AUTHORIZATIONS:</u></b> 365 Days</p> <p><b><u>NON-PREFERRED CRITERIA:</u></b></p> <ul style="list-style-type: none"> <li>• Must have had an inadequate clinical response to at least <u>one preferred</u> drug in this UPDL category and indicated for diagnosis</li> </ul> <p><b><u>SUBSEQUENT AUTHORIZATION CRITERIA:</u></b></p> <ul style="list-style-type: none"> <li>• Subsequent reauthorizations for expired epinephrine auto-injectors are allowable</li> </ul>

Respiratory Agents: Hereditary Angioedema		
PREFERRED AGENTS	NON-PREFERRED AGENTS	PA CRITERIA
ACUTE		<b>LENGTH OF AUTHORIZATIONS:</b> Acute: 30 days; Prophylaxis: 180 Days  <b>CLINICAL PA CRITERIA:</b> <ul style="list-style-type: none"><li>Acute Treatment<ul style="list-style-type: none"><li>Must provide documentation that diagnosis is verified by a C4 level below the lower limit of normal as defined by laboratory testing AND one of the following:<ul style="list-style-type: none"><li>C1 inhibitor (C1-INH) antigenic level below the lower limit of normal as defined by laboratory testing; OR</li><li>C1-INH functional level below the lower limit of normal as defined by laboratory testing</li></ul></li></ul></li><li>Prophylactic Treatment<ul style="list-style-type: none"><li>Must provide documentation that diagnosis is verified by a C4 level below the lower limit of normal as defined by laboratory testing AND one of the following:<ul style="list-style-type: none"><li>C1 inhibitor (C1-INH) antigenic level below the lower limit of normal as defined by laboratory testing; OR</li><li>C1-INH functional level below the lower limit of normal as defined by laboratory testing; OR</li><li>Presence of a known HAE-causing C1-INH mutation</li></ul></li></ul></li><li>All indications<ul style="list-style-type: none"><li>History of moderate or severe attacks such as airway swelling, severe abdominal pain, facial swelling, nausea and vomiting, or painful facial distortion</li></ul></li></ul> <b>NON-PREFERRED CRITERIA:</b> <ul style="list-style-type: none"><li>Must have had an inadequate clinical response of at least <u>3 days</u> with at least <u>one preferred</u> acute drug in this UPDL category and indicated for diagnosis to request a non-preferred acute drug.</li><li>Must have had an inadequate clinical response of at least <u>14 days</u> with at least <u>one preferred</u> prophylaxis drug in this UPDL category and indicated for diagnosis to request a non-preferred prophylaxis drug.</li></ul>
BERINERT <sup>PA</sup> icatibant acetate <sup>PA</sup>	KALBITOR RUCONEST	
PROPHYLAXIS		
TAKHZYRO <sup>PA</sup>	CINRYZE HAEGARDA ORLADEYO	

## Respiratory Agents: Inhaled Agents

PREFERRED AGENTS	NON-PREFERRED AGENTS	PA CRITERIA
ANTICHOLINERGIC BRONCHODILATORS/COMBINATIONS		LENGTH OF AUTHORIZATIONS: 365 Days  NON-PREFERRED CRITERIA: <ul style="list-style-type: none"><li>Must have had an inadequate clinical response of at least <u>14 days</u> with at least <u>two preferred</u> drugs in this UPDL category within the same sub-section classification and indicated for diagnosis</li></ul> ADDITIONAL STEROID-CONTAINING INHALER CRITERIA: <ul style="list-style-type: none"><li>Must have had an inadequate clinical response of at least <u>14 days</u> with at least one preferred steroid-containing drug</li></ul> ADDITIONAL BUDESONIDE/ALBUTEROL (AIRSUPRA) CRITERIA: <ul style="list-style-type: none"><li>Must have had an inadequate clinical response of at least <u>14 days</u> with either DULERA or SYMBICORT</li></ul> AR – albuterol nebulizer soln 0.021% (0.63mg/3mL), 0.042% (1.25mg/3mL): a PA is required for patients 13 years and older AR – budesonide nebulizer soln: a PA is required for patients 13 years and older
ANORO ELLIPTA <sup>BvG</sup> ATROVENT HFA COMBIVENT RESPIMAT INCRUSE ELLIPTA ipratropium ipratropium/albuterol neb soln SPIRIVA <sup>BvG</sup> STIOLTO	BEVESPI AEROSPHERE DUAKLIR PRESSAIR tiotropium inhaled caps TUDORZA umeclidinium/vilanterol YUPELRI	
ADRENERGIC BRONCHODILATORS		
albuterol HFA albuterol neb 0.021% (0.63mg/3mL), 0.042% (1.25mg/3mL) <sup>AR</sup> albuterol neb 0.083% (2.5mg/3mL) albuterol neb 0.5% (5mg/mL) conc arformoterol neb SEREVENT DISKUS STRIVERDI RESPIMAT VENTOLIN HFA XOPENEX HFA <sup>BvG</sup>	formoterol fumarate levalbuterol PROAIR DIGIHALER, RESPICLICK	
BRONCHODILATOR/GLUCOCORTICOID COMBINATIONS		
ADVAIR DISKUS <sup>BvG</sup> ADVAIR HFA <sup>BvG</sup> DULERA SYMBICORT <sup>BvG</sup>	AIRDUO DIGIHALER AIRSUPRA BREO ELLIPTA <sup>BvG</sup> BREYNA BREZTRI AEROSPHERE budesonide/formoterol fluticasone/salmeterol TRELEGY ELLIPTA WIXELA INHUB	
GLUCOCORTICOIDS		



ARNUITY ELLIPTA	ALVESCO	
ASMANEX TWISTHALER	ARMONAIR DIGIHALER	
budesonide neb susp <sup>AR</sup>	ASMANEX HFA	
FLOVENT		
fluticasone propionate		
PULMICORT FLEXHALER		
QVAR		
OTHER		
cromolyn neb soln		

Respiratory Agents: Leukotriene Receptor Modifiers & Inhibitors		
PREFERRED AGENTS	NON-PREFERRED AGENTS	PA CRITERIA
montelukast zafirlukast <sup>ST</sup>	zileuton ER ZYFLO	<p><b><u>LENGTH OF AUTHORIZATIONS:</u></b> 365 Days</p> <p><b><u>STEP THERAPY CRITERIA:</u></b></p> <ul style="list-style-type: none"> <li>• Must have had an inadequate clinical response of at least <u>30 days</u> with at least <u>one preferred</u> drug in this UPDL category</li> </ul> <p><b><u>NON-PREFERRED CRITERIA:</u></b></p> <ul style="list-style-type: none"> <li>• Must have had an inadequate clinical response of at least <u>30 days</u> with at least <u>two preferred</u> drugs in this UPDL category and indicated for diagnosis</li> </ul>

## Respiratory Agents: Monoclonal Antibodies-Anti-IL/Anti-IgE

PREFERRED AGENTS	NON-PREFERRED AGENTS	PA CRITERIA
DUPIXENT <sup>PA</sup> FASENRA <sup>PA</sup> XOLAIR <sup>PA</sup>	CINQAIR NUCALA TEZSPIRE	<p><b><u>LENGTH OF AUTHORIZATIONS:</u></b> Initial: 180 days; Subsequent: 365 days</p> <p><b><u>CLINICAL PA CRITERIA:</u></b></p> <ul style="list-style-type: none"> <li>• Must be prescribed by or in consultation with an applicable specialist (i.e., allergist/ immunologist, pulmonologist, or otolaryngologist)</li> <li>• For <b>Asthma</b> – Must have had uncontrolled asthma symptoms and/or exacerbations despite at least <u>30 days</u> with:               <ul style="list-style-type: none"> <li>○ Medium dose preferred ICS/LABA inhaler for 6 years and older <b>OR</b> medium dose preferred ICS/LABA inhaler with tiotropium or high dose ICS/LABA inhaler if 12 years and older</li> </ul> </li> <li>• For <b>Chronic Rhinosinusitis with Nasal Polyposis</b> – Must have had an inadequate clinical response of at least <u>30 days</u> to at least <u>one oral</u> corticosteroid <b>AND</b> <u>one nasal</u> corticosteroid spray</li> <li>• For <b>Chronic Urticaria</b> – Must have had an inadequate clinical response to at least <u>14 days</u> with at least <u>two different</u> second-generation antihistamines at 4 times standard dose</li> <li>• For <b>Chronic Obstructive Pulmonary Disease (COPD)</b>:               <ul style="list-style-type: none"> <li>○ The patient must have an eosinophilic count of greater than or equal to 300 cells per mcL within 12 months prior to initiation of therapy <b>AND</b></li> <li>○ The patient has a history of uncontrolled disease, as indicated by greater than or equal to 2 COPD exacerbations or greater than or equal to 1 COPD exacerbation resulting in a hospitalization despite being on standard of care, defined as triple therapy (LAMA+LABA+ICS) for at least 3 months prior to request, and at a stable dose for at least 1 month prior.</li> </ul> </li> </ul> <p><b><u>NON-PREFERRED CRITERIA:</u></b></p> <ul style="list-style-type: none"> <li>• Must have had an inadequate clinical response of at least <u>90 days</u> with at least <u>one preferred</u> drug in this UPDL category and indicated for diagnosis</li> </ul> <p><b><u>SUBSEQUENT AUTHORIZATION CRITERIA:</u></b></p> <ul style="list-style-type: none"> <li>• Must provide documentation of patient's clinical response to treatment and ongoing safety monitoring (i.e., PFT improvement, reduced affected BSA)</li> </ul>

Respiratory Agents: Nasal Preparations		
PREFERRED AGENTS	NON-PREFERRED AGENTS	PA CRITERIA
GLUCOCORTICOID/COMBINATIONS		LENGTH OF AUTHORIZATIONS: 365 days  NON-PREFERRED CRITERIA: <ul style="list-style-type: none"><li>Must have had an inadequate clinical response of at least <u>14 days</u> with at least <u>two preferred</u> drugs in this UPDL category within the same sub-section classification and indicated for diagnosis</li></ul>
flunisolide fluticasone (gen of FLONASE)	azelastine/fluticasone spray BECONASE AQ mometasone OMNARIS QNASL RYALTRIS XHANCE ZETONNA	
OTHER		
azelastine ipratropium olopatadine		

Respiratory Agents: Pulmonary Fibrosis		
PREFERRED AGENTS	NON-PREFERRED AGENTS	PA CRITERIA
OFEV <sup>PA</sup>	pirfenidone	<p><b><u>LENGTH OF AUTHORIZATIONS:</u></b> 365 Days</p> <p><b><u>CLINICAL PA CRITERIA:</u></b></p> <ul style="list-style-type: none"> <li>• Must be prescribed by or in consultation with a pulmonologist</li> </ul> <p><b><u>NON-PREFERRED CRITERIA:</u></b></p> <ul style="list-style-type: none"> <li>• Must have had an inadequate clinical response of at least <u>30 days</u> with at least <u>one preferred</u> drug in this UPDL category and indicated for diagnosis</li> </ul>

### Topical Agents: Antifungals

PREFERRED AGENTS	NON-PREFERRED AGENTS	PA CRITERIA
<p>ALEVAZOL</p> <p>butenafine</p> <p>ciclopirox</p> <p>clotrimazole</p> <p>clotrimazole/betamethasone</p> <p>econazole</p> <p>ketoconazole</p> <p>miconazole</p> <p>nystatin</p> <p>nystatin/triamcinolone</p> <p>terbinafine</p> <p>tolnaftate cream, powder</p>	<p>ciclopirox kit</p> <p>JUBLIA</p> <p>ketoconazole foam</p> <p>luliconazole</p> <p>miconazole/zinc/white petrolatum oint</p> <p>naftifine</p> <p>oxiconazole</p> <p>OXISTAT</p> <p>tavaborole</p> <p>tolnaftate soln</p>	<p><b><u>LENGTH OF AUTHORIZATIONS:</u></b> 365 days</p> <p><b><u>NON-PREFERRED CRITERIA:</u></b></p> <ul style="list-style-type: none"> <li>• Must have had an inadequate clinical response of at least <u>14 days</u> with at least <u>two preferred</u> drugs in this UPDL category and indicated for diagnosis</li> </ul> <p><b><u>ADDITIONAL EFINACONAZOLE (JUBLIA) CRITERIA:</u></b></p> <ul style="list-style-type: none"> <li>• Must have had an inadequate clinical response of at least 48 weeks of ciclopirox AND 6 weeks of oral terbinafine (if fingernail) <b>OR</b> 12 weeks of oral terbinafine (if toenail)</li> </ul>

### Topical Agents: Antiparasitics

PREFERRED AGENTS	NON-PREFERRED AGENTS	PA CRITERIA
NATROBA <sup>BvG</sup> permethrin piperonyl butoxide/pyrethrins VANALICE	CROTAN ivermectin lot malathion spinosad	<p><b><u>LENGTH OF AUTHORIZATIONS:</u></b> 14 Days</p> <p><b><u>NON-PREFERRED CRITERIA:</u></b></p> <ul style="list-style-type: none"> <li>Must have had an inadequate clinical response of at least <u>14 days</u> with at least <u>one preferred</u> drug in this UPDL category and indicated for diagnosis</li> </ul>

## Topical Agents: Corticosteroids

PREFERRED AGENTS	NON-PREFERRED AGENTS	PA CRITERIA
LOW POTENCY		<b>LENGTH OF AUTHORIZATIONS:</b> 365 days  <b>NON-PREFERRED CRITERIA:</b> <ul style="list-style-type: none"><li>Must have had an inadequate clinical response of at least <u>14 days</u> with at least <u>two preferred</u> drugs within the same sub-section classification in this UPDL category and indicated for diagnosis, if available</li></ul>
desonide cream, oint fluocinolone acetone 0.01% hydrocortisone	alclometasone desonide lotion TEXACORT	
MEDIUM POTENCY		
betamethasone valerate flurandrenolide fluticasone propionate cream, oint prednicarbate triamcinolone cream, lotion, oint	betamethasone val aerosol foam clocortolone pivalate fluocinolone acetone 0.025% fluticasone propionate lotion hydrocortisone butyrate, valerate PANDEL triamcinolone spray	
HIGH POTENCY		
betamethasone dip/calciptotriene oint fluocinonide 0.05% mometasone furoate	betamethasone dip betamethasone dip/calciptotriene susp desoximetasone diflorasone diacetate ENSTILAR halcinonide	
ULTRA HIGH POTENCY		
clobetasol propionate	APEXICON E BRYHALI fluocinonide 0.1% halobetasol propionate ULTRAVATE	



### Topical Agents: Immunomodulators

PREFERRED AGENTS	NON-PREFERRED AGENTS	PA CRITERIA
ELIDEL <sup>AR</sup> pimecrolimus <sup>AR</sup> [labeler 68682] tacrolimus <sup>AR</sup>	EUCRISA HYFTOR OPZELURA pimecrolimus <sup>AR</sup> [labeler 00591, 68462] VTAMA ZORYVE CREAM, FOAM	<p><b><u>LENGTH OF AUTHORIZATIONS:</u></b> 365 Days</p> <p><b><u>NON-PREFERRED CRITERIA:</u></b></p> <ul style="list-style-type: none"> <li>• Must have had an inadequate clinical response of at least <u>30 days</u> with at least <u>one preferred</u> drug in this UPDL category and indicated for diagnosis</li> </ul> <p><b><u>ADDITIONAL ROFLUMILAST (ZORYVE) CRITERIA:</u></b></p> <ul style="list-style-type: none"> <li>• <b><u>0.15% CREAM:</u></b> Must have had an inadequate clinical response of at least <u>30 days</u> with at least <u>one preferred</u> topical corticosteroid OR topical calcineurin inhibitor</li> <li>• <b><u>0.3% CREAM:</u></b> Must have had an inadequate clinical response of at least <u>30 days</u> with at least <u>one preferred</u> topical corticosteroid OR topical calcipotriene</li> <li>• <b><u>FOAM:</u></b> Must have had an inadequate clinical response of at least <u>30 days</u> with at least <u>one preferred</u> agent indicated for Seborrheic Dermatitis (such as a topical antifungal, topical calcineurin inhibitor, or topical corticosteroid)</li> </ul> <p><b>AR</b> – ELIDEL, pimecrolimus, and tacrolimus: a PA is required for patients younger than 2 years old</p>