


GUIDEBOOK NO. 13

SAMPLING THE LAYER CAKE THAT ISN'T: THE STRATIGRAPHY AND PALEONTOLOGY OF THE TYPE-CINCINNATIAN

Richard Arnold Davis and Roger J. Cuffey, Editors



originally prepared for the 1992 Annual Meeting of the Geological Society of America

Columbus 1998



DIVISION OF GEOLOGICAL SURVEY
4383 FOUNTAIN SQUARE DRIVE
COLUMBUS, OHIO 43224-1362
(614) 265-6576
(614) 447-1918 (FAX)
e-mail: geo.survey@dnr.state.oh.us
World Wide Web: http://www.dnr.state.oh.us/odnr/geo_survey/

OHIO GEOLOGY ADVISORY COUNCIL

Dr. E. Scott Bair, *representing Hydrogeology*
Ms. F. Lynn Kantner, *representing At-Large Citizens*
Mr. David Wilder, *representing Coal*

Mr. Mark R. Rowland, *representing Environmental Geology*
Dr. Mark R. Boardman, *representing Higher Education*
Mr. William M. Rike, *representing Oil and Gas*

Mr. Ronald M. Tipton, *representing Industrial Minerals*

SCIENTIFIC AND TECHNICAL STAFF OF THE DIVISION OF GEOLOGICAL SURVEY

ADMINISTRATION (614) 265-6988

Thomas M. Berg, MS, *State Geologist and Division Chief*
Robert G. Van Horn, MS, *Assistant State Geologist and Deputy Division Chief*
Dennis N. Hull, MS, *Assistant State Geologist and Deputy Division Chief*
Michael C. Hansen, PhD, *Senior Geologist, Ohio Geology Editor, and Geohazards Officer*
Janet H. Kramer, *Fiscal Officer*
Billie Long, *Fiscal Specialist*
Sharon L. Stone, AD, *Executive Secretary*

BEDROCK GEOLOGY MAPPING GROUP (614) 265-6473

Edward Mac Swinford, MS, *Geologist Supervisor*
Glenn E. Larsen, MS, *Geologist*
Gregory A. Schumacher, MS, *Geologist*
Douglas L. Shrake, MS, *Geologist*
Ernie R. Slucher, MS, *Geologist*

CARTOGRAPHY & EDITING GROUP (614) 265-6593

Edward V. Kuehnle, BA, *Cartographer Supervisor*
Merrienne Hackathorn, MS, *Geologist and Editor*
Ray O. Klingbeil, AD, *Cartographer*
Robert L. Stewart, *Cartographer*
Lisa Van Doren, BA, *Cartographer*

ENVIRONMENTAL & SURFICIAL GEOLOGY GROUP (614) 265-6599

Richard R. Pavey, MS, *Geologist Supervisor*
C. Scott Brockman, MS, *Geologist*
Kim E. Vorbau, BS, *Geologist*

PETROLEUM GEOLOGY GROUP (614) 265-6598

Lawrence H. Wickstrom, MS, *Geologist Supervisor*
Mark T. Baranoski, MS, *Geologist*
Michael R. Lester, BS, *Data Systems Coordinator*
James McDonald, MS, *Geologist*
Ronald A. Riley, MS, *Geologist*
Joseph G. Wells, MS, *Database Administrator*

LAKE ERIE GEOLOGY GROUP (419) 626-4296, (419) 626-8767 (FAX)

Scudder D. Mackey, PhD, *Geologist Supervisor*
Danielle A. Foye, BS, *Geologist*
Jonathan A. Fuller, MS, *Geologist*
Donald E. Guy, Jr., MS, *Senior Geologist*
Diane E. Honoshofsky, *Secretary*
Dale L. Liebenthal, *Operations Officer & Research Vessel Operator*
Richard M. Weekley, AD, *Survey Technician*

INDUSTRIAL MINERALS GROUP (614) 265-6602

David A. Stith, MS, *Geologist Supervisor*
Ronald G. Rea, MS, *Geologist and Sample Repository Manager*

COAL GEOLOGY GROUP (614) 265-6594

Douglas L. Crowell, MS, *Geologist Supervisor*
Richard W. Carlton, PhD, *Senior Geologist*
Mark E. Wolfe, BS, *Geologist*

GEOLOGIC RECORDS CENTER (614) 265-6585

Garry E. Yates, NZCS, *Supervisor*
Angelena M. Bailey, *Administrative Assistant*
Madge R. Fitak, BS, *Office Assistant*

An Equal Opportunity Employer - M/F/H



recycled paper

STATE OF OHIO
George V. Voinovich, Governor
DEPARTMENT OF NATURAL RESOURCES
Donald C. Anderson, Director
DIVISION OF GEOLOGICAL SURVEY
Thomas M. Berg, Chief

GUIDEBOOK NO. 13

SAMPLING THE LAYER CAKE THAT ISN'T: THE STRATIGRAPHY AND PALEONTOLOGY OF THE TYPE-CINCINNATIAN

Richard Arnold Davis and Roger J. Cuffey, editors

Contributors:

Roger J. Cuffey
Department of Geosciences
Pennsylvania State University

Steven M. Holland
Geology Department
University of Georgia

Benjamin F. Dattilo
Geoscience Department
Weber State University (Ogden, Utah)

Brian Kirchner
Geology Department
Earlham College

Richard Arnold Davis
Biology Department
College of Mount St. Joseph (Cincinnati)

Lynn A. Rockwell
Department of Geosciences
Pennsylvania State University

Sharon C. St. Louis Diekmeyer
Department of Geology
University of Cincinnati

Gregory A. Schumacher
Division of Geological Survey
Ohio Department of Natural Resources

Lawrence I. Goldman
MINEsoft, Ltd.
Lakewood, Colorado

Douglas L. Shrake
Division of Geological Survey
Ohio Department of Natural Resources

Donald E. Hattin
Department of Geological Sciences
Indiana University

Richard A. Spohn
Geology/Physics Library
University of Cincinnati


Helen B. Hay (deceased)
formerly Geology Department
Earlham College

E. Mac Swinford
Division of Geological Survey
Ohio Department of Natural Resources

Stanley M. Totten
Department of Geology
Hanover College

Guidebook originally prepared for Field Trip 1 for the Annual Meeting of the Geological Society of America,
Cincinnati, Ohio, October 26-29, 1992

Columbus
1998



Typesetting, layout, and revision of some figures by Lisa Van Doren
Revision of some figures by Robert L. Stewart

Cover photo: E. O. Ulrich, in the type-Cincinnatian along the Ohio River below Covington, Kentucky. The photograph was taken in 1901 by R. S. Bassler (see *Geotimes*, v. 30, no. 9, p. 12, and v. 30, no. 12, p. 2); according to Ellis L. Yochelson, a print of this photograph, from Dr. Bassler, is in the archives of the Smithsonian Institution. Photo courtesy of Tom Gathright, Virginia Division of Mineral Resources, and of the Virginia Department of Environmental Sciences, Virginia Museum of Natural History at the University of Virginia, who own a lantern slide of the image.

CONTENTS

	Page
Preface	iv
1. An introduction to this volume	1
2. An introduction to the type-Cincinnatian	2

LOCALITIES AND UNITS

3. Kope to Bellevue Formations: the Riedlin Road/Mason Road site (Upper Ordovician, Cincinnati, Ohio, region)	10
4. Upper Kope (McMicken) collecting on the Miamitown/Hamilton-Cleves exit ramp (Upper Ordovician, southwesternmost Ohio).....	36
5. The Maysville bryozoan reef mounds in the Grant Lake Limestone (Upper Ordovician) of north-central Kentucky	38
6. The Madison/U.S. Route 421 road cuts—Bellevue through Upper Whitewater strata (Upper Ordovician, southeastern Indiana)	45
7. The Miamitown Shale: stratigraphic and historic context (Upper Ordovician, Cincinnati, Ohio, region)	49
8. The Brookville Dam spillway—Miamitown through Waynesville Formations (Upper Ordovician, southeastern Indiana).....	60
9. The Corryville Member of the Grant Lake Formation (Upper Ordovician, southwestern Ohio).....	64
10. The “Brookville formation” (“Excello”, Waynesville, and Liberty Members) at Bon Well Hill, near Brookville (Upper Ordovician, southeastern Indiana).....	79
11. Liberty, lower and upper Whitewater, and Saluda strata at Garr Hill/Brookville North (Upper Ordovician, southeastern Indiana)	84
12. “Excello” (Arnheim) to basal Saluda strata on Indiana Route 1 at South Gate Hill (Upper Ordovician, southeastern Indiana)	89
13. Upper Arnheim through lower Whitewater strata at Caesar Creek Lake (Upper Ordovician, southwestern Ohio)	95
14. Arnheim through upper Whitewater strata at the Madison-Indiana Route 56 road cuts (Upper Ordovician, southeastern Indiana)	100
15. The Whitewater Formation on U.S. Route 27 near Richmond (uppermost Ordovician, east-central Indiana)	106

INTERPRETATIONS

16. A new look at the Cincinnati Series from a mapping perspective.....	111
17. Paleogeography and paleoenvironments, Fairview through Whitewater Formations (Upper Ordovician, southeastern Indiana and southwestern Ohio).....	120
18. Sequence stratigraphy of the Cincinnati Series (Upper Ordovician, Cincinnati, Ohio, region).....	135

APPENDIXES

A. Type-Cincinnatian localities	152
B. Bibliography on the type-Cincinnatian	167

PREFACE

This guidebook was produced for use on a two-day field trip held in conjunction with the 1992 Annual Meeting of the Geological Society of America in Cincinnati. At that time, because of time constraints, only enough copies of the guidebook were produced to distribute to field-trip participants. The intention was to publish a typeset version for general distribution after the meeting. Unfortunately, the process took rather longer than originally intended. In the intervening years, progress has been made in the study of type-Cincinnati rocks. However, we have not endeavored to update the entire contents of the volume. Citations of publications that were “in press” in 1992, but subsequently have made it “in print,” have been modified. Typographical errors have been corrected, and some authors have chosen to make a few, mostly minor, alterations in their respective chapters. Many of the figures have been redone by the cartographic staff of the Division of Geological Survey. The bibliography has been augmented, but no comprehensive literature search for recent publications was undertaken.

In the years subsequent to the field trip, Helen Hay, one of the authors of several chapters in this volume, died. She was certainly a leading authority on the geology of the Cincinnati rocks on the west side of the Cincinnati Arch. She will be missed. We dedicate this volume to her memory.

1. AN INTRODUCTION TO THIS VOLUME

by
Richard Arnold Davis

“Oh, No! Not more on the Cincinnati!”—I hear your plaintive cry. “It’s been done.” “We solved it all years ago.” Well, maybe.

This volume is not a standard road-log guidebook. Instead, the localities are arranged by the stratigraphic units to be seen, beginning at the bottom of the type-Cincinnati and ending, strangely enough, at the top. Each locality paper is modeled somewhat after those in the Decade of North American Geology Centennial Field Guides of the Geological Society of America.

Partly because various localities expose various thicknesses of the column and various units, a given locality may be mentioned in more than one paper in the volume. In order to keep track of these localities and to avoid ambiguity in citing them in the various papers, each locality has a locality number. The format for these designations is OH-HA-0001. The first two letters refer to the state in which the locality lies, in this case, Ohio. The state designation is followed by a county designation, here Hamilton County. The concluding four-digit number is the locality within the county. Only for the principal locality or localities (fig. 1-1) for each locality paper are location details given in that paper. Otherwise, such information is relegated to the locality list (Appendix A) at the end of the volume.

In addition to the several “locality papers,” there are three of what might be termed interpretational papers. The first of these, by Gregory A. Schumacher, summarizes the perspectives and conclusions that have underlain and resulted from the recent work by the Ohio Division of Geological Survey in the Cincinnati rocks of the state. Then follows like coverage by Helen B. Hay, dealing especially with the Cincinnati rocks of eastern Indiana and western Ohio. Finally, there is a discussion by Steven M. Holland of the rocks of the type-Cincinnati from the perspective of sequence stratigraphy.

The volume closes with a lengthy bibliography of works on the type-Cincinnati (Appendix B). Each paper also has its own list of references cited. The appendicular bibliography does not include all the references cited in the various papers in the volume, but, rather, only those on the type-Cincinnati. There are, however, many references in the volume bibliography that are not cited in any of the papers in the volume.

Some readers may click their tongues and shake their heads about the sloppy use of the term “shale” in this volume. Although it is a deeply seated tradition to speak of the rocks of the type-Cincinnati as a sequence of interbedded limestones and shales, the local “shales” most generally lack the fissility that characterizes shale, *sensu stricto*. In short, the local “shales” probably should be referred to as claystones or mudstones. Nonetheless, the various authors in this volume generally have bowed to the hoary, shaly tradition.

Another minor frustration will result from the issue of formal stratigraphic units vs. informal stratigraphic units. In the literature as a whole, different authors have chosen to designate informal units in different ways. Some workers have done so by putting the initial letter of the unit part of the name in lower case, for example, Brookville formation (as opposed to the formal unit, the Kope Formation). Un-

fortunately, subsequent authors (and editors?) not always have been so careful as to keep that initial letter lower case. Hence, we have chosen also to put quotation marks around the names of informal units, for example “Brookville formation,” so that it will be obvious to the reader that there is something unusual about the unit name so flagged (and not just sloppy proofreading).

On the other hand, we have chosen not to have a mere proposed change in rank affect the formality of a unit name; for example, neither Corryville Formation nor Corryville Member displays the visible stigmata of informality.

Another, larger dose of potential frustration almost certainly will result from the fact that various authors in the volume have chosen to use different schemes of stratigraphic nomenclature. This, too, is a well-established tradition in the study of the rocks of the type-Cincinnati.

Would that this volume be a real step toward a greater understanding of the geological reality recorded in the rocks of the type-Cincinnati and toward the adoption of a single scheme of stratigraphic nomenclature that truly and accurately reflects that reality!!

Address of author: Richard Arnold Davis, Biology Department, College of Mount St. Joseph, 5701 Delhi Avenue, Cincinnati, OH 45233-1670.




FIGURE 1-1.—Map of the tri-state area showing major locations described in the locality papers (papers 3-15) in this guidebook. ★ = central Cincinnati area; BD = Brookville Dam (paper 8); BWH = Bon Well Hill (paper 10); CC = Caesar Creek (paper 13); GH = Garr Hill/Brookville North (paper 11); MC = Miamitown-Cleves (papers 4, 7); MD = Madison (papers 6, 14); MY = Maysville (paper 5); RM = Richmond (paper 15); RR = Riedlin Road/Mason Road (papers 3, 7); SGH = South Gate Hill (paper 12); SL = Stonelick Creek (paper 9).

2. AN INTRODUCTION TO THE TYPE-CININNATIAN

by
Roger J. Cuffey

Paleontologists recently have developed great interest in detailed studies of the tempo, mode, rate, and pattern of evolution. Such studies require abundant fossils and thick stratigraphic sections representing millions of years of uninterrupted sedimentation; the type-Cincinnati should provide many excellent opportunities. Comparable sequences elsewhere have permitted initial documentation (Cuffey, 1967, p. 37-39, 85-86) and later confirmation (Pachut and Cuffey, 1991; Pachut and others, 1991) of gradual evolution within trepostome bryozoans. Preliminary studies of certain type-Cincinnati trepostomes (Brown and Daly, 1985), by methods whose value has been re-emphasized (Cuffey and Pachut, 1990), have indicated similar phyletic trends and so hold much interest for future investigations.

Type-Cincinnati fossils (fig. 2-1), even though well known, have yielded less of this kind of information than might be expected. Personal experience with Cincinnati bryozoans suggests that one significant cause of this situation has been confusion generated by inadequacies and inconsistencies in the type-Cincinnati stratigraphic literature and usage. However, careful examination of that literature and appropriate field localities indicates that the



FIGURE 2-1.—Diorama reconstruction of shallow marine habitat characteristic of the Cincinnati area during the Late Ordovician. Courtesy of Cincinnati Museum of Natural History; the diorama was constructed about 1959 by noted museum artist Paul Marchand, under the direction of K. E. Caster.

differences in terminology and classification actually have much in common. In fact, a coherent, though complicated integration of many previous studies does provide a pragmatically useable framework (Errett and Cuffey, 1989) for investigating type-Cincinnati bryozoans; this integration involves paleoecologic interpretations (fig. 2-2), stratigraphic correlations (fig. 2-3), paleoenvironmental distributions (fig. 2-4), and derivative conclusions such as depth curves (fig. 2-5) and regional profiles (such as fig. 5-3 of Cuffey, paper 5 in this volume).

The type-Cincinnati consists of thin, interbedded limestones and shales that appear much alike at first glance but which exhibit subtle differences that have paleoecologic implications (fig. 17-2, Hay, paper 17 in this volume; Schumacher, paper 16 in this volume; and references in both). The fossiliferous limestone beds commonly have been interpreted as storm-lag (tempestite) deposits, although such lags obviously were modified by between-storm invertebrate encrustation and growth so as to produce complex and varied carbonate beds (Anstey and Fowler, 1969; Harris and Martin, 1979; Kidwell and Aigner, 1985; Kidwell, Fürsich, and Aigner, 1986). If deposition of the limestones was indeed storm related, higher percentages of limestone beds within a succession indicate more frequent penetration by storm waves and, hence, shallower water depths, whereas higher shale percentages imply deeper waters (Hay, paper 17 in this volume, and references therein). "Deeper," is only relative; the entire Cincinnati region was quite shallow throughout the Cincinnati Epoch and was characterized by gentle slopes, holes, and shoals, and depths ranging from as little as one or a very few meters (or feet) (Hatfield, 1968, p. 12-13, 20; Etnensohn and others, 1986, p. 210, 214; Cuffey, paper 5 in this volume) to on the order of 20, 25, 30, or perhaps 50 meters (60-150 ft) (Bucher, 1917, p. 288, 290; Anstey and Fowler, 1969, p. 674-675; Anstey and others, 1987, p. 166-167; Brown and Daly, 1987; Weir and others, 1984, p. 96-97).

Critical to evolutionary studies is clarification of type-Cincinnati stratigraphic relationships, including vertical successions, lateral facies, biochronologic correlations, and extent of disconformities. Moreover, the ranges of a great number of Cincinnati species are recorded in terms of the




FIGURE 2-2.—Diagrammatic summary of paleoecologic interpretations of the major type-Cincinnati facies groups; the stimuli of Hay and others (1981) and Weir and others (1984) are acknowledged.




FIGURE 2-3.—Correlation chart showing the type-Cincinnatian stratigraphic units that have been employed in recording bryozoan species ranges; the influences of Pojeta (1984), Sweet (1979), Hay and others (1981), and Caster, Dalvé, and Pope (1955) are acknowledged.




FIGURE 2-4.—Distribution of paleoenvironments within the type-Cincinnatian; patterns are those shown on figure 2-2; unit boundaries and unit names are abbreviated from figure 2-3.

diverse named stratigraphic units that have appeared on regional, local, and partial columns and charts over many years. These units can be coordinated consistently, as shown in figure 2-3, which can be compared and contrasted with the similar but nonduplicative charts in Hay (paper 17 in this volume), Holland (paper 18), and Schumacher (paper 16), as well as with earlier comprehensive portrayals (figs. 2-6 and 2-7).

Most of the type-Cincinnatian formations and members are characterized by one or another of the facies groups noted earlier (fig. 2-2). Thus, their paleoecologic implications can be plotted for the entire type-Cincinnatian (fig. 2-4), so that one easily can see the fluctuating mosaic of paleoenvironments

that produced this complicated set of deposits.

One can derive additional relevant information from figure 2-4. Moving horizontally across the chart permits construction of a regional profile or cross section of the adjacent paleoenvironments existing across the Cincinnatian region at a given moment in time, as for example at the time when the Maysville bryozoan reef mounds were growing (see Cuffey, paper 5 in this volume, fig. 5-3). Moving vertically up the chart enables construction of a depth curve for a given location within the region (fig. 2-5); it is interesting to note that the same major shallowing-upward cycles are evident in the papers by Hay, by Holland, and by Schumacher in this volume (papers 17, 18, and 16, respectively).




FIGURE 2-5.—Depth curve for the area between Cincinnati, Madison, and Richmond, inferred from the succession shown in figure 2-4 and using paleoenvironments appearing in figure 2-2.






FIGURE 2-6.—Stratigraphic units and fossil ranges for Edenian, Maysvillian, and Richmondian strata

Lamborg

SYSTEM	SERIES	STAGE	EXPLANATION:		SECTION	"MEMBER" and THICKNESS	"FORMATION"	
				= LIMESTONE				
ORDOVICIAN	CINCINNATIAN	MAYSVILLIAN			<i>Homotrypa</i> sp. <i>Platystrophia ponderosa auburnensis</i>	Mt. AUBURN 4.5m	McMILLAN	
					<i>Glyptocrinus dyeri</i> <i>Plectrothis</i> sp. <i>Rafinesquina nasuta</i> <i>Flexicalymene, Isotelus</i>	CORRYVILLE 13.5m (44')		
					<i>Platystrophia cypha</i> <i>Onniella</i> sp. "SHINGLED" <i>RAFINESQUINA</i>	BELLEVUE 8.5m (28')		
					<i>Glyptocrinus</i> sp. <i>Onniella</i> sp. <i>Rafinesquina</i> sp. <i>Ambonychia</i> sp. <i>Caritodens</i> sp. <i>Cyclonema inflatum</i> ? <i>Trigrammaria planoconvexa</i>	FAIRMOUNT "HILL QUARRY BEDS" 18m (60')		
					RECURRENCE OF <i>ONNIELLA</i> <i>Batostoma</i> sp. <i>Escharopora falciformis</i> <i>Platystrophia hopensis</i> <i>Cyclonema gracile</i>	Mt. HOPE 16m (53')		
			EDENIAN			<i>ONNIELLA</i> "ZONE" large bryozoa fauna <i>Plectrothis</i> sp. <i>Sinuities cancellatus</i>	McMICKEN 21m (69')	LATONIA
					RECURRENCE OF <i>TRIARTHURUS</i> <i>Homotrypa</i> sp. <i>Onniella emacerata</i> <i>Cyrtolites</i> sp. <i>Loxoplocus</i> sp. <i>Sinuities</i> sp. large pelecypod fauna <i>Flexicalymene granulosa</i> trilobite tracks <i>Climacograptus typicalis</i>	SOUTHGATE 37m (122')		
					<i>Onniella</i> sp. <i>Trigrammaria</i> sp. <i>Caritodens</i> sp. <i>Triarthrus</i> sp.	ECONOMY 16m (52')		
						FULTON BEDS		
					<i>Escharopora</i> sp. <i>Platystrophia</i> sp. <i>Cryptolithus tessellatus</i> <i>Onniella</i> sp. <i>Whiteavesia</i> sp. <i>Cyclonema</i> sp. <i>Triarthrus eatoni</i>	Pt. PLEASANT "RIVER QUARRY BEDS" 10.5m (34')		
	CHAMPLAINIAN				BROMLEY SHALE 4.5m exposed			

Lamborg

at and near Cincinnati (from Davis, 1992, p. 21-22; revised from Caster, Dalvé, and Pope, 1955, p. 12-13).




FIGURE 2-7.—Stratigraphic relationships of principal type-Cincinnatian lithostratigraphic units as determined from conodont biostratigraphy (from Pojeta, 1984, p. 80; modified from Sweet, 1979, p. 13).

REFERENCES CITED

- Anstey, R. L., and Fowler, M. L., 1969, Lithostratigraphy and depositional environment of the Eden Shale (Ordovician) in the tri-state area of Indiana, Kentucky, and Ohio: *Journal of Geology*, v. 77, p. 668-682.
- Anstey, R. L., Rabbio, S. F., and Tuckey, M. E., 1987, Bryozoan bathymetric gradients within a Late Ordovician epeiric sea: *Paleoceanography*, v. 2, no. 2, p. 165-176.
- Brown, G. D., Jr., and Daly, E. J., 1985, Analysis of evolution of structural characters of *Parvohallopora* Singh from the Dillsboro Formation, Cincinnati Series, Indiana, in Nielsen, Claus, and Larwood, G. P., eds., *Bryozoa: Ordovician to Recent: papers presented at the 6th International Conference on Bryozoa*, Vienna, 1983: Fredensborg, Denmark, Olsen and Olsen, p. 51-58.
- , 1987, Trepostome bryozoan distributions in the Dillsboro Formation (Cincinnati Series), Indiana: water depth controls?, in Ross, J. R. P., ed., *Bryozoa: present and past: Western Washington University*, p. 41-47.
- Bucher, W. H., 1917, Large current-ripples as indicators of paleogeography: *Proceedings of the U.S. National Academy of Sciences*, v. 3, p. 285-291.
- Caster, K. E., Dalvé, E. A., and Pope, J. K., 1955, Elementary guide to the fossils and strata of the Ordovician in the vicinity of Cincinnati, Ohio: Cincinnati Museum of Natural History, 47 p.
- Cuffey, R. J., 1967, Bryozoan *Tabulipora carbonaria* in Wreford Megacyclothem (Lower Permian) of Kansas: *University of Kansas Paleontological Contributions*, art. 43 (Bryozoa 1), 96 p.
- Cuffey, R. J., and Pachut, J. F., 1990, Clinal morphological variation along a depth gradient in the living scleractinian reef coral *Favia pallida*: effects on perceived evolutionary tempos in the fossil record: *Palaios*, v. 5, p. 580-588.
- Davis, R. A., ed., 1992, Cincinnati fossils: an elementary guide to the Ordovician rocks and fossils of the Cincinnati, Ohio, region: Cincinnati Museum of Natural History, 61 p.
- Errett, D. H., and Cuffey, R. J. 1989, Bryozoan mode of occurrence relative to individual beds in the type Cincinnati (Upper Ordovician; Ohio, Indiana, Kentucky) (abs.): *Geological Society of America Abstracts with Programs*, v. 21, no. 4, p. 11.
- Ettensohn, F. R., Amig, B. C., Pashin, J. C., Greb, S. F., Harris, M. Q., Black, J. C., Cantrell, D. J., Smith, C. A., McMahan, T. M., Axon, A. G., McHargue, G. J., 1986, Paleocology and paleoenvironments of the bryozoan-rich Sulphur Well Member, Lexington Limestone (Middle Ordovician), central Kentucky: *Southeastern Geology*, v. 26, no. 4, p. 199-219.
- Harris, F. W., and Martin, W. D., 1979, Benthic community development in limestone beds of the Waynesville (upper Dillsboro) Formation (Cincinnati Series, Upper Ordovician) of southeastern Indiana: *Journal of Sedimentary Petrology*, v. 49, p. 1295-1306.
- Hatfield, C. B., 1968, Stratigraphy and paleoecology of the Saluda Formation (Cincinnati) in Indiana, Ohio, and Kentucky: *Geological Society of America Special Paper* 95, 34 p.
- Hay, H. B., Pope, J. K., and Frey, R. C., 1981, Lithostratigraphy, cyclic sedimentation, and paleoecology of the Cincinnati Series in southwestern Ohio and southeastern Indiana, in Roberts, T. G., ed., *GSA Cincinnati '81 Field Trip Guidebooks: American Geological Institute*, v. 1, p. 73-86.
- Kidwell, S. M., and Aigner, T., 1985, Sedimentary dynamics of complex shell beds: Implications for ecologic and evolutionary patterns, in Bayer, U., and Seilacher, A., eds., *Sedimentary and evolutionary cycles: Berlin, Springer-Verlag*, p. 382-395.
- Kidwell, S. M., Fürsich, F. T., and Aigner, T., 1986, Conceptual framework for analysis and classification of fossil concentrations: *Palaios*, v. 1, p. 228-238.
- Pachut, J. F., and Cuffey, R. J., 1991, Clinal variation, intraspecific heterochrony, and microevolution in the Permian bryozoan *Tabulipora carbonaria*: *Lethaia*, v. 24, no. 2, p. 165-185.
- Pachut, J. F., Cuffey, R. J., and Anstey, R. L., 1991, The concepts of astogeny and ontogeny in stenolaemate bryozoans, and their illustration in colonies of *Tabulipora carbonaria* from the Lower Permian of Kansas: *Journal of Paleontology*, v. 65, p. 213-233.
- Pojeta, John, Jr., 1984, Biostratigraphy, in Weir, G. W., Peterson, W. L., and Swadley, W C, *Lithostratigraphy of Upper Ordovician strata exposed in Kentucky: U.S. Geological Survey Professional Paper* 1151-E, p. 77-82.
- Sweet, W. C., 1979, Conodonts and conodont biostratigraphy of post-Tyrone Ordovician rocks of the Cincinnati region: *U.S. Geological Survey Professional Paper* 1066-G, 26 p.
- Weir, G. W., Peterson, W. L., Swadley, W C, and Pojeta, J., Jr., 1984, Lithostratigraphy of Upper Ordovician strata exposed in Kentucky; with a section on biostratigraphy: *U.S. Geological Survey Professional Paper* 1151-E, 121 p.

Address of author: Roger J. Cuffey, Department of Geosciences (Deike Bldg.), Pennsylvania State University, University Park, PA 16802.

3. KOPE TO BELLEVUE FORMATIONS: THE RIEDLIN ROAD/MASON ROAD SITE (UPPER ORDOVICIAN, CINCINNATI, OHIO, REGION)

by
Sharon C. St. Louis Diekmeyer

SIGNIFICANCE

The Riedlin Road/Mason Road site (fig. 3-1) provides a well-exposed sequence of strata including the upper Kope, Fairview, and Bellevue Formations and at least two intervals of the Miamitown Shale facies (fig. 3-2). This road cut allows easy and safe access to all the formations present. Many recent studies have used this section for these reasons (Tobin, 1982; Jennette, 1986; Diekmeyer, 1990; Holland, 1990, 1993).




FIGURE 3-1.—Location of the Riedlin Road/Mason Road site (KY-KE-0001) (modified from Jennette, 1986).

LOCATION

This site (KY-KE-0001) is located in Kenton County, Kentucky, at the intersection of Kentucky Route 16 (Taylor Mill Road) and Riedlin Road/Mason Road, 0.4 mile (0.6 km) north of I-275 (Exit 79), at 39°01'40"N latitude, 84°30'38"W longitude on the Covington, Kentucky–Ohio, 7.5-minute quadrangle (Ford, 1974; Luft, 1971).

TYPE SECTIONS

The abundant exposures of fossiliferous strata in the Cincinnati area have provided opportunity for intense study since before the middle of the 19th century, when these rocks were called the “Blue Limestone” (Locke, 1838; Hall, 1842). The first designation of these beds as the Cincinnati Group (Meek and Worthen, 1865) referred to strata bounded below by the Trenton Limestone and above by the Silurian formations (Orton, 1873). This group was divided on the basis of stratigraphic position (Orton, 1873). These divisions included, in ascending order, the Point Pleasant Beds, the Cincinnati Beds, and the Lebanon Beds. The Cincinnati Beds, in turn, were subdivided stratigraphically into the River Quarry Beds, the Middle or Eden Shales, and the Hill Quarry Beds (on the top), on the basis of lithologic and faunal characteristics. Typical fossils from the Riedlin Road/Mason Road locality are illustrated in figures 3-16, 3-17, and 3-18 at the end of this paper.

KOPE FORMATION

The Eden Shales were named for 250 ft (76.2 meters) of blue clay and limestone interbeds (limestone-to-clay ratio 1:10; Orton, 1873) exposed in Eden Park in Cincinnati (fig. 3-3; this outcrop no longer is accessible because of development). Nickles (1902) used the term Utica Group for the Middle Shales on the basis of James Hall’s (1842) correlation with the Utica Shales of New York and divided these shales using lithologic and faunal variations. These beds were termed the “Lower Utica or *Aspidopora newberryi* Beds” (about 80 ft/24 meters of greenish-gray, drab, or yellowish shales, more indurated than those occurring stratigraphically higher), the “Middle Utica or *Batostoma jamesi* Beds” (about 120 ft/36 meters of rather unfossiliferous strata with less limestone than the other two units), and the “Upper Utica or *Dekayella ulrichi* Beds” (about 60 ft/18 meters of more fossil-rich shales with increased abundance of thin, slabby limestones). Bassler (1906), in turn, gave these members local geographic names—Economy, Southgate, and McMicken, respectively. Foerste (1905) named the lowermost dark-shale zone containing abundant specimens of the trilobite *Triarthrus eatoni* the “Fulton Beds.” Weiss and others (1965) concluded that all these divisions were based upon bryozoan zones found at Cincinnati and were not useful in subdividing the Eden Formation in areas adjacent to the Cincinnati area. On the basis of clastic ratio and bedding logs, Weiss and Sweet (1964) correlated Eden-like strata in the Maysville area of Ohio and Kentucky with the Eden (Edenian) and Mt. Hope (Maysvillian) Formations at Cincinnati. Because of confusion brought on by the usage of Eden for both a stage and a rock unit, especially in reference to a section where the rock unit transgressed the Edenian-Maysvillian boundary, the rock-unit name Kope Formation was proposed by Weiss and Sweet (1964) to include those strata falling between the Point Pleasant limestones and beds having a clastic ratio of 2.5 to 3.8 that they didn’t name but correlated with the Fairview Formation of the Cincinnati area. The name Kope was derived from Kope Hollow, in Brown County, Ohio, nearly 50 miles (75 km) to the southeast (fig. 3-4). There is no one complete section known. However, Weiss and Sweet (1964) established a composite section thickness of 73.5 meters (241 ft). Kope Hollow, Ohio (OH-BR-0003), Red Oak Creek, Ohio (OH-BR-0004), and Maysville, Kentucky (KY-MS-0004), were designated co-equal Kope type sections.

FAIRVIEW FORMATION

Orton (1873) named the Hill Quarry Beds for 125 ft (38.1 meters) of strata lying between the Eden Shales and the Lebanon Beds. These beds provided abundant limestone for building and for lime production. The Hill Quarry Beds make up the tops of the hills around the downtown Cincinnati area. Orton (1873) defined the contact with the underlying Eden Shales on the basis of the increase in the limestone-to-clay ratio (1:5 or 1:6). Winchell and Ulrich (1897) proposed the name Lorraine Group in their correlation of fauna from the




FIGURE 3-2.—Riedlin Road/Mason Road locality. Center photo (3) is a small-scale view of most of the section exposed at the locality; photographs 1, 2, 4, and 5 are keyed to it. 1, upper part of the Fairview Formation. 2, Miami Shale/Bellevue interval. 4, top three cycles in the upper part of the Kope Formation; the limestone bed indicated is just below the “two foot shale” that underlies the Kope-Fairview contact. 5, three cycles in the lower part of the Fairview Formation.




FIGURE 3-3.—Location of Eden Park, for which the Eden Shales were named. The arrow indicates the approximate location of the original type section of the Eden Shales. Newport, Kentucky–Ohio, 7.5-minute quadrangle.




FIGURE 3-4.—Location of Kope Hollow (OH-BR-0003), for which the Kope Formation was named. The position of the most obvious exposure is indicated by the arrow. Russellville, Ohio–Kentucky, 7.5-minute quadrangle.

Illinois area. Nickles (1902) divided the Lorraine Group into six intervals on the basis of faunal and lithologic character. These included, from bottom to top: the “Mt. Hope or *Amplexopora septosa* Beds,” the “Fairmount or *Dekayia aspera* Beds,” the “Bellevue or *Monticulipora molesta* Beds,” the “Corryville or *Chiloporella nicholsoni* Beds,” the “Mt. Auburn or *Platystrophia lynx* Beds,” and the “Warren or *Homotrypa bassleri* Beds.”

Mt. Hope beds

The Mt. Hope beds were thought to range between the “heavy, irregularly bedded limestones” containing abundant *Dalmanella multisecta* above the Utica Shale and the *Strophomena planoconvexa* zone (Nickles, 1902, p. 76). The type exposure was designated as the southeastern portion of Price Hill known as Mt. Hope (fig. 3-5).

Fairmount beds

The Fairmount beds consist of that portion of the Lorraine known as the Stone Quarries or Hill Quarry Beds of Orton (1873). Nickles (1902) proposed the name Fairmount, because all the hilltops in the Fairmount area (north of Price Hill and south or southwest of Cumminsville and west of Mill Creek) exhibited these beds (fig. 3-5). Nickles (1902) determined that these strata also occupied the hilltops around Covington and Newport, Kentucky. Bassler (1906) lumped the Mt. Hope and Fairmount beds into the Fairview Formation because of the difficulty in discerning faunal differences and mapping the units. Bassler (1906) named the Fairview for about 100 ft (30 meters) of strata at Fairview Heights, Cincinnati (OH-HA-0001). Ford (1965, 1967) redefined the Fairview Formation as a rock-stratigraphic unit and designated the south- and west-facing cliff below Bellevue Hill Park along Clifton Avenue (OH-HA-0003) as the type section (fig. 3-5). The boundaries of the Fairview were based on a mean clastic content of 64 percent and bedding-index characteristics (Ford, 1967).

BELLEVUE FORMATION

Nickles (1902) distinguished the “Bellevue or *Monticulipora molesta* Beds” as the 15 ft (4.6 meters) of strata exposed at the top of the cliff below the site of the old Bellevue House (OH-HA-0003). Lithologically, these beds exhibit a shelly appearance and contain abundant bryozoans. *Monticulipora molesta* was considered one of the most characteristic bryozoan species of these strata. At the top of the Bellevue, Nickles (1902) noted about 5 ft (1.5 meters) of section containing abundant specimens of what he called *Rafinesquina alternata*. Bassler (1906) chose to incorporate the Bellevue, Corryville, and Mt. Auburn beds of Nickles (1902) into the McMillan Formation, thus reducing them to member status. He felt these members were closely related and “not of sufficient status to be mapped separately” (Bassler, 1906, p. 10). He chose McMillan Avenue in the Clifton portion of Cincinnati as the type section for this formation. Peck (1966), however, considered the boundaries to be poorly defined, and he rejected the McMillan Formation and its three members as rock-stratigraphic units. Using lithologic characteristics, Peck (1966) assigned rocks in the Maysville, Kentucky, area that previously had been referred to the McMillan Formation to the Grant Lake Limestone. He



FIGURE 3-5.—Locations of several sites within the type-Cincinnati. Covington, Kentucky—Ohio, Cincinnati West, Ohio, and Newport, Kentucky—Ohio, 7.5-minute quadrangles. **BH** = Bellevue Hill (OH-HA-0003), **BK** = Bald Knob (OH-HA-0013), **CHS** = Rice and Gage Streets/Christ Hospital (OH-HA-0004), **EMG** = Emming Street (OH-HA-0002), **FM** = Fairmount section of Cincinnati, **FV** = Fairview Park (OH-HA-0001), **JH** = Jackson Hill (OH-HA-0018), **MH** = Mt. Hope, **MM** = McMillan Street (OH-HA-0027).

designated road cuts along Kentucky Route 1449 about 6.5 km (4.0 miles) north-northwest of Orangeburg, in Mason County, Kentucky (Sleepy Hollow, KY-MS-0005), as the type section. Previously, other workers had correlated the beds of the Grant Lake Limestone with the McMillan Formation in the Cincinnati area (Palmquist and Hall, 1960; McFarlan and Nosow, 1961; Carpenter and Ory, 1961). Ford (1967) formally defined the Bellevue Limestone as a rock unit based on lithologic character and designated the type section as the cliff at the intersection of Rice and Gage Streets, Cincinnati (OH-HA-0004; fig. 3-5). He suggested that the Bellevue Limestone may represent a northwesterly tongue of the Grant Lake Formation of Peck (1966). Tobin (1982) compared the Grant Lake Limestone and the Bellevue Limestone and found the rock types, fauna, bedding character, and shale content of the two to be similar.

MIAMITOWN SHALE

Many early workers noted the presence of a shaly interval containing abundant molluscan remains associated with the Bellevue Limestone. Ford (1967) proposed the name Miami town Shale to denote the thin shale unit that overlies the Fairview Formation. He noted that it is lithologically and faunally different from the surrounding strata and designated the highway cut along Interstate 74, 1.6 km (1 mile) west of Miami town as the type section (OH-HA-0008). Please see Dattilo (paper 7 in this volume) for details on the historical development of the Miami town Shale.

DESCRIPTIVE STRATIGRAPHY

Previous lithostratigraphic studies of the Cincinnati exhibit a diversity of opinions on the placement of boundaries and the criteria useful for separating units. Many authors stated that original unit boundaries were based solely on faunal zones (Cumings, 1922; Gutstadt, 1958; Brown and Lineback, 1966). Assessment of the original literature by others (Weiss, 1961; Krumpolz, 1980; Tobin, 1982, 1986) drew attention to the use of both lithologic and faunal character for distinguishing unit boundaries. Tobin (1982, 1986) revised the lithostratigraphy of the Cincinnati, basing his scheme on the original unit descriptions and the recognition of sequences of related facies within each unit. This approach opened the way for study of the variability of units on a wider geographic scale and reduced the parochialism of geographically isolated studies. Lithologic designations used in this paper follow the classification of Dunham (1962), as modified by Tobin (1982).

KOPE FORMATION

The Riedlin Road/Mason Road outcrop exposes the upper 8.5 meters (27.9 ft) of the Kope Formation. This sequence contains medium-bedded, generally unfossiliferous, blue-gray calcareous (locally silty) shales. These shales are interbedded with medium (3-4 cm) calcisiltites, thin, lenticular (1-2 cm) skeletal packstone/wackestone beds, and medium (less than 8 cm) to thick (more than 8 cm) planar skeletal packstone/grainstone beds. Many of the thick packstone/grainstone beds are laterally continuous and have been traced across the Cincinnati area (Jennette, 1986). The shale component makes up 75-80 percent of the unit's total thickness (shale interval thicknesses of

approximately 1 meter). There is a general increase in the number and thickness of the packstone/wackestone beds approaching the top of the formation. Sedimentary features noted in Kope strata include symmetrical and asymmetrical ripples, starved ripples, planar lamination and cross-lamination, tool marks, flute casts, gutter casts, load structures, and ball-and-pillow structures (Tobin, 1982). Ford (1967, p. 929) originally defined the Kope/Fairview contact as the base of the limestone bed above the "...uppermost terrigenous stratum greater than 2 feet thick." Tobin (1982) interpreted the terrigenous stratum to be the shale bed, and this interval has come to be termed the "two foot shale" that is readily observable throughout the Cincinnati area. Alternatively, Jennette (1986) placed the contact at the top of the limestone bed above this shale unit; this was done on the basis of a different depositional interpretation of the facies packages (refer to paleoenvironmental interpretation section).

FAIRVIEW FORMATION

Ford (1967) originally defined the Fairview Formation as interbedded limestones, siltstones, and shales with bedding thicknesses of generally less than 0.6 ft (0.2 meter). His emphasis on the "close spacing" of the limestone beds indicates the trend toward decreased shale occurring in thinner beds than those of the underlying Kope Formation (Tobin, 1982). At the Riedlin Road/Mason Road site the entire Fairview Formation is exposed. The 31.7 meters (104 ft) of section is approximately 50 percent planar to lenticular, gray to blue-gray, skeletal limestone beds. These limestones are generally packstones to grainstones with fewer wackestones (although wackestone lenses occur within and around thicker packstone/grainstone beds). Grainstones are more abundant than in the underlying Kope strata. Compound beds containing both a packstone or wackestone component and a grainstone component are common. Limestone beds commonly are blanketed with a thin (1-2 mm) to medium (1-2 cm) bioturbated silt layer. Calcareous, silty shales and calcareous siltstones constitute the remaining beds in this interval. The lower half of the Fairview Formation contains about 55 percent shale, whereas the upper half contains 45 percent (Tobin, 1982). Shales are generally less silty than those of the Kope beds.

The upper contact of the Fairview originally was defined by Ford (1967) as the top of the limestone bed that occurs below a shale unit that is more than 0.6 meter (2 ft) thick and exhibits a very different lithology (Miami town Shale). Imbricated (shingled) brachiopods occur in several beds toward the top of the Fairview Formation. Many workers have used the "shingled *Rafinesquina* zone" of Caster, Dalvé, and Pope (1955) as the boundary. However, there is much controversy regarding the regional extent of this "zone." The problem with this contact is the intertonguing nature of the Fairview, Miami town Shale, and the Bellevue beds (see Dattilo, paper 7 in this volume). Tobin (1982) redefined the top of the Fairview Formation as the top of the uppermost planar, medium-bedded limestones and shales. Either the shale-rich, nodular Miami town facies or the wavy-bedded, limestone-dominated Bellevue beds may occur above this contact. Designating a contact at the Riedlin Road/Mason Road site is difficult. However, the contact probably exhibits an intertonguing between the Fairview beds and a Miami town interval (see below).

MIAMITOWN SHALE

The Miamitown Shale contains shales interbedded with thin- to medium-bedded limestone lenses and beds and abundant limestone nodules (Tobin, 1982). This unit thickens north-northwestward from 0.4 meter (1.3 ft) in northern Kentucky to more than 5 meters (16.4 ft) at the type section (Miamitown West, OH-HA-0008). At Riedlin Road/Mason Road there appear to be two intervals of Miamitown Shale lithology. The first occurs above the Fairview Formation boundary, is about 0.5 meter (1.6 ft) thick, and has abundant siltstone layers. This corresponds temporally to the Miamitown Shale exposed at the type section (OH-HA-0008). The 0.4-meter (1.3-ft)-thick second interval is about 2.5 meters (8.2 ft) above the "upper shingled *Rafinesquina* zone." It contains an average of 76 percent shale, less silty than underlying shales (Tobin, 1982). The dominant limestone types are packstones, wackestones, and micstones (Tobin, 1982). Sedimentary structures include gutter casts and cross-lamination. The limestone nodules exhibit ovoid and ripple shapes, are not more than 15 cm (4 inches) in length, and commonly occur along discrete horizons. These nodules commonly are cross-laminated internally. The relationship of the Miamitown Shale facies to the Bellevue and Fairview Formations is discussed in depth by Dattilo (paper 7 in this volume).

BELLEVUE FORMATION

The Bellevue interval at Riedlin Road/Mason Road is characteristic of exposures in the Cincinnati area. It consists of irregularly bedded, gray, argillaceous limestones (70-90 percent, Tobin, 1982) composed of whole and broken brachiopod valves and bryozoan colonies. The rubbly or "crenulated" appearance of the beds results from the weathering of thin, shale partings and the convexity and concavity of the brachiopod valves (Ford, 1965, 1967). Many beds are not laterally traceable for more than a few meters, although some thicker beds may be followed along the length of the outcrop. Beds range from poorly sorted to well-sorted packstones and grainstones. Thin beds and partings of gray, calcareous shales contain more abundant fossil material than do underlying beds. Fining-upward and some coarsening-upward sequences are common within beds. Discrete horizons of fossils and fossil fragments within beds exhibit planar to indistinguishable orientation trends. The bases of beds commonly are composed of abundant concave-down brachiopod valves. Mud sheltering beneath valves and spar filling of brachiopods and bryozoan colonies are characteristic features. Mud within and surrounding valves and fragments may be pelletal. Increased size and robustness of faunal elements, dominated by brachiopods and bryozoans, characterize this interval. These fossils exhibit varying degrees of breakage, encrustation, boring, and corrosion. Encrusting bryozoans are much more abundant than in the Fairview or Kope Formations. Many bryozoan colonies are preserved as phosphatic internal molds of zoecia. There is also increased preservation of pelecypod internal molds and body fossils.

Because of the difficulty in finding exposures of the top of the Bellevue strata, Ford (1965, 1967) designated the contact as a gradational "zone" between the uppermost Bellevue "coquinite limestone" and the "overlying weak shale and thin-bedded limestone" (Ford, 1967, p. 934) of his "unnamed beds." Tobin (1982) determined that this contact

was not gradational but was the "sharp termination of thin, wavy-bedded, shale-poor strata characteristic of the Bellevue Limestone, and the beginning of the thicker, planar-bedded limestones and shales of the Corryville Formation" (Tobin, 1982, p. 166). Although not exposed at Riedlin Road/Mason Road, the Bellevue/Corryville contact in other outcrops is sharp to gradational (see Goldman, paper 8 in this volume).

PALEOENVIRONMENTAL INTERPRETATION

The local Cincinnati is characterized by alternating beds of relatively unfossiliferous shales and fossil-rich limestones. Many researchers have tried to explain the origin of this characteristic arrangement of strata. Bucher (1917) was the first worker to ascribe this pattern to the introduction of storm-derived shelly accumulations onto normal mud sedimentation. For many years, researchers proposed depositional models that depended on variations within the depositional system (autocyclic mechanisms). These models included variability in water temperature or rate of sediment supply (Fox, 1962). Weiss and others (1965) and Osborne (1971) suggested a model of shifting silt-dispersal systems that allowed blanketing of benthic communities on submarine highs, producing autochthonous, skeletal accumulations. Shifting sediments exposed other areas for colonization.

STORM-DOMINATED SYSTEMS

Over the past 25 years, storm events have been singled out as a predominant influence on Cincinnati depositional patterns. Anstey and Fowler (1969) interpreted the depositional pattern of the Eden Shale (now designated the Kope Formation) to be the result of an environment of quiet mud sedimentation interrupted by periodic storms that deposited winnowed, coarse shell lags. Harris and Martin (1979) suggested colonization of muddy substrates by flat-valved brachiopods as the initial stage in short-term faunal succession within the Cincinnati limestone beds. Increased stabilization of the substrate by brachiopods allowed colonization by trepostome bryozoans and, subsequently, by crinoids, trilobites, gastropods, and pelecypods. Storm-induced influx of terrigenous material smothered the community and provided a muddy substrate for inception of a new successional cycle. Examination of the limestone beds within the Kope Formation led Harrison and Mahan (*in Meyer and others*, 1981) to conclude that, commonly, only partial successional sequences were evident. The occurrences of sequences that originated after the pioneer stage indicate that postdepositional transport of shelly debris may have been responsible for second-generation successional communities. Modification of the substrate by accumulation of skeletal debris and the influence this has on community structure (taphonomic feedback) has been studied recently in relation to storm-dominated systems (Kidwell and Jablonski, 1983; Kidwell, 1986b; Kidwell and others, 1986). Tobin (1982) noted the arrangement of Kope through Bellevue strata into limestone-shale couplets. Resuspension of offshore muds by periodic storms and redeposition onto nearshore carbonates by waning storm conditions was the model used to explain these couplets.

PALEO GEOGRAPHIC INTERPRETATION

Recent studies of Late Ordovician paleogeography locate

the Cincinnati region between 15° and 25° south latitude and rotated 45° clockwise of present-day orientation (Kriese, 1981a; Tobin, 1982; Weir and others, 1984; Jennette, 1986; Jennette and Pryor, 1993). Many researchers have inferred a northward-dipping ramp during this time (Weiss and others, 1965; Hoffman, 1966; Ford, 1967; Anstey and Fowler, 1969; Hay, 1981; Weir and others, 1984). Reconstructed ancient oceanic- and wind-circulation patterns indicate a storm track moving toward the south-southwest and impinging on the ramp (Ziegler, 1981; Tobin, 1982) (fig. 3-6). Idealized storm sequences from the Middle and Upper Ordovician Martinsburg and Reedsville Formations on the eastern margin of the Appalachian Basin (Kriese, 1981a) are comparable to the patterns found in Cincinnati depositional sequences (Tobin, 1982).

STORM-DEPOSITIONAL MODEL

Many recent studies of storm-dominated systems on modern and ancient shelves use a proximal-distal model to explain the depositional patterns noted (Kriese, 1981a, 1981b; Brett, 1983; Aigner, 1985). This model has been used to explain the sequences found within the Kope-Fairview interval (Jennette, 1986; Jennette and Pryor, 1993). Hur-

ricanes or winter storms moved toward the south-southwest across the adjacent ocean and intersected the north-northwestward-sloping ramp perpendicularly or at a high angle to the shoreline. Offshore winds set up a gradient onshore that resulted in an associated offshore storm surge with enough velocity to entrain abundant terrigenous and bioclastic material on the ramp (fig. 3-7). The storm surge, moving beneath the onshore-trending currents, carried the coarser material offshore, depositing sorted skeletal material below fair-weather wave base. In turn, waning storm conditions carried finer grained material and deposited it farther down ramp (distal tempestites). Renewed sedimentation below fair-weather wave base resulted in mud accumulations blanketing shell lags and tempestites (fig. 3-7).

MEGACYCLES

Tobin and Pryor (*in* Meyer and others, 1981) and Tobin (1982) noted the arrangement of beds into repeating vertical sequences within the Kope-through-Bellevue interval. These megacycles are represented by a basal, limestone-rich hemicycle composed of grainstones, packstones, wackestones, and calcisiltites interbedded with shales and a capping, shale-rich hemicycle containing interbed-



FIGURE 3-6.—Paleogeographic reconstruction showing the location of the study area during the Late Ordovician, 438 million years ago. Arrows indicate the inferred paths of hurricanes and severe storms as they approached the Cincinnati ramp (modified from Scotese and Denham, 1988).




FIGURE 3-7.—Proximal-distal storm model showing the resultant lateral and vertical distribution of facies (from Aigner, 1985b).

ded thin carbonate beds (Meyer and others, 1981; Tobin, 1982). Changes in supply of terrigenous sediment resulting from delta switching, lateral changes in sediment supply, or tectonic and/or volcanic activity were suggested as possible explanations for these megacycles. Jennette (1986) and Jennette and Pryor (1993) re-interpreted these cycles as shallowing sequences, because the shale hemicycles appear to coarsen rather than fine upward, and the carbonate component exhibits thickening beds toward the top of each megacycle. The shale hemicycle represents offshore deposition that aggraded, shifting proximal deposition basinward. Thin carbonate beds within the shale component represent influxes of nearshore material during storm-surge activity. The capping, amalgamated grainstones (proximal tempestites) resulted from winnowing of previously deposited shell accumulations nearshore. Deposition of shales directly atop capping, amalgamated grainstones is thought to have resulted from a relatively rapid transgressive pulse related to glacio-eustatic sea-level rise (Jennette, 1986; Jennette and Pryor, 1993) (fig. 3-8). This asymmetrical regressive-transgressive pattern has been noted in other studies at a similar scale (Goodwin and Anderson, 1980; Aigner, 1982, 1985; Bayer and others, 1985; Einsele, 1985; Brett and others, 1986b).

LARGE-SCALE, SHOALING-UPWARD CYCLES

Three packages of thinning shale hemicycles and thickening limestone hemicycles were noted in the Cincinnati by Tobin and Pryor (*in* Meyer and others, 1981) and Tobin (1982); they interpreted these to be large-scale regressions. Holland (paper 18 in this volume) notes the presence of four shallowing-upward cycles. The Kope-through-Bellevue sequence is the first of these regressions. Location of individual megacycles within this large-scale shoaling sequence determines their character (fig. 3-9). Megacycles within the deeper water Kope Formation exhibit thick shale intervals (approximately 1 meter) that have thick amalgamated limestones restricted to the top of each cycle. Four megacycles can be delineated in the Kope Formation at the Riedlin Road/Mason Road site. Megacycles within the Fairview Formation are thinner (approximately 0.5 meter or less), and accompanying capping amalgamated limestone packages are thicker. Jennette (1986) and Jennette and Pryor (1993) documented four megacycles within the lower Fairview Formation. The upper Fairview Formation and the Bellevue Formation appear to represent deposition at or near fair-weather wave base where signatures of both the storm-induced cycles and the larger megacycles are overprinted by nearshore reworking.

FAUNAL ANALYSIS

Investigation of the faunal assemblages in the Kope-through-Bellevue sequence reveals a complex set of factors influencing their distribution. Environmental parameters such as food availability, substrate, depth, salinity, and energy (among others) affected their arrangement on the shelf. Additionally, interspecific and intraspecific interactions were factors in faunal distribution patterns. Post-mortem processes that modified the already-existing faunal assemblages were especially important in storm-dominated systems. Many researchers have documented a relationship between the storm and shoaling cycles within the Cincinnati and the changes in faunal assemblages (Anstey and




FIGURE 3-8.—Stratigraphic column showing megacycles interpreted to be shallowing sequences in the upper Kope and lower Fairview Formations (from Jennette, 1986). The scale is in meters.

Perry, 1972; Alexander, 1975; Brandt, 1980; Meyer and others, 1981, 1990; Schumacher, 1984; Jennette, 1986). Four major aspects of faunal analysis that provide insight into this relationship are abundance, diversity, paleoecol-




FIGURE 3-9.—Interpretation of the lateral relationship of the units in the Kope-through-Bellevue sequence (modified from Tobin, 1982).

ogy, and taphonomy. Faunal-abundance data from closely spaced samples of the Kope-through-Bellevue sequence at Riedlin Road/Mason Road, in association with previously established stratigraphic control (Tobin, 1982; Jennette, 1986), paleoecology, and taphonomic indicators, were used to ascertain environmentally meaningful patterns associated with storm events and transgressive/regressive sequences (Diekmeyer, 1990). The patterns resulting from this analysis are treated in the following sections.

Cluster analysis and polar ordinations were carried out on faunal-abundance data gathered from counts of whole fossils, grid counts, and percent-rock-volume approximations (shale samples) for fragmental faunal elements in 107 samples. Percent-rock-volume approximations for fossil fragments were ascertained by weighing shale samples, washing and sieving the samples, and weighing the individual taxa (bryozoans, crinoids, and trilobites) for each size fraction. The section was sampled at approximately 0.5-meter (1.6-ft) intervals from the base to the top of the outcrop. Cluster analysis delineated seven groups of samples distinguished on the basis of faunal content. Table 3-1 describes each cluster group based upon lithologic type and characteristics, stratigraphic location, and major faunal elements. The cluster groups, hereafter just called groups, are named for their most abundant faunal elements (fig. 3-10). They are:

- A. the *Onniella*-bryozoan group,
- B. the *Onniella*-crinoid-trilobite group,
- C. the *Rafinesquina*-bryozoan-crinoid group,
- D. the *Platystrophia*-bryozoan group,
- E. the *Zygospira*-bryozoan group,
- F. the *Plectrothis*-*Rafinesquina*-crinoid group, and
- G. the *Zygospira*-graptolite-trilobite group.

Although these groups are defined on fossil content, the beds within each group commonly have characteristic lithologies.

These groups were plotted onto the stratigraphic column to distinguish faunal transitions through the section (fig. 3-11). The distribution of these groups indicates a transition in faunal components from the bottom to the top of the section associated with the large-scale regression (Tobin, 1982).

FAUNAL PATTERNS ASSOCIATED WITH THE LARGE-SCALE REGRESSION

Faunal transitions from the upper Kope through the exposed Bellevue at Riedlin Road/Mason Road correspond to an increase in energy related to shallowing conditions. Successively shallowing environments were susceptible to more frequent reworking events that modified the substrate from a muddy bottom to shelly pavements. These shelly accumulations became increasingly amalgamated up section. In response, faunal assemblages range from organisms

associated with relatively deep, low-energy, soft-substrate conditions to those better suited to higher energy, firmer substrate environments.













In general, smaller-valved brachiopods (*Zygospira* and *Onniella*) are replaced by larger-valved, more robust specimens (*Rafinesquina*, *Platystrophia*, *Plectrothis*, and *Hebertella*) (fig. 3-12), although *Zygospira* is an important component in discrete intervals throughout the section. The trend toward increased size and shell strength suggests increasingly energetic conditions. Increased abundance of crinoid fragments in the upper Fairview (fig. 3-13) indicates increased shelly substrate resulting from shallowing conditions. The taphonomic character of crinoid fragments ranges from well-abraded, single columnals (transported into the muddy Kope environment) through articulated stem and arm fragments (rapid burial by storm events during the time in which the upper Fairview was deposited) to abundant stem and arm fragments of varying taphonomic state (in-place winnowing and reworking during deposition of the lower Bellevue). Assemblages of crinoids indicate a relationship between environmental conditions and crinoid diversity and morphology (Meyer and others, 1990). Crinoid fragments and, more rarely, individuals are represented primarily by a low-diversity fauna, including the small, slender inadunates *Ectenocrinus* and *Cincinnatiocrinus* in the Kope Formation. In the overlying Fairview, the more robust inadunate *Iocrinus* increases in abundance and is accompanied by abundant arms, stems, crowns, and individuals of the camerate *Pycnocrinus* (traditionally called *Glyptocrinus*). Stem fragments and cemented holdfasts of the large, robust inadunate *Anomalocrinus* are abundant in the Bellevue Formation (commonly associated with hardgrounds).






Morphologic variations of bryozoans also may indicate an overall shallowing trend. Assessment of general growth morphologies for each sample indicates a lack of encrusting forms in the Kope Formation, but abundance increases into the upper Fairview. Small, ramose bryozoans transcend the depth gradient, occurring in most samples. There appears to be a zone, approximately 8 meters (26 ft) thick, where massive, boxwork bryozoans occur. These colonies are mud filled, indicating burial in place, and their massive form suggests an environment turbulent enough to have maintained large numbers of filter-feeding individuals, but deep enough for them to have remained relatively undisturbed by storm events. Large, ramose bryozoans occur in the upper Kope (*Batostoma*, *Dekayia*, and *Parvohallopora* spp.), lower Fairview (*Escharopora*, *Constellaria*, *Parvohallopora* spp., *Heterotrypa*, and *Dekayia*), and upper Bellevue (*Parvohallopora ramosa*). Those in the Kope and Fairview deposits do not appear abraded and are surrounded by large amounts of mud, indicating smothering of the colonies by influxes of sediment. However, specimens from the Bellevue are abraded and lack a mud matrix, suggesting increased winnowing and possible transport.

Fragments of the trilobites *Flexicalymene* and *Isotelus* are most abundant in the Kope Formation, but have a fairly low abundance throughout the remaining section (fig. 3-14). This distribution may indicate increased disturbance frequency associated with a shallowing environment that inhibited the long-term establishment of a muddy bottom advantageous to trilobites.

Trace-fossil assemblages show a transition from a *Cruziana*-dominated ichnofacies through a mixed *Skolithos*-*Cruziana*-dominated ichnofacies associated with the large-

TABLE 3-1.—Faunal groups, showing their lithologies, sedimentologic features, stratigraphic locations, and major faunal elements

GROUP NAME	DISTINGUISHING CHARACTERISTICS	STRATIGRAPHIC LOCATION	MAJOR FAUNAL COMPONENTS	
GROUP A	AMAL GS SH 	THK PL 	KOPE CYCLE TOPS TRANSITIONAL ABOVE KOPE/FAIRVIEW	HIGH ONNIELLA HIGH BRYOZOAN INTERMEDIATE ZYGOSPIRA
GROUP B	SH SS PK WK PL BIOT		BETWEEN KOPE CYCLE TOPS	HIGH ONNIELLA INTERMEDIATE BRYOZOAN INTERMEDIATE ZYGOSPIRA INTERMEDIATE MODIOLOPSIS INTERMEDIATE TRILOBITE INTERMEDIATE CRINOID
GROUP C	PK PK/GR 	THN - THK UND 	UPPER FAIRVIEW	HIGH BRYOZOAN HIGH RAFINESQUINA INTERMEDIATE CRINOID
GROUP D	GS PK WK 	AUTOCH MED - THK UND 	LOWER FAIRVIEW TOPS OF FAIRVIEW CYCLES B & C TRANSITIONAL L - U FAIRVIEW UPPER BELLEVUE	HIGH BRYOZOAN INTERMEDIATE PLATYSTROPHIA INTERMEDIATE ZYGOSPIRA LOW - INTERMEDIATE RAFINESQUINA
GROUP E	GS PK SH 	 YY 	TRANSITIONAL KOPE/LOWER FAIRVIEW UPPER FAIRVIEW LOWER BELLEVUE FAIRVIEW TONGUE	HIGH BRYOZOAN HIGH ZYGOSPIRA LOW - INTERMEDIATE CRINOID
GROUP F	GS PK SH THN - THK 	YY  	LOWERMOST UPPER FAIRVIEW	HIGH PLECTORTHIS INTERMEDIATE - HIGH CRINOID INTERMEDIATE RAFINESQUINA
GROUP G	AUTOCH PK/WK SH SS		MIAMITOWN SHALE BETWEEN KOPE CYCLE TOPS	HIGH ZYGOSPIRA HIGH TRILOBITE HIGH GRAPTOLITE

KEY:	PL-PLANAR BEDS		FINING UPWARD SEQUENCE
AMAL- AMALGAMATED BEDS	THN -THIN BEDS		COARSENING UPWARD SEQUENCE
GS-GRAINSTONE	MED-MEDIUM BEDS		SHALE RIP-UP CLASTS
PK-PACKSTONE	THK-THICK BEDS		BRACHIOPODS
WK-WACKESTONE	BIOT-BIOTURBATION		BRYOZOANS
SH-SHALE	AUTOCH-AUTOCHTHONOUS FOSSIL ACCUMULATIONS		
SS-SILSTONE	BRACHIPOD PAVEMENTS		
UND-UNDULATING BEDS			

scale regression (Tobin, 1982). Abundant *Diplocraterion*, representative of the *Skolithos* facies, are present in the upper Kope Formation. Three separate ichnospecies occur in a calcareous siltstone near the top of the formation (Meyer and others, 1981). This bed is laterally traceable across the outcrop. Present-day organisms living in U-shaped burrows commonly occur in relatively shallow, well-oxygenated and well-lighted environments. Fürsich (1974, 1975) attributed *Diplocraterion* to a high-energy intertidal or very shallow subtidal environment. The abundance of *Diplocraterion* in an environment interpreted to have been at or below storm wave base requires further study.

FAUNAL PATTERNS WITHIN THE KOPE FORMATION

Megacycles in the Kope and lower Fairview distinguished by Tobin (1982), Jennette (1986), and Jennette and Pryor (1993) are represented by a repetitive succession of certain faunal groups. The *Onniella*-bryozoan group (A) is restricted to the Kope megacycle tops and lowermost Fairview. The *Onniella*-crinoid-trilobite group (B) is associated with the multiple-event packstones and wackestones between the Kope megacycle tops. The *Zygospira*-graptolite-trilobite group (G) occurs in the thin packstones/wackestones and shales between Kope megacycle tops (high trilobite




FIGURE 3-10.—The seven faunal groups showing the relative abundances of major faunal elements within each group. Percent abundances are indicated by the lengths of the bars for the faunal elements. The bar graph on the left shows the level of similarity among samples within each group and between groups. Similarity ranges from perfect correspondence, with a value of 1.0, to completely dissimilar, with a value of 0. Samples not falling within any group also are indicated.

abundance) and in the Miamitown Shale (high graptolite abundance). Polar ordination reveals a nearly linear distribution of groups A, B, and G along a gradient associated with *Zygospira* and *Onniella* relative abundances. The lowest relative abundance of *Onniella* and highest relative abundance of *Zygospira* are in the packstones, wackestones, and shales of group G. These strata represent smothering of distal communities by shale and silt during waning storm conditions or transport of bioclastic material into deeper muds by storm surge. As depth became increasingly shallow, reworking of carbonate materials produced the amalgamated packstones and wackestones of group B that contain an intermediate abundance of *Onniella*. The amalgamated grainstones (group A) contain the highest abundance of *Onniella* and represent the shallowest conditions in the Kope Formation. However, analysis of within-bed distribution of *Onniella* and *Zygospira* reveals increasing *Zygospira* from bed bottom to bed top. Morphologic assessment of these two brachiopods suggests that *Onniella*'s flat, more delicate valves were more suited to a muddier substrate where

they could "float" at the sediment-water interface. The common disarticulation and the abraded appearance of the valves may indicate that they were winnowed from the muds and reworked into the amalgamated beds by storm events. In contrast, the stronger articulation and greater density of the shells of *Zygospira* would have allowed animals of that genus to live on substrates associated with increased energy. Shells of *Zygospira* commonly are attached to bioclasts, including other *Zygospira* shells and crinoid stems (Meyer and others, 1981), suggesting that the animals had the ability to live in association with the shelly accumulations characteristic of shallowing conditions.

FAUNAL PATTERNS WITHIN THE FAIRVIEW FORMATION

Tobin (1982) interpreted the Fairview Formation to be the result of an environment intermediate between the deeper water Kope Formation and the shallower





FIGURE 3-11.—Stratigraphic column of the Riedlin Road/Mason Road site; lithologies are noted by patterns and widths of those in figure 3-10.



beds. Patterned blocks on the left of the column indicate the location of each sample and its faunal group. These patterns correspond to




FIGURE 3-12.—Distribution of five major brachiopod genera from the base (left) to the top (right) of the section, based on a five-point moving average of percent relative abundance.




FIGURE 3-13.—Percent relative abundance of crinoid fragments from the base (left) to the top (right) of the section.




FIGURE 3-14.—Percent relative abundance of trilobite fragments from the base (left) to the top (right) of the section.

water Bellevue Formation. On the basis of multivariate analyses of faunal-abundance data, several faunal patterns emerge (fig. 3-11). The *Zygospira*-bryozoan group (E) appears to be transitional between the Kope and lower Fairview and is intercalated with the upper Kope groups. The *Platystrophia*-bryozoan group (D) dominates the lower Fairview, stratigraphically replacing group E. The upper Fairview is characterized by alternating limestone beds containing groups C, D, E, and F, representing dominance by *Rafinesquina*, *Platystrophia*, *Zygospira*, and *Plectorthis*, respectively. *Zygospira*-dominated group E becomes increas-

ingly common approaching the Bellevue contact and may be related to the lower tongue of the Miamiotown Shale facies. Throughout the Fairview, the trend appears to be one of frequent disruption of community structure by events that concentrated and enhanced relative abundances of distinct faunal components. Following the final event represented in each amalgamated bed, there developed a new community that may or may not have contained the same dominant faunal elements as in the previous bed. As a result of the amalgamating nature of storm events, fine-scale faunal patterns (megacycles) are difficult to distinguish within

the Fairview Formation. However, a general pattern of increased size and robustness of brachiopod valves through the Fairview is evident and corresponds to decreased thicknesses of intervening shales associated with thinner, more abundant limestone beds.

FAUNAL PATTERNS OF THE BELLEVUE/ MIAMITOWN INTERVAL

The Bellevue interval represents the shallowest environment of the Kope-through-Bellevue large-scale regression of Tobin (1982). Researchers have interpreted the Miamitown Shale to be the result of one of the following: (1) a deepening event within the shallow Bellevue environment, (2) an increased clastic supply to the shelf, (3) variation in water chemistry, or (4) deposition in a lagoon (see Tobin, 1982, for summary). Continuing research points to the intimate relationship between the Miamitown and Bellevue facies based upon their interfingering relationship on a regional scale (see Dattilo, paper 7 in this volume). The overlap of faunal elements within these facies at the Riedlin Road/Mason Road site lends credence to this interpretation (fig. 3-11). The "Middle Shale Zone," "upper shingled *Rafinesquina* Zone," and the Miamitown Shale ("Gastropod Shale") intervals are characterized by the *Zygospira*-bryozoan group (E). Further, *Cyclonema* exhibits increased abundance in this interval, as do *Ambonychia* and *Caritodens*. Samples from the upper Miamitown facies fall into the *Zygospira*-graptolite-trilobite group (G). Samples in group G with high abundances of graptolites are found in this Miamitown interval, whereas samples with high abundances of trilobites are found in the Kope shales (fig. 3-10). The "Lower Massive Zone" of the Bellevue Formation contains a relatively high abundance of the large brachiopods *Platystrophia*, *Plectrothis*, *Hebertella*, and, less commonly, *Zygospira* and *Rafinesquina*, indicating shallowing conditions. Because of their large size and robustness, they were better suited for withstanding the higher energy associated with an environment at or near fair-weather wave base. Bryozoans are more abundant than in underlying beds and are represented by large and small ramose morphotypes and abundant encrusting and foliaceous types. Size sorting is common in the upper Bellevue; large brachiopod valves dominate the bottoms of beds, and *Zygospira* is abundant on the bed tops. Fossil fragments give additional evidence for shallowing conditions. Crinoid columnals in the lower Bellevue are associated with abundant arm plates, indicating possible winnowing but very little, if any, transport (Meyer and Meyer, 1986). The winnowing model for the Bellevue is substantiated by increased sorting of skeletal fragments, variable states of abrasion, and small average size of bioclasts.

Analyses of faunal-abundance data, taphonomy, and paleoecology, in association with increased understanding of facies relationships within storm-dominated systems, allow the reconstruction of an integrated depositional model. Figure 3-15 shows the distribution of faunal groups along the Cincinnati ramp for the Kope, Fairview, and Bellevue/Miamitown intervals at the Riedlin Road/Mason Road site. Gathering and analyzing faunal, stratigraphic, and lithologic data on a regional scale (Holland, 1990, 1993) provides a model for large-scale depositional patterns within Upper Ordovician strata. In addition, future research using faunal data from closely spaced lateral and vertical samples collected from Cincinnati and adjacent areas will provide the

basis for interpreting the emerging, smaller scale patterns and for "fine tuning" regional models.

ADDITIONAL LOCALITIES

For more information on localities, see Appendix A at the end of this volume.

"Lexington Limestone"

Sugar Creek (KY-GA-0001), contact with Kope

Point Pleasant

Bear Creek (OH-CT-0010) - 69 ft (21 meters); contact with overlying Kope
 Boat Run (OH-CT-0011) - 52 ft (15.8 meters); contact with overlying Kope
 Bradford (KY-BK-0001) - 63 ft (19.2 meters); base of Kope estimated by Weiss and others (1965) to be 15 ft (4.6 meters) higher than exposed section
 Chilo (OH-CT-0012) - 11 ft (3.4 meters)
 Foster (KY-BK-0003) - 60 ft (18.3 meters)
 Kope Hollow (OH-BR-0003) - 32 ft (9.8 meters)
 North Point Pleasant (OH-CT-0016) - 53 ft (16.2 meters)
 Slickaway Run (OH-CT-0017) - 4 ft (1.2 meters)
 Twelvemile Creek (OH-CT-0018) - 6.5 ft (2.0 meters), upper contact

Kope Formation

Arnold Creek (IN-O-0002)
 Aurora (IN-D-0001)
 Backbone Creek (OH-CT-0006) - 30 meters (98.4 ft)
 Bald Knob (OH-HA-0013) - 7.6 meters (24.9 ft)
 Batavia #1 (OH-CT-0001) - 16 meters (52.5 ft) of upper Kope
 Bear Creek (OH-CT-0010) - 70 ft (21.3 meters); contact with underlying Point Pleasant
 Bellevue Hill (OH-HA-0003) - 37 ft (11.3 meters)
 Boat Run (OH-CT-0011) - 72 ft (21.9 meters); contact with underlying Point Pleasant
 Briarly Creek (OH-HA-0014) - 19 ft (5.79 meters)
 Camp Claybanks (OH-HA-0015) - 12 ft (3.7 meters)
 Chilo (OH-CT-0012) - 30 ft (9.1 meters)
 Congress Run (OH-HA-0016)
 Covington (KY-KE-0004) - 30.5 meters (100.1 ft)
 Crosby Road (OH-HA-0017) - 32 ft (9.7 meters)
 Eightmile Creek (OH-HA-0034)
 Fort Thomas (KY-CP-0001) - 59 meters (193.6 ft)
 Foster (KY-BK-0003) - 10 ft (3.05 meters) + another 50 ft (15.2 meters) poorly exposed above that
 Galbraith Road (OH-HA-0032) - 2 ft (0.6 meter) to base of Fairview
 Galbraith Road, cliff section (OH-HA-0033) - 34 ft (10.4 meters)
 Georgetown (OH-BR-0006) - 8.9 meters (29.2 ft)
 Grand Avenue (OH-HA-0006) - 94 ft (28.7 meters) from quarry floor to Maysville
 Happy Hollow (OH-CT-0013)
 I-275/Turkeyfoot Road (KY-KE-0005) - 12.5 meters (41.0 ft)
 Indian Creek (OH-CT-0014)
 Jackson Hill Park (OH-HA-0018) - 6.1 meters (20 ft)
 Johns Hill (KY-CP-0005) - 8.7 meters (28.5 ft)
 Kope Hollow (OH-BR-0003) - 151 ft (46.0 meters); one of three co-equal type sections of Kope
 Maysville (KY-MS-0004) - one of three co-equal type




FIGURE 3-15.—Reconstructed distribution of five of the faunal groups along the Cincinnati ramp, showing the characteristic lithologies and fauna (based on Brett and others, 1986).

sections of Kope

- Miami Station (OH-HA-0019) - 53 ft (16.2 meters)
- Miamitown/Hamilton-Cleves exit (OH-HA-0009) - 19.9 meters (65.2 ft)
- Millvale (OH-HA-0035) - 106 ft (32.3 meters) in middle and upper parts
- Montana Avenue (OH-HA-0038) - 15.6 meters (51.2 ft)
- Muddy Creek (OH-HA-0020) - 106 ft (32.3 meters)
- Newport-Grand Avenue (KY-CP-0004) - 26 meters (85.3 ft)
- Newport Shopping Center (KY-CP-0002) - 164 ft (50.0 meters); upper contact
- Ninemile Creek (OH-CT-0015) - about 177 ft (54 meters), upper contact
- North Point Pleasant (OH-CT-0016) - 65 ft (19.8 meters)
- Orphanage Road (KY-KE-0003) - 30 meters (98.4 ft)

- Pabco/Timberview Apartments (OH-HA-0037) - 29.4 meters (96.5 ft), upper contact
- Rabbit Hash (KY-BE-0002)
- Rapid Run (OH-HA-0036) - 202 ft (61.57 meters), upper contact
- Red Oak Creek (OH-BR-0004) - one of three co-equal type sections of Kope
- Rising Sun (IN-O-0001) - 52 ft (15.8 meters), upper contact
- Riverside (OH-HA-0021) - 81 ft (24.7 meters), both contacts
- Sekitan Road (OH-HA-0022) - 62 ft (18.9 meters)
- Slickaway Run (OH-CT-0017) - 171 ft (52.1 meters)
- Southgate (KY-CP-0003) - 23.2 meters (76.1 ft)
- Sugar Creek (KY-GA-0001), contact with "Lexington"
- Tanners Creek (IN-D-0003) - 22 ft (6.7 meters), up-

per contact
 Twelvemile Creek (OH-CT-0018) - 108.5 ft (33.07 meters), lower contact
 West Fork (OH-HA-0024) - 110 ft (33.5 meters)
 West Harrison (IN-D-0004)
 White Swan Run (OH-BR-0010) - 161 ft (49.1 meters), to base of Maysvillian
 Wiley Branch (IN-SW-0001)

Grand Avenue Member of Kope Formation

Bald Knob (OH-HA-0013)
 Grand Avenue (OH-HA-0006) - type section
 Miami Station (OH-HA-0019) - 11 ft (3.4 meters)
 Miamitown/Hamilton-Cleves exit (OH-HA-0009)
 Muddy Creek (OH-HA-0020) - 12 ft (3.7 meters)
 Riverside (OH-HA-0021) - 11 ft (3.4 meters), both contacts
 Sekitan Road (OH-HA-0022)
 West Fork (OH-HA-0024)

Wesselman Tongue of Kope Formation

Briarly Creek (OH-HA-0014) - 5 ft (1.52 meters)
 Congress Run (OH-HA-0016)
 Miami Station (OH-HA-0019) - 27 ft (8.2 meters)
 Miamitown/Hamilton-Cleves exit (OH-HA-0009)
 Muddy Creek (OH-HA-0020) - 4 ft (1.2 meters)
 Riedlin Road/Mason Road (KY-KE-0001) - 8.5 meters (27.9 ft), upper contact
 Sekitan Road (OH-HA-0022) - 14 ft (4.3 meters)
 Wesselman (OH-HA-0011) - bounded above and below by Fairview; type section
 West Fork (OH-HA-0024) - 11 ft (3.4 meters)

Fairview Formation

Aiken School (OH-HA-0012) - 26 ft (7.9 meters)
 Aurora (IN-D-0001)
 Backbone Creek (OH-CT-0006) - 30 meters (98.4 ft)
 Bald Knob (OH-HA-0013) - 9.7 meters (31.8 ft)
 Batavia #2 (OH-CT-0002) - 12.1 meters (39.7 ft)
 Bellevue Hill (OH-HA-0003) - 108 ft (32.9 meters)
 Briarly Creek (OH-HA-0014)
 Brookville Dam spillway (IN-FR-0002) - in highwall
 Camp Claybanks (OH-HA-0015) - 84 ft (25.6 meters)
 Congress Run (OH-HA-0016)
 Covington (KY-KE-0004) - 3 meters (9.8 ft)
 Crosby Road (OH-HA-0017) - 63 ft (19.2 meters)
 East Fork Reservoir spillway/Wm. A. Harsha Lake (OH-CT-0007)
 Eightmile Creek (OH-HA-0034)
 Emming Street (OH-HA-0002)
 Fairview Park (OH-HA-0001) - original type section of Fairview Formation
 Fort Thomas (KY-CP-0001)
 Galbraith Road, cliff section (OH-HA-0033) - 73 ft (22.3 meters)
 Georgetown (OH-BR-0006) - 9.2 meters (30.2 ft)
 Happy Hollow (OH-CT-0013)
 I-275/Turkeyfoot Road (KY-KE-0005) - 0.7 meter (2.3 ft)
 Jackson Hill Park (OH-HA-0018) - 33.2 meters (109 ft)
 Johns Hill (KY-CP-0005) - 8.0 meters (26.2 ft)
 Miami Station (OH-HA-0019) - 29 ft (8.8 meters)
 Miamitown/Hamilton-Cleves exit (OH-HA-0009) - 0.2 meter (0.7 ft)
 Montana Avenue (OH-HA-0038) - 0.3 meter (1.0 ft)

Muddy Creek (OH-HA-0020) - 67 ft (20.4 meters)
 Newport-Grand Avenue (KY-CP-0004) - 2.9 meters (9.5 ft)
 Pabco/Timberview Apartments (OH-HA-0037) - 0.5 meter (1.6 ft), lower contact
 Riedlin Road/Mason Road (KY-KE-0001) - 32.6 meters (107 ft), both contacts
 Riverside (OH-HA-0021) - 39 ft (11.9 meters), lower contact
 Sekitan Road (OH-HA-0022) - 26 ft (7.9 meters)
 Sheits Road (OH-HA-0023) - 38 ft (11.6 meters), upper contact
 Southgate (KY-CP-0003) - 0.5 meter (1.6 ft)
 Stonelick Creek (OH-CT-0004) - Fairview-Bellevue contact at Belfast-Owensville Road bridge over Stonelick Creek, about 1.2 miles (2 km) southeast of Ohio Rte. 131 at 39°09'40"N, 84°07'35"W
 Tanners Creek (IN-D-0003) - 41 ft (12.5 meters), lower contact
 Wesselman (OH-HA-0011) - above and below rock of the Wesselman Tongue of Kope Formation
 West Fork (OH-HA-0024) - 43 ft (13.11 meters)

North Bend Tongue of Fairview Formation

Briarly Creek (OH-HA-0014) - 11 ft (3.35 meters)
 Congress Run (OH-HA-0016)
 Galbraith Road, cliff section (OH-HA-0033) - 18 ft (5.5 meters)
 Miami Station (OH-HA-0019) - 8 ft (2.4 meters)
 Miamitown/Hamilton-Cleves exit (OH-HA-0009)
 Muddy Creek (OH-HA-0020) - 20 ft (6.1 meters)
 North Bend (OH-HA-0010) - type section
 Sekitan Road (OH-HA-0022) - 10 ft (3.0 meters)
 West Fork (OH-HA-0024) - 19 ft (5.8 meters)

Mt. Hope Member of Fairview Formation

Maysville (KY-MS-0004) - 40.8 meters (133.9 ft)

Fairmount Member of Fairview Formation

Maysville (KY-MS-0004) - 21.6 meters (70.9 ft)

Miamitown Shale

Aiken School (OH-HA-0012) - 5 ft (1.5 meters)
 Bald Knob (OH-HA-0013)
 Batavia #2 (OH-CT-0002) - 0.9-1.2 meters (3.0-3.9 ft)
 Bellevue Hill (OH-HA-0003) - 5 ft (1.5 meters)
 Briarly Creek (OH-HA-0014)
 Brookville Dam spillway (IN-FR-0002)
 Camp Claybanks (OH-HA-0015) - 8 ft (2.4 meters)
 Congress Run (OH-HA-0016)
 Crosby Road (OH-HA-0017) - 21 ft (6.4 meters)
 Emming Street (OH-HA-0002)
 Fairview Park (OH-HA-0001)
 Galbraith Road, cliff section (OH-HA-0033) - 7 ft (2.1 meters)
 Jackson Hill Park (OH-HA-0018) - 1.2 meters (4.0 ft)
 Miami Station (OH-HA-0019)
 Miamitown West (OH-HA-0008) - 5.2 meters (17.0 ft)
 Muddy Creek (OH-HA-0020) - 5 ft (1.5 meters)
 Rice and Gage Streets (OH-HA-0004)
 Riedlin Road/Mason Road (KY-KE-0001) - 1.0 meter (3.3 ft), both contacts
 Sekitan Road (OH-HA-0022)
 Sheits Road (OH-HA-0023) - 14 ft (4.3 meters), both contacts

West Fork (OH-HA-0024)
Wynbrook Apartments (OH-HA-0025)

Bellevue Limestone Member of Grant Lake Formation

Aiken School (OH-HA-0012) - 23 ft (7.0 meters)
Aurora (IN-D-0001)
Backbone Creek (OH-CT-0006) - 8 meters (26.2 ft)
Batavia #2 (OH-CT-0002) - 7.9 meters (25.9 ft)
Bellevue Hill (OH-HA-0003) - 22 ft (6.7 meters)
Brookville Dam spillway (IN-FR-0002)
Camp Claybanks (OH-HA-0015) - 3 ft (0.9 meter)
Emming Street (OH-HA-0002)
Fairview Park (OH-HA-0001)
Galbraith Road, cliff section (OH-HA-0033) - 20 ft (6.1 meters)
Georgetown (OH-BR-0006) - 14.8 meters (48.6 ft)
Jackson Hill Park (OH-HA-0018) - 6.71 meters (22 ft)
Madison South road cut (IN-JE-0003) - 0.25 meter (0.8 ft)
Maysville (KY-MS-0004) - 22.6 meters (74.1 ft)
Miamiwest West (OH-HA-0008) - 2.1 meters (7 ft)
Muddy Creek (OH-HA-0020) - 29 ft (8.8 meters)
Rice and Gage Streets (OH-HA-0004)
Riedlin Road/Mason Road (KY-KE-0001) - 4.5 meters (14.7 ft), lower contact
Sheits Road (OH-HA-0023) - 42 ft (12.8 meters)
Stonelick Creek (OH-CT-0004) - Fairview-Bellevue contact at Belfast-Owensville Road bridge over Stonelick Creek, about 1.2 miles (2 km) southeast of Ohio 131 at 39°09'40"N, 84°07'35"W
Wynbrook Apartments (OH-HA-0025)

REFERENCES CITED

- Aigner, Thomas, 1982, Calcareous tempestites: Storm-dominated stratification in Upper Muschelkalk limestones (Middle Trias, SW-Germany), *in* Einsele, G., and Seilacher, A., eds., *Cyclic and event stratification*: Berlin, Springer-Verlag, p. 180-198.
- _____, 1985a, Dynamic stratigraphy of the Upper Muschelkalk, South-German Basin, *in* Bayer, U., and Seilacher, A., eds., *Sedimentary and evolutionary cycles*: Berlin, Springer-Verlag, p. 342-346.
- _____, 1985b, Storm depositional systems: dynamic stratigraphy in modern and ancient shallow-marine sequences: Berlin, Springer-Verlag, 174 p.
- Alexander, R. R., 1975, Phenotypic lability of the brachiopod *Rafinesquina alternata* (Ordovician) and its correlation with the sedimentologic regime: *Journal of Paleontology*, v. 49, p. 607-618.
- Anstey, R. L., and Fowler, M. L., 1969, Lithostratigraphy and depositional environment of the Eden Shale (Ordovician) in the tri-state area of Indiana, Kentucky, and Ohio: *Journal of Geology*, v. 77, p. 668-682.
- Anstey, R. L., and Perry, T. G., 1972 (1973), Eden Shale bryozoans: a numerical study (Ordovician, Ohio Valley): Michigan State University, Publications of the Museum, Paleontological Series, v. 1, no. 1, 80 p.
- Bassler, R. S., 1906, A study of the James types of Ordovician and Silurian Bryozoa: *Proceedings of the U.S. National Museum*, v. 30, no. 1442, p. 1-66.
- Bayer, U., Althamer, E., Deutschle, W., 1985, Environmental evolution in shallow epicontinental seas: sedimentary cycles and bed formation, *in* Bayer, U., and Seilacher, A., eds., *Sedimentary and evolutionary cycles*: Berlin, Springer-Verlag, p. 347-381.
- Brandt, D. S., 1980, Phenotypic variation and paleoecology of *Flexicalymene* (Arthropoda: Trilobita) in the Cincinnati Series (Upper Ordovician) near Cincinnati, Ohio: M.S. thesis (unpub.), University of Cincinnati, 148 p.
- Brett, C. E., 1983, Sedimentology, facies and depositional environments of the Rochester Shale (Silurian; Wenlockian) in western New York and Ontario: *Journal of Sedimentary Petrology*, v. 53, p. 947-971.
- Brett, C. E., Speyer, S. E., and Baird, G. C., 1986, Storm-generated sedimentary units: Tempestite proximity and event stratification in the Middle Devonian Hamilton Group of New York: *New York State Museum Bulletin* 457, p. 129-156.
- Brown, G. D., Jr., and Lineback, J. A., 1966, Lithostratigraphy of the Cincinnati Series (Upper Ordovician) in southeastern Indiana: *Bulletin of the American Association of Petroleum Geologists*, v. 50, p. 1018-1023.
- Bucher, W. H., 1917, Large current-ripples as indicators of paleogeography: *Proceedings of the U.S. National Academy of Sciences*, v. 3, p. 285-291.
- Carpenter, J. W., and Ory, T. R., 1961, The American Upper Ordovician Standard. VI. The Covington Sequence at Maysville, Kentucky: *Ohio Journal of Science*, v. 61, p. 372-378.
- Caster, K. E., Dalvé, E. A., and Pope, J. K., 1955, Elementary guide to the fossils and strata of the Ordovician in the vicinity of Cincinnati, Ohio: Cincinnati Museum of Natural History, 47 p.
- Cummings, E. R., 1922, Nomenclature and description of the geological formations of Indiana, *in* Logan, W. N., and others, *Handbook of Indiana geology*: Indiana Department of Conservation Publication 21, p. 403-570.
- Diekmeyer, S. C., 1990, Quantitative analysis of cyclical faunal patterns in the Upper Ordovician Kope through Bellevue sequence, Cincinnati, Ohio: M.S. thesis (unpub.), University of Cincinnati, 168 p.
- Dunham, R. J., 1962, Classification of carbonate rocks according to depositional texture, *in* Ham, W. E., ed., *Classification of carbonate rocks*: American Association of Petroleum Geologists Memoir 1, p. 108-121.
- Einsele, G., 1985, Response of sediments to sea-level changes in differing subsiding storm-dominated marginal and epeiric basins, *in* Bayer, U., and Seilacher, A., eds., *Sedimentary and evolutionary cycles*: Berlin, Springer-Verlag, p. 68-97.
- Foerste, A. F., 1905, The classification of the Ordovician rocks of Ohio and Indiana: *Science*, new ser., v. 22, no. 553, p. 149-152.
- Ford, J. P., 1965, Bedrock geology in southwest Hamilton County, Ohio: Ph.D. dissertation (unpub.), The Ohio State University, 103 p.
- _____, 1967, Cincinnati geology in southwest Hamilton County, Ohio: American Association of Petroleum Geologists Bulletin, v. 51, p. 918-936.
- _____, 1974, Bedrock geology of the Cincinnati West quadrangle and part of the Covington quadrangle, Hamilton County, Ohio: Ohio Division of Geological Survey Report of Investigations 93, 1 sheet.
- Fox, W. T., 1962, Stratigraphy and paleoecology of the Richmond Group in southeastern Indiana: *Geological Society of America Bulletin*, v. 73, p. 621-641.
- Fürsich, F. T., 1974, On *Diplocraterion* Torell, 1870 and the significance of morphological features in vertical spreiten-bearing U-shaped trace fossils: *Journal of Paleontology*, v. 48, p. 952-962.
- _____, 1975, Trace fossils as environmental indicators in the Corallian of England and Normandy: *Lethaia*, v. 8, p. 151-173.
- Goodwin, P. W., and Anderson, E. J., 1980, Punctuated aggradational cycles: a general hypothesis of stratigraphic accumulation (abs.): *Geological Society of America Abstracts with Programs*, v. 12, p. 436.
- Gutstadt, A. M., 1958, Upper Ordovician stratigraphy in eastern interior region: American Association of Petroleum Geologists Bulletin, v. 42, p. 513-547.
- Hall, James, 1842, Notes upon the geology of the western states: *American Journal of Science*, v. 42, p. 51-62.
- Harris, F. W., and Martin, W. D., 1979, Benthic community development in limestone beds of the Waynesville (upper Dillsboro) Formation (Cincinnati Series, Upper Ordovician) of southeastern Indiana: *Journal of Sedimentary Petrology*, v. 49, p. 1295-1306.
- Hay, H. B., 1981, Lithofacies and formations of the Cincinnati

- Series (Upper Ordovician), southeastern Indiana and southwestern Ohio: Ph.D. dissertation (unpub.), Miami University, 236 p.
- Hofmann, H. J., 1966, Ordovician paleocurrents near Cincinnati, Ohio: *Journal of Geology*, v. 74, p. 868-890.
- Holland, S. M., 1990, Distinguishing eustasy and tectonics in foreland basin sequences: the Upper Ordovician of the Cincinnati Arch and Appalachian Basin: Ph.D. dissertation (unpub.), University of Chicago, 390 p.
- _____, 1993, Sequence stratigraphy of a carbonate-clastic ramp: the Cincinnati Series (Upper Ordovician) in its type area: *Geological Society of America Bulletin*, v. 105, p. 306-322.
- Jennette, D. C., 1986, Storm-dominated cyclic ramp deposits of the Kope-Fairview transition (Upper Ordovician), southwestern Ohio and northern Kentucky: M.S. thesis (unpub.), University of Cincinnati, 210 p.
- Jennette, D. C., and Pryor, W. A., 1993, Cyclic alternation of proximal and distal storm facies: Kope and Fairview Formations (Upper Ordovician), Ohio and Kentucky: *Journal of Sedimentary Petrology*, v. 63, p. 183-203.
- Kidwell, S. M., 1986, Taphonomic feedback in Miocene assemblages: testing the role of dead hardparts in benthic communities: *Palaios*, v. 1, p. 239-255.
- Kidwell, S. M., Fürsich, F. T., and Aigner, T., 1986, Conceptual framework for the analysis and classification of fossil concentrations: *Palaios*, v. 1, p. 228-238.
- Kidwell, S. M., and Jablonski, D., 1983, Taphonomic feedback: ecological consequences of shell accumulation, in Tevesz, M. J., and McCall, P. L., eds., *Biotic interactions in recent and fossil benthic communities*: New York, Plenum, p. 195-248.
- Kreisa, R. D., 1981a, Origin of stratification in a Paleozoic epicontinental sea: the Cincinnati Series (abs.): *Geological Society of America Abstracts with Programs*, v. 13, no. 7, p. 491.
- _____, 1981b, Storm-generated sedimentary structures in subtidal marine facies with examples from the Middle and Upper Ordovician of southwestern Virginia: *Journal of Sedimentary Petrology*, v. 51, p. 823-843.
- Krumpolz, B. J., 1980, Paleogeology, biostratigraphy, and sedimentology of the Mt. Auburn Member (McMillan Formation, uppermost Maysvillian) in the area of Cincinnati, Ohio: M.S. thesis (unpub.), University of Cincinnati, 196 p.
- Locke, John, 1838, Prof. Locke's geological report [on southwestern Ohio]: Ohio Division of Geological Survey Second Annual Report, p. 201-286.
- Luft, S. J., 1971, Geologic map of part of the Covington quadrangle, northern Kentucky: U.S. Geological Survey Geologic Quadrangle Map GQ-955, scale 1:24,000.
- McFarlan, A. C., and Nosow, Edmund, 1961, [Ordovician section] in Schmidt, R. G., McFarlan, A. C., Nosow, Edmund, Bowman, R. S., and Alberts, Robert, Examination of the Ordovician through Devonian stratigraphy and the Serpent Mound chaotic structure area: Guidebook for Field Trips, Cincinnati meeting, 1961, Geological Society of America, p. 270-280.
- Meek, F. B., and Worthen, A. H., 1865, Descriptions of new species of Crinoidea, etc., from the Paleozoic rocks of Illinois and some of the adjoining states: *Academy of Natural Sciences of Philadelphia Proceedings*, v. 17, p. 143-155.
- Meyer, D. L., and Meyer, K. B., 1986, Biostratigraphy of recent crinoids (Echinodermata) at Lizard Island, Great Barrier Reef, Australia: *Palaios*, v. 1, p. 294-302.
- Meyer, D. L., Diekmeyer, S. D., and Holterhoff, P. F., 1990, Crinoid distribution in shoaling-upward carbonate-clastic sequences: Upper Ordovician (Maysvillian) of Ohio and Kentucky, U.S.A. (abs.): Atami, Japan, Seventh International Echinoderm Conference, Abstracts with Programs, p. 29.
- Meyer, D. L., Schumacher, G. A., Swinford, E. M., Jennette, D. C., and Brockman, C. S., 1985, Upper Ordovician stratigraphy, sedimentology, and paleontology along Backbone Creek, Clermont County, Ohio: Field Trip Guidebook for Annual Meeting of Ohio Academy of Science, Geology Section, 14 p.
- Meyer, D. L., Tobin, R. C., Pryor, W. A., Harrison, W. B., Osgood, R. G., Hinterlong, G. D., Krumpolz, B. J., and Mahan, T. K., 1981, Stratigraphy, sedimentology, and paleoecology of the Cincinnati Series (Upper Ordovician) in the vicinity of Cincinnati, Ohio, in Roberts, T. G., ed., *GSA Cincinnati '81 Field Trip Guidebooks*: American Geological Institute, v. 1, p. 31-72.
- Nickles, J. M., 1902, The geology of Cincinnati: Cincinnati Society of Natural History Journal, v. 20, p. 49-100; reprint edition, p. 1-53.
- Orton, Edward, 1873, Report on the Third Geological District; geology of the Cincinnati Group; Hamilton, Clermont, Warren, and Butler Counties: Ohio Division of Geological Survey, v. 1, pt. 1, Geology, p. 365-480.
- Osborne, R. H., 1971, The American Upper Ordovician Standard. XIV. Markov analysis of typical Cincinnati sedimentation: *Journal of Sedimentary Petrology*, v. 41, p. 444-449.
- Palmquist, W. N., Jr., and Hall, F. R., 1960, Availability of ground water in Bracken, Harrison, Mason, Nicholas, and Robertson Counties, Kentucky: U.S. Geological Survey Hydrologic Investigations Atlas HA-16, scale 1:125,000.
- Peck, J. H., 1966, Upper Ordovician formations in the Maysville area, Kentucky: U.S. Geological Survey Bulletin 1244-B, 30 p.
- Schumacher, G. A., 1984, Sedimentology, biostratigraphy, and paleoecology of Cincinnati (Upper Ordovician) *Iocrinus subcrassus* crinoid (Echinodermata) assemblages: M.S. thesis (unpub.), University of Cincinnati, 210 p.
- Scotese, C. R., and Denham, C. R., 1988, User's manual for TERRA MOBILIS: plate tectonics for the Macintosh: privately printed by Earth in Motion Technologies, 6601 Kirby Drive, Suite 529, Houston, Texas, 77005, 43 p.
- Tobin, R. C., 1982, A model for cyclic deposition in the Cincinnati Series of southwestern Ohio, northern Kentucky, and southeastern Indiana: Ph.D. dissertation (unpub.), University of Cincinnati, 483 p.
- _____, 1986, An assessment of the lithostratigraphic and interpretive value of the traditional "biostratigraphy" of the type Upper Ordovician of North America: *American Journal of Science*, v. 286, p. 673-701.
- Tobin, R. C., and Pryor, W. A., 1985, Cincinnati Series—model for cyclic and episodic deposition of carbonates and shales on a storm-dominated ramp (abs.): *American Association of Petroleum Geologists Bulletin*, v. 69, p. 311-312.
- Weir, G. W., Peterson, W. L., Swadley, W. C., and Pojeta, John, Jr., 1984, Lithostratigraphy of Upper Ordovician strata exposed in Kentucky; with a section on biostratigraphy: U.S. Geological Survey Professional Paper 1151-E, 121 p.
- Weiss, M. P., 1961, The American Upper Ordovician Standard. V. A critical appraisal of the classification of the typical Cincinnati beds: *Geological Society of America Bulletin*, v. 72, p. 645-647.
- Weiss, M. P., Edwards, W. R., Norman, C. E., and Sharp, E. R., 1965, The American Upper Ordovician Standard. VII. Stratigraphy and petrology of the Cynthiana and Eden Formations of the Ohio Valley: *Geological Society of America Special Paper* 81, 76 p.
- Weiss, M. P., and Sweet, W. C., 1964, Kope Formation (Upper Ordovician): Ohio and Kentucky: *Science*, v. 145, no. 3638, p. 1296-1302.
- Winchell, N. H., and Ulrich, E. O., 1897, The Lower Silurian deposits of the upper Mississippi province: a correlation of the strata with those in the Cincinnati, Tennessee, New York and Canadian Provinces, and the stratigraphic and geographic distribution of the fossils, in Ulrich, E. O., and others, *The geology of Minnesota: Geological and Natural History Survey of Minnesota*, v. 3, pt. 2, Paleontology, p. lxxxiii-cxxviii.
- Ziegler, A. M., 1981, Paleozoic biogeography and climatology, in Niklas, K. J., ed., *Paleobotany, paleoecology, and evolution*: New York, Praeger Publishing, p. 231-266.

Address of author: Sharon C. St. Louis Diekmeyer, Department of Geology, University of Cincinnati, Cincinnati, OH 45221-0013.

FIGURE 3-16.—Cincinnatian brachiopods, crinoids, trilobites, and bryozoans. Scales bars = 1 cm; UCGM = University of Cincinnati Geology Museum; UCGM PT = University of Cincinnati Paleontology Teaching Collection.

- 1 *Zygospira cincinnatiensis* Meek, 1873, UCGM 7351.
- 2 *Onniella meeki* (S. A. Miller, 1875), University of Cincinnati Teaching Collection.
- 3 *Ectenocrinus* sp.
- 4 *Isotelus maximus* Locke, 1838, UCGM PT 220; note hypostome (circled).
- 5 *Flexicalymene meeki* (Foerste, 1910), UCGM 40681; note disarticulation in the specimen on the right.
- 6 *Batostoma jamesi* (Nicholson, 1874), UCGM.
- 7 *Homotrypa wortheni* (James, 1882), UCGM P527.




FIGURE 3-17.—Cincinnatian brachiopods, crinoids, and bryozoans. Scales bars = 1 cm; UCGM = University of Cincinnati Geology Museum; UCGM PT = University of Cincinnati Paleontology Teaching Collection.

- 1a, b *Platystrophia laticosta* (Meek, 1873), brachial and pedicle valve, respectively.
- 2 *Plectorthis fissicosta* (Hall, 1847), University of Cincinnati Teaching Collection.
- 3a *Strophomena planoconvexa* (Hall, 1847), brachial valve, exterior, UCGM 8-260.
- 3b *Strophomena planumbona* (Hall, 1847), pedicle valve, interior; note prominent muscle scars, University of Cincinnati Teaching Collection.
- 4 *Iocrinus subcrassus* (Meek & Worthen, 1865), part of stem.
- 5 *Pycnocrinus dyeri* (Meek, 1872), calyx, UCGM 17596.
- 6 *Dekayia aspera* Milne-Edwards & Haime, 1851, UCGM 20398.
- 7 *Escharopora falciformis* (Nicholson, 1875), UCGM.
- 8 *Constellaria florida* Ulrich, 1882.
- 9 *Heterotrypa frondosa* (d'Orbigny, 1850), UCGM 20390.





FIGURE 3-18.—Cincinnatian brachiopods, mollusks, and bryozoans. Scales bars = 1 cm; UCGM = University of Cincinnati Geology Museum; UCGM PT = University of Cincinnati Paleontology Teaching Collection.

- 1a, b *Hebertella occidentalis* Hall, 1847, brachial and pedicle valve, respectively, University of Cincinnati Teaching Collection.
- 2 *Platystrophia ponderosa* Foerste, 1909, pedicle valve, University of Cincinnati Teaching Collection.
- 3 *Rafinesquina alternata* (Emmons, 1842), pedicle valve, UCGM P238.
- 4 *Caritodens demissa* (Conrad, 1842), UCGM.
- 5 Modiolopsid pelecypod, internal mold.
- 6 *Cyclonema inflatum* Ulrich, 1897.
- 7 *Ambonychia* sp., internal mold; note cyclostome bryozoans on surface of mold, University of Cincinnati Teaching Collection.
- 8 ?*Orthonybyoceras* sp., internal mold with encrusting bryozoans.
- 9 *Parvohallopore ramosa* (d'Orbigny, 1850).
- 10 *Monticulipora molesta* Nicholson, 1881, UCGM 20356.



4. UPPER KOPE (MCMICKEN) COLLECTING ON THE MIAMITOWN/HAMILTON-CLEVES EXIT RAMP (UPPER ORDOVICIAN, SOUTHWESTERMOST OHIO)

by
Roger J. Cuffey

SIGNIFICANCE

This locality (OH-HA-0009) provides excellent access to a prolific bryozoan fauna of late Edenian age. This fauna has been thoroughly studied by Anstey and Perry (1972) and thus serves as a comparative standard for that phylum and age within the type-Cincinnatian region.

LOCATION

The strata yielding this bryozoan fauna are exposed in the road cut along the south side (away from the main highway) of the exit ramp in the southwestern quadrant of the interchange between I-74 (here also I-275) and Ohio Route 128 (Hamilton-Cleves Road), 0.8 mile (1.3 km) south-southwest of the crossroads in the center of Miamitown, Hamilton County, Ohio (fig. 4-1). This exit ramp takes the eastbound traffic off the interstate. The shoulders of both the ramp itself and the crossroad at the bottom of the ramp are wide enough to provide parking out of the way of any exiting traffic. Permission to examine the cut should be obtained from the Ohio Department of Transportation. The ramp lies in the center SE¹/₄SE¹/₄ sec. 1, T. 1 N., R. 1 E., on the Addyston, Ohio–Kentucky, 7.5-minute quadrangle, at

39°12'21"N latitude, 84°42'39"W longitude (4341928 m N, 697649 m E, UTM zone 16).

The heavy traffic on both lanes of the interstate above effectively separates this exit ramp from the road cuts along the north side of the interstate. Those cuts include the type section of the Miamitown Shale (Ford, 1967), discussed fully by Dattilo (paper 7 in this volume; OH-HA-0008).

STRATIGRAPHY AND PALEONTOLOGY

The road cut (fig. 4-2) exposes about 20 meters (60 ft) of gray shale with minor, interbedded, thin, bioclastic to barren limestones, characteristic of the upper part of the Kope Formation, equivalent to the McMicken of the traditional type-Cincinnatian units (Anstey and Perry, 1972, p. 35, 77; they used the term Eden Shale). Most abundant and most conspicuous of the numerous fossils weathering out on the outcrop surface are branching bryozoan fragments, mostly trepostomes. Only five species are present; this is a surprisingly low diversity for such an abundant fauna. The fauna has been interpreted as indicating deep-water conditions (Anstey and Perry, 1972, p. 37, 38, 42, 44). The species documented here by Anstey and Perry (1972, p. 28, 35, 49-52, 59-61, 63-75) are *Amplexopora septosa* (Ulrich,




FIGURE 4-1.—Locations of the road cut in the Kope Formation at the Hamilton-Cleves exit ramp (OH-HA-0009; XR on map) and of the Miamitown Shale type section on north side of the Interstate (OH-HA-0008; MTS on map). Addyston, Ohio–Kentucky, 7.5-minute quadrangle.



FIGURE 4-2.—Bryozoan-rich strata of the upper Kope along the south side of Hamilton-Cleves exit ramp; photograph taken in 1988; the thicker bedded limestones in the upper part of the cut could be the Grand Avenue Member of the Kope Formation, although that unit reputedly pinches out east of this site (Ford, 1967).

1879) Cumings, 1908, *Dekayia aspera* Milne-Edwards and Haime, 1851, *Hallopora nodulosa* (Nicholson, 1874) Anstey and Perry, 1973, *Heterotrypa ulrichi* Nicholson, 1881, and *Peronopora vera* Ulrich, 1888.

REFERENCES CITED

- Anstey, R. L., and Perry, T. G., 1972 (1973), Eden Shale bryozoans: a numerical study (Ordovician, Ohio Valley): Michigan State University, Publications of the Museum, Paleontological Series, v. 1, no. 1, 80 p.
- Ford, J. P., 1967, Cincinnati geology in southwest Hamilton County, Ohio: American Association of Petroleum Geologists Bulletin, v. 51, p. 918-936.

Address of author: Roger J. Cuffey, Department of Geosciences (Deike Bldg.), Pennsylvania State University, University Park, PA 16802.

5. THE MAYSVILLE BRYOZOAN REEF MOUNDS IN THE GRANT LAKE LIMESTONE (UPPER ORDOVICIAN) OF NORTH-CENTRAL KENTUCKY

by
Roger J. Cuffey

Editor's note: Unfortunately, between 1992 and 1998, highway widening destroyed the part of the road cut that contained these unusual reef mounds. However, this paper is presented here as it was written in 1992.

SIGNIFICANCE

Two small reef mounds in the Grant Lake Limestone (upper Maysvillian) south of Maysville, Kentucky, were built almost entirely by the trepostome bryozoan *Stigmarella personata* Ulrich and Bassler, 1904. These Maysville bryozoan bioherms are significant because they represent one of the best developed and most accessible examples of bryozoan reefs known anywhere in the world and because they constitute a mode of bryozoan occurrence that is unique within the type-Cincinnatian.

Bryozoan reefs represent ecological extremes for their phylum and exotic rarities among bioherms in general (Cuffey, 1977, 1985, 1987). Only about 100 are known, especially from the mid-Ordovician, late Paleozoic, and mid-Cenozoic, and largely in North America and Eurasia; many are in remote or hazardous locations. In some, the importance of the bryozoans may be obscured by intermixture with animals of other phyla or by later development of diagenetic cements. As a result, the well-preserved, completely bryozoan, easily accessible (drive up and park!) Maysville mounds are exquisite examples of these rare structures.

The type-Cincinnatian is famous for its bryozoan fossils, but none are reefal except for this Maysville occurrence, which is therefore of exceptional interest in the region. The normal mode of bryozoan occurrence is as broken branch fragments and small crusts or masses; all individual colonies are not interconnected, even though they may be abundant (Anstey and Fowler, 1969; Harris and Martin, 1979; Errett and Cuffey, 1989). Moreover, in contrast to the preceding epoch, the Late Ordovician in North America overall saw virtually no bryozoan reef construction; the only other instance known so far from this age is at Mullet Creek near Toronto, Ontario (Kobluk, 1980).

HISTORY OF STUDY

The Maysville bryozoan bioherms were first reported by Kissling and Turonis (1977). Later, their bryozoan species composition, constructional roles, reef-rock classification, and bryozoan reef type were noted (Cuffey, 1985, p. 309; Cuffey and Kamandulis, 1985). The Maysville mounds are under long-term detailed study (Roger A. Clark and Don L. Kissling, written communication, 1990).

LOCATION AND DIRECTIONS

The Maysville bryozoan reef mounds (KY-MS-0006) are two low limestone mounds, each about 3 meters (10 ft) wide by 0.3 meter (1 ft) high and separated by 2 meters (6 ft). They are exposed at road level in the ditch at the base of the 6-meter-high (20-ft) outcrop face on the west side of U.S. Route 68 0.1 mile (0.2 km) south of the junction of U.S. Route 68 and U.S. Route 62 at the south edge of the village of Wash-




FIGURE 5-1.—Location of the Maysville bryozoan reef mounds (MBR and arrow). Topographic map is from Mays Lick, Kentucky, 7.5-minute quadrangle.

ington, Kentucky (fig. 5-1). That junction is 5.0 miles (8.0 km) by road south of the southern end of the U.S. 68 bridge over the Ohio River at Maysville, or equivalently 4.3 miles (6.9 km) by road south of the junction of U.S. Route 68 and Kentucky Route 11 at the south-central edge of Maysville. One can park on the road shoulder next to the reefs; both the passing traffic and the roadside ditch must be avoided carefully. The condition of the ditch has varied considerably over the years; it was full of rubble which practically buried the reefs in 1982, but had been scooped out by 1984 and 1988 so that the reefs and their substratum were exposed as they had been in 1975 and 1976.

The Maysville bryozoan bioherms (or "bryoherms"; Cuffey,

1977, p. 185) are in Mason County, Kentucky, on the Mays Lick, Kentucky, 7.5-minute quadrangle (also available as a geologic quadrangle, Gibbons, 1968), at 38°36'36"N latitude and 83°48'39"W longitude (4277049 m N and 255255 m E, UTM zone 17). Neither section-township-range coordinates nor political township names are available for this part of Kentucky.

Note that the Maysville bryoherms are small, as well as quite unusual, and are under current study. Examination of these intriguing structures therefore should be cautious and conservation oriented in order to maximize their potential contribution to our understanding of bryozoans, reefs, and the type-Cincinnatian. At the same time, however, making these reef mounds readily available to visiting scientists also serves those goals; hence, this present brief paper.

STRATIGRAPHY AND REGIONAL PALEOENVIRONMENTS

The Maysville bryozoan mounds are stratigraphically within the upper Maysvillian, in the Grant Lake Limestone, approximately at the level of the Mt. Auburn-Corryville contact to the northwest (fig. 5-2) (Gibbons, 1968; Schilling and Peck, 1967; Don L. Kissling, written communication, 1984). The outcrop face above the reefs exposes rubbly, nodular, irregularly bedded limestone with minor shale partings characteristic of the upper member of the Grant Lake (Mt. Auburn equivalent). The reefs are founded upon a flat-bedded limestone and buried under a thin calcareous shale; such evenly bedded limestones with more shale beds characterize the middle member of the Grant Lake (Corryville equivalent). The lower member of the Grant Lake, equivalent to the Bellevue, is well below the road bed at the site of the reefs and not visible, but is rubbly limestone with little or no shale. The Mt. Auburn equivalent in the Maysville area has been renamed the Straight Creek Member by Schumacher and others (1991).

Five km (3 miles) northeast of this reef locality, essen-




FIGURE 5-2.—Stratigraphic succession containing the Maysville bryozoan reef mounds. Nomenclature and lithologic symbols after Gibbons (1968) and Schilling and Peck (1967); the Mt. Auburn Member is the Straight Creek Member of Schumacher and others (1991). Unit thicknesses were derived from elevations of contacts mapped on Gibbons (1968) at and near the location of the reefs.

tially the same horizon as that of the mounds—the basal Mt. Auburn or Straight Creek—carries massive bryozoans, including the same genus which is dominant here (Schmidt and others, 1961, p. 272-273, 279-280), but they are not developed into biohermal accumulations. Similar colonies but not bioherms also were noted there by those authors at a lower horizon, the basal Corryville or topmost Bellevue. Likewise, no additional bryozoan bioherms thus far have been found at either horizon within several kilometers from the reef-mound locality, in spite of extensive searches by both Kissling and Cuffey.

As noted in the introductory paper for this volume (Cuffey, paper 2), stratigraphic correlations and inferred paleoenvironments of the type-Cincinnatian can be coordinated and summarized as background specifically for bryozoan investigations. Plotting the environments indicated by the rock types of the formations correlated with the horizon of the Maysville bryoherms yields a regional cross section or profile (fig. 5-3) within which the reef mounds can be seen in broad paleoenvironmental context: out on a tropical, shallow, offshore, carbonate platform, flanked by deeper, muddy-bottomed water farther offshore to the northwest.

REEF-MOUND CHARACTERISTICS AND LOCAL PALEOECOLOGY

REEF-MOUND GEOMETRY

Both mounds are bisected by the vertical face of the outcrop and so reveal their cross-sectional shape well but not much of their plan views. Overall, each appears as a low, wide, partially flat topped dome (fig. 5-4), with a possibly circular, oval, or even irregular outline horizontally. When measured in 1984, following at least one burial and re-exposure by highway-department road-ditch scraping, the outcrop had been cut back enough to show the reef mounds well (figs. 5-5A, 5-5B); the southern mound then was 3 meters (10 ft) wide at its base, the northern one 3.4 meters (11.2 ft) wide, and both only 0.3 meter (1.0 ft) high. In life, therefore, the mounds would have appeared as low rocky rises several cm above the surrounding shallow sea floor, whether muddy or hardground.

Such low relief over the reef mounds is not likely to represent the total water depth around the mounds during their growth, especially because the tops of the mounds show no effects of having been battered by surf action. Instead, note that the living Bahamian bryozoan reefs (Cuffey and others, 1977), which are roughly similar in size to the Maysville ones, lie under 3-4 meters (10-12 ft) of water even at low tide, a depth which has also been suggested for the Cincinnatian lithofacies enclosing the mounds from regional paleoenvironmental interpretations (Cuffey, paper 2 in this volume).

REEF-ROCK CHARACTERISTICS

The Maysville bioherms are massive to nodular limestones, even viewed from a distance, but appear quite different from the typical Cincinnatian limestone layers all around them. Close examination reveals that the bioherms consist of very large, solid, tubulobryozoan crusts, masses, and domes, overgrowing and attached one atop another (figs. 5-5C, 5-5D, 5-6A, and 5-6B). In places, they are tightly cemented to each other; in others, they are only loosely attached, because of an intervening thin film of mud. Thus,




FIGURE 5-3.—Generalized paleoenvironmental profile across the Cincinnati region at the time of growth of the Maysville bryozoan reef mounds. The scale is approximate. Lithofacies and paleoenvironments from figure 2-2 and correlations from figure 2-4 of Cuffey (paper 2 in this volume).




FIGURE 5-4.—Vertical cross section of the two Maysville bryozoan reef mounds, as measured by Cuffey and Kamandulis in 1984; no vertical exaggeration.

the biohermal rock ranges from solidly massive to rubbly nodular along the exposure. In the expanded petrographic reef-rock and reef-type classification of Cuffey (1985), the Maysville bryozoan bioherms are crust mounds composed of trepostome globstones and cruststones. They exemplify the reef-constructive role of principal frame building by bryozoans alone (Cuffey, 1977).

The individual colonies vary greatly in size, but the majority are 5-20 cm wide by 1-4 cm high. Maximum observed size is 95 cm across and 15 cm thick (Kissling and Turonis, 1977; Don L. Kissling, written communication, 1984). Broken surfaces of colonies appear finely striated, due to the thin, elongate, tubular zooecia. The broken colonies range from light gray to pink or white, in contrast to the medium or dark gray of the surrounding thin-bedded limestones and shales. Many successive colonies are separated by thin dark-gray bands; these consist of the mud-filled, distal ends of the zooecial tubes immediately below an exterior or upper growing surface, which had received a dusting of detrital sediment before being overgrown by the next colony.

In a few places within each mound, the edges of the bryozoan colonies curve or bend away from each other, thus creating small, irregular growth cavities inside the reefs. These cavities are now filled with detrital mud that contains a few little rhynchonellid brachiopods which originally might have lived in the cavities or might have been washed in after death. Unlike many reef cavities elsewhere, the walls and roofs of these cavities are bare; only the primary frame-building bryozoans are visible, and they are not encrusted by other bryozoans or animals of other phyla. This lack of a coelobitic, cavity-dwelling fauna suggests that mud filled the cavities as fast as the reefs grew (Kissling and Turonis, 1977).

The Maysville reefs appear to be essentially homogeneous throughout their extent; they do not exhibit any vertical or lateral ecozonation. This lack of zonation probably is due to their small size; they simply were not large enough to

develop different growing conditions across their surfaces, nor to influence water movements and physical sedimentation around them.

BRYOZOAN FRAME BUILDERS

A hundred reef-rock samples were collected according to a uniformly rectangular grid across both mounds. Each sample contained several bryozoan colonies. Every sample was sectioned in order to identify the species responsible for building the Maysville bioherms (Cuffey and Kamandulis, 1985). Only two species were found, both trepostomes.

One species is overwhelmingly predominant, constituting 98 percent of the reef-mound frame builders: *Stigmatella personata* (figs. 5-6C and 5-6D). It occurs as small to very large colonies of crustose, tabular, lenticular, domal, hemispherical, and massive forms. These all consist of zooecia packed tightly together without any intervening mesopores. The zooecia are elongate, prismatic, and thin walled, and have polygonal, angular apertures and a few tiny obscure acanthorods. The zooecia appear mostly empty, but some are crossed in places by a few diaphragms. Considerable morphologic variability is displayed among the many colonies, but extreme variants which might be misidentified as other species are all interconnected with the typical population by continuous series of intermediates. This species was described thoroughly by Cumings (1908, p. 884, pl. 24, fig. 3-3d) and by Utgaard and Perry (1964, p. 82-85, pl. 15, fig. 3-8, pl. 16, fig. 1-4). It was known previously from the Whitewater Formation down into the Waynesville Formation, so its presence at this locality represents a slight range extension downward. It has been suggested that delayed maturation due to a medical-type heterochronic mutation within the local population of this species may have been responsible for development of these Maysville reefs (Cuffey and Pursell, 1995).




FIGURE 5-5.—Photographs of the Maysville bryozoan reef mounds taken in 1984. **A**, at the base of the vertical outcrop face, Michael Kamandulis between the mounds; **B**, from the road ditch, a geologic hammer on each mound (southern mound on the left in both **A** and **B**); **C**, bryozoan globstone; and **D**, bryozoan cruststone, in place in the field (hammer for scale in both **C** and **D**).




FIGURE 5-6.—Maysville bryozoan reef rock and reef-building species; **A**, vertically broken surface, and **B**, polished surface of bryozoan cruststone pieces (ruler in cm and mm); **C**, **D**, *Stigmatella personata* (the dominant species) in tangential and longitudinal peel sections; **E**, **F**, *Atactoporella mundula* (the rare species) in tangential and longitudinal peel sections; figured sections (all X29) are in the KETMAS-1 suite (S5b, S12c, S7u, S7u, respectively) in the Paleobryozoological Research Collection at Pennsylvania State University.

A few thin encrusting colonies, only 2 percent of the reef-mound frame builders, represent *Atactoporella mundula* (Ulrich, 1879) Ulrich, 1883 (figs. 5-6E and 5-6F). Its zooecia are cylindrical and thin walled, and have round apertures, numerous large acanthorods, and some diaphragms. Large, angular mesopores containing many closely spaced diaphragms are packed in among the zooecia. Cumings (1908, p. 770, pl. 7, figs. 3-3a, pl. 26, fig. 6) and Brown and Daly (1985, p. 25, pl. 1, figs. 9-12) furnished additional descriptive data. This species has been found from the Bellevue Member of the Grant Lake Formation up through the Arnheim Formation, a range that includes the stratigraphic level of the present locality.

The extremely low biodiversity shown by the frame builders of the Maysville reefs is surprising, especially in view of the great number of bryozoan species in the type-Cincinnatian overall. Bryozoans in contemporaneous rocks surrounding the Maysville reefs (Pursell and Cuffey, 1995) are highly diverse and comprise abundant, normal-sized, isolated colonies, with well-developed mature exozones. Species identified are mostly trepostomes (the monticuliporids *Monticulipora mammulata**, *Atactoporella mundula**, *Homotrypa grandis*, *H. nodulosa*, *Homotrypella hospitalis*, *Peronopora decipiens**, *P. dubia*, and *P. pachymura**; the heterotrypids *Heterotrypa frondosa**, *H. solitaria*, *Dekayia aspera**, *D. inflecta*, *D. magna*, *D. mesospinosa*, and *D. prolifica*; the diplotrypid *Batostoma* sp.; the amplexoporids *Amplexopora cingulata*, *A. filiosa**, *A. parva*, and *A. winchelli*; the halloporids *Parvohallopora ramosa** and *P. rugosa**; and the batostomellid *Bythopora dendrina**) plus one ceramoporoid (*Ceramopora invenusta**). In addition, a few broken fragments of the reefs' principal frame builder, the heterotrypid *Stigmatella personata**, were found north and east of the reef mounds, suggestive of prevailing currents of the kind expected in such shallow-water habitats as that represented here. Many of these species—those marked with an asterisk (*)—also occur in the rock immediately underlying the reefs, as do the monticuliporid *Homotrypa flabellaris*, the heterotrypid *Leptotrypa minima*, the diplotrypid *Batostoma implicatum*, and a halloporid transitional between *Parvohallopora ramosa* and *P. subnodosa*. The few specimens of *Stigmatella personata* (the principal frame builder in the reefs) that have been found beneath the reefs are dome shaped, but normal sized, giving no hint of the reef building of the species to come. Moreover, this predominant frame-building species has been found in the region, but only as normal-sized colonies occurring individually rather than in biohermal accumulations. Recall, too, that very few bryozoan reefs are known from the North American Upper Ordovician. In our present state of incomplete knowledge concerning bryozoan reefs worldwide through time, possible explanations for such observations remain only speculative. (Note that *Parvohallopora ramosa* and *P. rugosa* may be conspecific; their distributions do overlap, but they do not coincide. It is important to record each variant wherever it turns up, because that eventually may tell us what each was. For example, perhaps the variants are a reflection of local variation in microhabitat.)

SURROUNDING SEDIMENTS

Reef foundation

The Maysville bryozoan bioherms arise from a well-indurated, calcarenitic limestone bed, 0.1 meter (0.3 ft) thick. The upper surface of the limestone apparently was a rocky, solid hardground by the time the bryozoans began to

encrust it. That surface is ferruginous in places, obscurely ripple marked in others, and extensively perforated by narrow, cylindrical, vertical borings 2-4 mm in diameter. The fact that some of those borings are directly covered by bryozoan colonies at the base of the reefs confirms that the original lime sands had lithified enough to support such borings without collapsing before encrustation by the bryozoans. Kissling and Turonis (1977) thought that the hardground might have been relatively local and that its limits might have controlled the plan-view outlines of the Maysville bioherms. However, the same foundation bed is exposed roughly 30 meters (100 ft) away on the other side of the road, but there lacks any biohermal development, although a few small zoaria of the dominant species occur widely scattered across its upper surface. The hardground also extends well beyond the edges of the mounds on the west side of the highway, although its degree of development there does vary considerably.

Among bryozoan reefs so far known, a majority are founded on sand-textured substrates (Cuffey, 1987), but a few are on surfaces that were clearly already lithified as a hardground like here.

Below the hardground surface, the foundation bed is dark-gray grainstone composed of finely comminuted shell fragments that are flat or flake shaped, lie parallel to bedding, and are spar cemented. This bed in turn is underlain by characteristic Cincinnatian limestones and shales.

Lateral flanks

As seen in the outcrop along the west side of the highway, thin-bedded limestones and shales are laterally adjacent to the Maysville bioherms and occupy the same interval. Approaching the low-domed reefs, several thin limestone beds bend slightly upward into a flanking dip of about 10°. Some of this dip may have been original depositional dip, some may result from postburial compactional drape over the solid mounds, but how much is due to each cause is unclear. Immediately adjacent to the bioherms are mostly shaly mudstones, within which continuous bedding planes are so difficult to follow that the geometric relationships between the reef rock and the flanking sediments remain obscure. Thus, it is uncertain whether the reef mounds during life were surrounded by accumulating soft muds or by the bare rock surface of the underlying hardground. Clarification of these relationships is one goal of the continuing studies of this site.

The sediments around the mounds have yielded only one or two eroded cobbles of the bryozoan reef rock and no broken fragments of the frame-building species. These small bryozoan reefs thus did not contribute any significant quantity of bioclastic debris to the surrounding sea floor.

Overburden

The two Maysville reef mounds are overlain by dark-gray shaly mudstone that is calcareous but relatively unfossiliferous. Influx of this detrital sediment, whether quickly as a result of storm activity or more slowly as a consequence of increased runoff from far-away land masses, smothered and thereby terminated reef growth here. A majority of other known bryozoan reefs likewise met their ends due to the smothering effect of burying muddy sediments (Cuffey, 1987).

This overlying mudstone in turn is succeeded by the usual type-Cincinnatian thin-bedded limestones and shales, well exposed in the outcrop face above the bioherms.

REEF DURATION

The length of time necessary for the Maysville bryozoan reefs to grow to their observed size can be estimated from two lines of evidence. First, normal-sized, robust bryozoan colonies have lifespans of several years, unusually large ones probably as much as several decades; hence, a succession of several such colonies encrusting upon one another quite possibly would represent several centuries or a millenium or two of growth. Second, the three known living bryozoan reefs—of roughly comparable size to the Maysville structures—have growth durations of several centuries or a millenium as determined from radiocarbon dating or historical records: Joulters Cays in the Bahamas, almost 1,000 years (Cuffey and Fonda, 1979); coastal Netherlands, about 300 years (Bijma and Boekschoten, 1985); and Coorong Lagoon in Australia, 700 years (Bone and Wass, 1990). Consequently, the Maysville bryozoans probably represent growth on the order of a thousand years or so.

REFERENCES CITED

- Anstey, R. L., and Fowler, M. L., 1969, Lithostratigraphy and depositional environment of the Eden Shale (Ordovician) in the tri-state area of Indiana, Kentucky, and Ohio: *Journal of Geology*, v. 77, p. 668-682.
- Bijma, J., and Boekschoten, G. J., 1985, Recent bryozoan reefs and stromatolite development in brackish inland lakes, SW Netherlands: *Senckenbergiana maritima*, v. 17, no. 1/3, p. 163-185.
- Bone, Y., and Wass, R. E., 1990, Sub-Recent bryozoan-serpulid buildups in the Coorong Lagoon, South Australia: *Australian Journal of Earth Sciences*, v. 37, p. 207-214.
- Brown, G. D., Jr., and Daly, E. J., 1985, Trepostome Bryozoa from the Dillsboro Formation (Cincinnatian Series) of southeastern Indiana: *Indiana Geological Survey Special Report 33*, 95 p.
- Cuffey, R. J., 1977, Bryozoan contributions to reefs and bioherms through geologic time: *American Association of Petroleum Geologists Studies in Geology* 4, p. 181-194.
- _____, 1985, Expanded reef-rock textural classification and the geologic history of bryozoan reefs: *Geology*, v. 13, p. 307-310.
- _____, 1987, Bryozoan reefs in relation to detrital versus carbonate sediments through geologic time (abs.): *Geological Society of America Abstracts with Programs*, v. 19, no. 7, p. 632-633.
- Cuffey, R. J., and Fonda, S. S., 1979, Bryozoan species assemblages in the modern bryozoan-rich reefs at Joulters Cays (Bahamas)—general character and paleoenvironmental implication (abs.): *Geological Society of America Abstracts with Programs*, v. 11, no. 7, p. 407-408.
- Cuffey, R. J., Gebelein, C. D., Fonda, S. S., Bliefnick, D. M., Kosich, D. F., and Soroka, L. G., 1977, Modern tidal-channel bryozoan reefs at Joulters Cays (Bahamas): *Miami, 3rd International Coral Reef Symposium, Proceedings*, v. 2, p. 339-345.
- Cuffey, R. J., and Kamandulis, M. A., 1985, Trepostome bryozoan reef-mounds in the upper Ordovician near Maysville, Kentucky (abs.): *Geological Society of America Abstracts with Programs*, v. 17, no. 5, p. 283.
- Cuffey, R. J., and Pursell, B. F., 1995, A possible causal explanation for the Maysville bryozoan reefs (Upper Ordovician, north-central Kentucky) (abs.): *Geological Society of America Abstracts with Programs*, v. 27, no. 3, p. 45.
- Cummings, E. R., 1908, The stratigraphy and paleontology of the Cincinnati Series of Indiana: *Indiana Department of Geology and Natural Resources, Thirty-second Annual Report for 1907*, p. 605-1189.
- Errett, D. H., and Cuffey, R. J. 1989, Bryozoan mode of occurrence relative to individual beds in the type Cincinnati (Upper Ordovician; Ohio, Indiana, Kentucky) (abs.): *Geological Society of America Abstracts with Programs*, v. 21, no. 4, p. 11.
- Gibbons, A. B., 1968, Geologic map of the Mays Lick quadrangle, Mason County, Kentucky: *U.S. Geological Survey Geologic Quadrangle Map GQ-784*, scale 1:24,000.
- Harris, F. W., and Martin, W. D., 1979, Benthic community development in limestone beds of the Waynesville (upper Dillsboro) Formation (Cincinnatian Series, Upper Ordovician) of southeastern Indiana: *Journal of Sedimentary Petrology*, v. 49, p. 1295-1306.
- Kissling, D. L., and Turonis, L. J., 1977, Bryozoan minibioherms in the Upper Ordovician of north-central Kentucky (abs.): *Geological Society of America Abstracts with Programs*, v. 9, no. 3, p. 284-285.
- Kobluk, D. R., 1980, Upper Ordovician (Richmondian) cavity-dwelling (coelobiontic) organisms from southern Ontario: *Canadian Journal of Earth Sciences*, v. 17, p. 1616-1627.
- Pursell, B. F., and Cuffey, R. J., 1995, Bryozoan species surrounding and below the Maysville bryozoan reefs (Upper Ordovician, north-central Kentucky) (abs.): *Geological Society of America Abstracts with Programs*, v. 27, no. 3, p. 81.
- Schilling, F. A., and Peck, J. H., 1967, Geologic map of the Orangeburg quadrangle, northeastern Kentucky: *U.S. Geological Survey Geologic Quadrangle Map GQ-588*, scale 1:24,000.
- Schmidt, R. G., McFarlan, A. C., Nosow, Edmund, Bowman, R. S., and Alberts, Robert, 1961, Examination of Ordovician through Devonian stratigraphy and the Serpent Mound chaotic structure area, in *Guidebook for Field Trips, Cincinnati meeting*, 1961, *Geological Society of America*, p. 259-293.
- Schumacher, G. A., Swinford, E. M., and Shrake, D. L., 1991, Lithostratigraphy of the Grant Lake Limestone and Grant Lake Formation (Upper Ordovician) in southwestern Ohio: *Ohio Journal of Science*, v. 91, p. 56-68.
- Utgaard, J. E., and Perry, T. G., 1964, Trepostomatous bryozoan fauna of the upper part of the Whitewater Formation (Cincinnati) of eastern Indiana and western Ohio: *Indiana Geological Survey Bulletin* 33, 111 p.

Address of author: Roger J. Cuffey, Department of Geosciences (Deike Building), Pennsylvania State University, University Park, PA 16802.

6. THE MADISON/U.S. ROUTE 421 ROAD CUTS—BELLEVUE THROUGH UPPER WHITEWATER STRATA (UPPER ORDOVICIAN, SOUTHEASTERN INDIANA)

by
Helen B. Hay, Stanley M. Totten, and Roger J. Cuffey

SIGNIFICANCE

The Ordovician outcrops at Madison, Indiana, are among the classic Cincinnati fossil-collecting localities of the Ohio-Kentucky-Indiana tri-state area. "In addition, the distinctive dolomitic and sparsely fossiliferous Saluda Formation near the top of the Ordovician is well exposed and reaches its maximum thickness in the Madison area." (Totten and Hay, 1987, p. 365). Relatively recent road cuts along U.S. Route 421, north from the east side of Madison, expose the upper part of the Cincinnati and provide significant insight into the stratigraphy of those units. The outcrops here can be compared instructively to those seen along Indiana Route 56 a few miles to the southwest (Hattin and Cuffey, paper 14 this volume). The Ordovician-Silurian contact, overlain by about 7 meters (23 ft) of Silurian rock, is exposed in the northernmost outcrop along U.S. 421.

LOCATION

This locality consists of three road cuts along U.S. Route 421, at various distances south from its junction with Indiana Route 62 and over 3 miles (4.8 km) north of historic downtown Madison, Jefferson County, Indiana (fig. 6-1). These localities were referred to as the north, middle, and south road cuts, respectively, by Totten and Hay (1987) and given letter designations in their locality map (Totten and Hay, 1987, fig. 1):

- Madison North road cut (= their A), 0.2-0.7 mile (0.3-1.1 km) south of intersection;
- Madison Middle road cut (= their B), 1.1-1.2 miles (1.8-1.9 km) south of intersection; and
- Madison South road cut (= their C), 2.8-2.9 miles (4.5-4.6 km) south of intersection.

The massive Saluda Dolomite is nicely exposed in the north cut 0.5 mile (0.8 km) south of that junction. The north road cut (IN-JE-0001) occupies the SW corner sec. 13 and NW¹/₄NW¹/₄ and center NW¹/₄ sec. 24; the middle road cut (IN-JE-0002) is in the center SW¹/₄ sec. 24; and the south road cut (IN-JE-0003) is in the center W¹/₂NE¹/₄ sec. 35; all are in T. 4 N., R. 10 E., the first two on the Canaan, Indiana, 7.5-minute quadrangle, the third on the Madison West, Indiana-Kentucky, 7.5-minute quadrangle. The Saluda exposure is located at 38°46'45"N, 85°21'58"W (4293308 m N, 641928 m E, UTM zone 16).

STRATIGRAPHY

The Madison/U.S. Route 421 road cuts have been described by Totten and Hay (1987). The following discussion of the stratigraphy is only slightly altered from that work.

Cincinnati rock-stratigraphic classification for southeastern Indiana and southwestern Ohio has undergone substantial revision during the past 30 years. Hay (1981) and Hay and others (1981) extended the Fairview, Miamitown, and Bellevue formation names throughout the outcrop area




FIGURE 6-1.—Location of the U.S. Route 421 road cuts (N, M, S = north, middle, and south cuts, respectively). The three N arrows indicate (from north to south) the northern end of the north cut, the Saluda outcrop, and the southern end of the north cut. Topographic map is from the Canaan, Indiana, 7.5-minute quadrangle.

of Indiana and Ohio and named a new unit, the "Brookville formation," for strata between the top of the Bellevue and the base of the Saluda or Whitewater Formation. The "Brookville formation," equivalent to the upper part of the Dillsboro, is divided into five members, three of which occur in the Madison area. Hay (fig. 17-3 in paper 17 of this volume) compares these new units with previous rock-stratigraphic classifications.

All of the Cincinnatian Series of Ohio and Indiana, with the exception of the Saluda, consists of interbedded thin- to medium-bedded fossiliferous limestones and thin to thick calcareous shales. The limestones include a variety of types (Martin, 1975), but the most common is biomicrosparrudite. Lithofacies are distinguished by percentages of shale, bedding characteristics, textures and structures of the limestones, and character of the shales, including the presence or absence of conspicuous lenses and/or nodules of limestone in the shales (referred to as limy shales).

Six lithofacies types occur in the Madison road cuts: 1a, 2a, 2b, 3a, 3b, and 3c (fig. 6-2). See Hay (paper 17 in this volume) for fuller discussion of these lithofacies, which are summarized here in figure 6-3. The numbers 1, 2, and 3 of the first part of the facies code refer to shale percentage: 1 = greater than 70 percent shale; 2 = 55 to 70 percent; and 3 = less than 55 percent. The second part of the facies code refers to other stratigraphic and lithologic aspects: the letter "a" indicates rather even-bedded limestones and shales without limestone lenses and/or nodules; "b," well-bedded limestones interbedded with limy shales, although some of the limestone beds of these facies may be argillaceous and rubbly; and "c," poorly bedded, rubbly, argillaceous limestones and very limy shales such that a given bed may be called either argillaceous limestone or limy shale.

Cincinnatian strata are exposed in each of the three road cuts along U.S. Route 421 north of Madison (fig. 6-2). The southernmost section, 0.5 mile (0.8 km) north of Madison, includes the upper 7.6 ft (2.3 meters) of the Bellevue Limestone and about 99 ft (30 meters) of the "Excello member" of the "Brookville formation." Toward the north is a small road cut which exposes 16 ft (4.9 meters) of the "Excello member." This section overlaps the top of the south road cut and is close to the top of the "Excello member." The Waynesville Shale Member lies in a covered stratigraphic interval of about 60 ft (18.3 meters) between the top of the middle road cut and the base of the long road cut just to the north. The third and northernmost road cut exposes the Liberty Member of the "Brookville formation," the Saluda and Whitewater Formations, and the Silurian formations.

The Cincinnatian fauna includes abundant and diverse representatives of many phyla. The major rock formers are bryozoans, brachiopods, echinoderms, and mollusks. Trilobites are common, and corals occur in the upper part of the section. Ostracodes are abundant in parts of the Saluda. An excellent and inexpensive reference for identification of these and many other Ordovician fossils is available (Davis, 1992).

BELLEVUE LIMESTONE

The Bellevue Limestone is approximately 40 ft (12 meters) thick in the Madison area, but only the top 7.6 ft (2.3 meters) are exposed in the south road cut, where it is assigned to rubbly facies 3c (fig. 6-2). Notable fossils include the brachiopods *Platystrophia ponderosa*, *P. laticosta*, *Hebertella sinuata*, *Plectorthis*, and *Rafinesquina*; *Rafinesquina* occurs throughout the Cincinnatian. Massive and encrusting bryo-




FIGURE 6-2.—Composite section representing road cuts along U.S. Route 421 north of Madison, Indiana (from Totten and Hay, 1987, p. 368, fig. 3, with permission).

zoans and *Parvohallopora*, a genus of dendritic bryozoans, are major faunal components.

The Bellevue was probably deposited in very shallow, slightly agitated, normal-marine water. The fine-grained terrigenous clastics were not winnowed out, and the en-

Principal diagnostic characteristics of lithofacies	
Group 1 >70% shale	1a Well-bedded, mostly thin limestones and medium to thick, fissile to blocky shales. Includes massive units of 100% shale. Low-diversity fauna.
	1b Rubbly, shaly, nodular limestones; medium to thick shales with limy stringers and layers of fossils in shale matrix. May have some well-bedded limestones as in 1a. Low-diversity fauna. Most easily seen in cores.
Group 2 55-70% shale	2a Like 1a except has high diversity of fauna. Intermediate in shale percentage between 1a and 3a.
	2b Well-bedded limestones, rubbly nodular limestones, and limy shales. Allochems tend to be more finely broken than 2a. High-diversity fauna. Intermediate between 3a and 3c.
Group 3 <55% shale	3a Well-bedded, thin to medium limestones and thin to medium, fissile to blocky shales. High-diversity fauna.
	3b Well-bedded, thin to medium limestones with or without rubbly or nodular shales. High-diversity fauna. Intermediate between 3a and 3c.
	3c Thin-bedded, rubbly weathering, argillaceous limestones, very thin limy shales. High-diversity fauna.
	3d Limestones and shales as in 3a except >20% of limestones are fine-grained laminated burrowed, barren of large fossils; trails on tops. Grade to siltstones.
Group 4 Miscellaneous	4a High-angle cross-bedded calcarenites, bimodal dip. Little or no shale.
	4b Granule-size fossils in thick to massive beds; little or no shale. May be like 4a if seen in similar outcrops.
	4c Massive <i>Loxoplocus</i> limestone bed. Marble Hill Bed.
	4d Bioturbated light-gray limestone with low-diversity fauna of ostracodes, algae, snails, and <i>Tetradium</i> . Related to Saluda facies complex.




FIGURE 6-3.—Lithofacies characteristics (see Hay, paper 17 in this volume).

environment, therefore, was not likely to have been one of extremely high energy.

“EXCELLO MEMBER” OF THE “BROOKVILLE FORMATION”

The “Excello member” constitutes the remainder of the south road cut and the middle road cut. As figure 6-2 indicates, the “Excello” is composed of intervals of facies 1a, 2a, and 3a. Most of the outcrop by far is classified as facies 2a but could be more finely subdivided into smaller intervals of 3a alternating with 1a. Compared with the Bellevue, the limestones are generally thicker and more evenly bedded, and the shales are fissile or blocky, not limy. The south road cut has four levels separated by three benches. *Platystrophia ponderosa* and *P. laticosta* are found as high as the second level but not higher. Other brachiopods are *Rafinesquina*, *Zygospira*, and *Plectorthis*; *Leptaena* occurs in level three. The “Excello” has abundant and diverse mollusks. *Ctenodonta*, a small clam, is particularly abundant at the top of the middle road cut. Large, coarse-ribbed clams, gastropods, including *Cyclonema inflatum*, and cephalopods

also occur. Echinoderms, bryozoans, and trilobites occur in the “Excello.”

The “Excello” sediments indicate a transgressive phase with sediments deposited in slightly deeper water than the Bellevue, probably below normal wave base but subject to periodic reworking by storm waves and strong currents that accounted for the interbedding of limestones and shales. Martin (1975) and Harris and Martin (1979) believed that the limestones are in situ biogenic accretions representing progressive community development after colonization of the muddy substrate.

LIBERTY MEMBER OF THE “BROOKVILLE FORMATION”

The lower half of the Liberty Member (fig. 6-2), exposed at the base of the north road cut, consists of several facies intervals (2b, 3c, 2b, 3b upward), which vary in shale percentage but share the characteristic of limy shales. Except for a generally higher percentage of shale, these intervals are similar to the facies of the Bellevue and probably represent a similar depositional environment. The upper half

of the Liberty Member has more limestone than the lower half and has nonlimy shales. These upper beds were probably deposited in a higher energy environment close to normal wave base. The Liberty-Saluda sequence indicates shoaling of the sea, a regressive phase. Prominent Liberty brachiopods include *Thaerodonta*, *Plaesiomys*, *Hebertella*, *Rafinesquina*, *Catazyga*, *Strophomena*, and *Hiscobeccus*. The horn coral *Grevingkia* is present.

SALUDA FORMATION

The Saluda contains a number of facies but is distinguished from the other formations by the preponderance of fine-grained dolomite and dolomitic mudstones, having lesser amounts of dolomitic limestone in the upper and lower parts. The base consists of 7.5 ft (2.3 meters) of slabby dolomitic mudstone that appears quite massive in fresh cuts. This lithology is overlain by interbedded limestone, dolomite, and mudstone containing the large colonial corals *Favistina* and *Tetradium*. The thickest unit of the Saluda is mud-cracked and ripple-marked, argillaceous and silty dolomite that exhibits color mottling in its lower part. This laminated, rather homogeneous rock appears massive in the fresh road cut. The total thickness of the Saluda is 51 ft (15.5 meters).

The main body of the Saluda is nearly barren of fossils, but some beds and bedding planes bear a normal-marine fauna that perhaps was washed into the area or that may indicate times of normal-marine salinity. Ostracodes and burrows are the dominant fossils in the lower and upper dolomitic and calcareous zones. Hatfield (1968) interpreted the paleogeographic setting of the Saluda as a hypersaline lagoon rimmed by colonial coral reefs, a couple of which are exposed in these road cuts (Larabee, 1994). The mud-cracked sediment was intermittently exposed to the atmosphere. Regionally, it is a lens-shaped body of rock that pinches out northward between Brookville and Richmond, Indiana (Hay, 1981). The Saluda is time equivalent to the mid-Whitewater strata to the east at Caesar Creek, where the bryozoans are highly diverse (Rockwell and Cuffey, 1996; Schumacher and others, paper 13 of this volume) and thus contrast with the greatly reduced bryozoan fauna here and associated with the penesaline Saluda shelf lagoon (Butler and Cuffey, 1994, 1996; Cuffey, Butler, and Rockwell, 1996).

UPPER? WHITEWATER FORMATION

The Whitewater Formation is 12.5 ft (3.8 meters) thick at Madison, if, as we consider it here, the base is taken to be at the change from Saluda dolomite to dolomitic limestone. If the *Lophospira hammelli* bed is taken as the base of the Whitewater (Foerste, 1903), the Saluda is 59.4 ft (18 meters) thick and the Whitewater is 4.6 ft (1.4 meters) thick at Madison. The Whitewater lithology farther north in Indiana is typically that of rubbly facies 3c, but at the road cut at Madison it consists, in descending order, of the following six units (modified from Conklin, 1977) that do not fit the facies classification used for other Ordovician strata: (1) fine-grained mottled dolomite (1.8 ft/0.56 meter), (2) black shale (2.56 inches/6.4 cm), (3) fossiliferous micritic limestone with the gastropod *Lophospira hammelli* (2.9 ft/0.87 meter), (4) black shale (0.2 inch/0.4 cm), (5) silty dolomitic biomicrosparite with ostracodes (9 inches/22.9 cm), and (6) ostracode-bearing, coarse-grained, greenish-gray, burrowed dolomitic limestone (8 ft/2.44 meters) placed by some authors at the top of the Saluda rather than in the Whitewater.

The Whitewater represents a return to normal-marine conditions, although the origin of the fine-grained dolomite at the top is uncertain.

REFERENCES CITED

- Butler, K. L., and Cuffey, R. J., 1994, Bryodiversity and paleostress in the Saluda dolomitic shelf lagoon (latest Ordovician, southeastern Indiana) (abs.): Geological Society of America Abstracts with Programs, v. 26, no. 5, p. 7.
- _____, 1996, Reduced bryozoan diversity and paleoenvironmental stress in the Saluda Dolomite (uppermost Ordovician, southeastern Indiana), in Gordon, D. P., Smith, A. M., and Grant-Mackie, J. A., eds., Bryozoans in space and time: Wellington, New Zealand, Proceedings of the 10th International Bryozoology Conference (1995), National Institute of Water & Atmospheric Research, Ltd., p. 55-61.
- Conklin, B. J., 1977, Stratigraphy and petrography of the Madison, Indiana, U.S. 421 road cut, Upper Ordovician through Middle Silurian: B.S. thesis (unpub.), Earlham College, 75 p.
- Cuffey, R. J., Butler, K. L., and Rockwell, L. A., 1996, Bryozoan diversity and salinity in the uppermost Ordovician (Saluda and Whitewater Formations) across southeastern Indiana and southwestern Ohio (abs.): Geological Society of America Abstracts with Programs, v. 28, no. 6, p. 34-35.
- Davis, R. A., ed., 1992, Cincinnati fossils; an elementary guide to the Ordovician rocks and fossils of the Cincinnati, Ohio, region: Cincinnati Museum of Natural History, 61 p.
- Foerste, A. F., 1903, The Richmond Group along the western side of the Cincinnati Anticline in Indiana and Kentucky: American Geologist, v. 31, p. 333-361.
- Harris, F. W., and Martin, W. D., 1979, Benthic community development in limestone beds of the Waynesville (upper Dillsboro) Formation (Cincinnatian Series, Upper Ordovician) of southeastern Indiana: Journal of Sedimentary Petrology, v. 49, p. 1295-1306.
- Hatfield, C. B., 1968, Stratigraphy and paleoecology of the Saluda Formation (Cincinnatian) in Indiana, Ohio, and Kentucky: Geological Society of America Special Paper 95, 34 p.
- Hay, H. B., 1981, Lithofacies and formations of the Cincinnatian Series (Upper Ordovician), southeastern Indiana and southwestern Ohio: Ph.D. dissertation (unpub.), Miami University, 236 p.
- Hay, H. B., Pope, J. K., and Frey, R. C., 1981, Lithostratigraphy, cyclic sedimentation, and paleoecology of the Cincinnatian Series in southwestern Ohio and southeastern Indiana, in Roberts, T. G., ed., GSA Cincinnati '81 Field Trip Guidebooks: American Geological Institute, v. 1, p. 73-86.
- Larabee, J. H., 1994, Stratigraphy and paleoecology of coralline biostromes and patch reefs, Upper Ordovician of southeastern Indiana (abs.): Geological Society of America Abstracts with Programs, v. 26, no. 5, p. 49.
- Martin, W. D., 1975, The petrology of a composite vertical section of Cincinnatian Series limestones (Upper Ordovician) of southwestern Ohio, southeastern Indiana, and northern Kentucky: Journal of Sedimentology Petrology, v. 45, p. 907-925.
- Rockwell, L. A., and Cuffey, R. J., 1996, Great bryozoan diversity and sampling in the uppermost Ordovician (Whitewater Formation) at Caesar Creek (southwestern Ohio) (abs.): Geological Society of America Abstracts with Programs, v. 28, no. 6, p. 62.
- Totten, S. M., and Hay, H. B., 1987, Madison, Indiana: geomorphology, and Paleozoic and Quaternary geology, in Biggs, D. L., ed., North-Central Section of the Geological Society of America: Geological Society of America Centennial Field Guide, v. 3, p. 365-370.
- Addresses of authors:* Helen B. Hay, deceased, formerly Geology Department, Earlham College, Richmond, IN 47374; Stanley M. Totten, Department of Geology, Hanover College, Hanover, IN 47243; and Roger J. Cuffey, Department of Geosciences (Deike Bldg.), Pennsylvania State University, University Park, PA 16802.

7. THE MIAMITOWN SHALE: STRATIGRAPHIC AND HISTORIC CONTEXT (UPPER ORDOVICIAN, CINCINNATI, OHIO, REGION)

by
Benjamin F. Dattilo

INTRODUCTION

In 1958, Gutstadt observed that the original stratigraphic nomenclature of the Cincinnati rocks (for example, Nickles, 1902) was defined by a mixture of paleontologic and lithologic criteria. He concluded that it was necessary to separate lithostratigraphic and chronostratigraphic terminology for these strata. This sentiment was an important factor in the development of new lithostratigraphic units to replace the old, largely biostratigraphic units.

More recent studies, however, indicated the possibility that eustatically induced lithologic cyclicity is present in the Cincinnati (Tobin, 1982; Jennette, 1986). In other words, lithologic characteristics can be valid criteria in the definition of chronostratigraphic units. From this realization it can be concluded that an integrated lithologic-paleontologic approach is potentially a powerful tool in the delineation of fine-scale chronostratigraphic correlations.

I began research on the Miamitown Shale with such an integrated study of cyclicity in mind (Dattilo, 1996). Having completed a number of fine-scale, lithology-based correlations across the Miamitown-Fairview facies transition, I discovered that a number of thin units already had been recognized within the same stratigraphic interval in the immediate vicinity of Cincinnati. One author (Hyde, 1959) also had traced one of these thin units into the area that later would be designated the type area of the Miamitown Shale (Ford, 1967). Whereas Hyde's (1959) correlations apparently had been intended as chronostratigraphic, and because they were roughly comparable to my own, I began using the old stratigraphic names in my own correlations. This paper presents a documentation of this informal chronostratigraphic nomenclature as seen in five outcrop sections, three of which will be discussed in detail. In no way are the designations of the units intended as formal stratigraphic names; rather, they denote testable hypotheses.

UNIT OVERVIEW

The Miamitown Shale is a thin, mollusk-dominated unit between the Fairview Formation and the Bellevue Member of the Grant Lake Limestone, both of which are considerably more limestone rich and are brachiopod dominated. The interval is intriguing because of this faunal-lithologic contrast and because of the relatively complex geometry of the rapid Miamitown-Fairview facies transition. In the 16 km (10 miles) between Miamitown and Cincinnati (fig. 7-1), the Miamitown Shale thins from about 5 meters (16 ft) to less than 1 meter (3 ft).

STRATIGRAPHIC SEQUENCE AND NOMENCLATURE

A brief history of Cincinnati fine-scale stratigraphic nomenclature will acquaint the reader with the complexities of the Miamitown interval in Cincinnati. The Miamitown was formally named by Ford in 1967, but a perusal of largely unpublished manuscripts and theses reveals that the gastropod-rich shale has been recognized since the 1930's and perhaps longer. Figure 7-2 compares the finer scale




FIGURE 7-1.—Map showing locations of the Miamitown outcrops discussed in this paper and shown on the stratigraphic cross section through these outcrops (fig. 7-4). **CHS** = Rice and Gage Streets/Christ Hospital (OH-HA-0004), **MAF** = Mt. Airy Forest (OH-HA-0039), **MTN** = Miamitown West (OH-HA-0008), **RDR** = Riedlin Road/Mason Road (KY-KE-0001), and **WYN** = Wayne Road (KY-KE-0006).

classification by Desjardins, as reported by Forsyth (1946) and published by Hyde (1959), with the more formally published classifications of Nickles (1902) and of Ford (1967). An unpublished U.S. Geological Survey manuscript by Ulrich and Bassler (1914) also contains an interesting reference to a shaly interval between the Fairmount and the Bellevue, the "*Stigmatella irregularis* Zone."

Desjardins (1933, 1934, and 1935) studied the physiography of the Cincinnati area, but also made detailed notes on the bedrock stratigraphy of the area (see Forsyth, 1946). Unfortunately, the only Desjardins stratigraphic column yet located is the generalized one published in Bucher, Caster, and Jones (1939, 1945) (reproduced in part in fig. 7-3). In this column, what is now known as the Miamitown Shale is clearly delineated just above the "*fracta* Zone" in the lowest part of the Bellevue Member. A second shaly horizon, Hyde's "B," is also visible in this rather coarse overview.

Forsyth (1946) described these units in more detail. At the top of the Fairmount Member there are three "*fracta*" or "shingled" zones, each 0.7 meter (2 ft) thick; each successive pair is separated by approximately 1.5 meters (5 ft) of more shaly, even-bedded strata. These zones are characterized by abundant fragmented valves of *Rafinesquina alternata*, *Hebertella occidentalis*, and *Platystrophia*. The uppermost ("first '*fracta*' zone" or "upper shingled") is the most extensive and continuous, whereas the lowest ("third '*fracta*' zone" or "lower shingled") is the least continuous.

Above the shingled zones is a shaly interval 2 meters (5 to 7 ft) thick with a few gastropod-rich limestones called the "gastropod zone" or "gastropod shale" or "zone of abundant gastropods and pelecypods" (Forsyth, 1946). This zone is capped by the 2- to 3-meter (7- to 8-ft) "lower very massive shelly/rubby" or "lower massive" horizon, which consists mostly of limestone and contrasts markedly with the underlying units. The base of the Bellevue was drawn by Des-

Nickles, 1902	Ulrich and Bassler, 1914	Desjardins <i>vide</i> Forsyth, 1946		Hyde, 1959	Ford, 1967	This paper (informal)		
Corryville	Corryville	Corryville Member		Undesignated	Unnamed	Corryville		
Bellevue Limestone	Bellevue Limestone	Bellevue Limestone	upper massive shelly/rubbly horizon	Bellevue Limestone	Bellevue Limestone	upper massive		
			5'-8'			"A"	Bellevue Limestone	<i>Rafinesquina</i> shale
			medium light-buff-yellow shaly horizon w/ large full <i>Rafinesquina alternata</i>			5'		"B"
?	?	lower very massive shelly/rubbly horizon	7'-8'	"C"				
Fairmount Member	<i>Stigmatella irregularis</i> Zone	?	gastropod zone light-buff-yellow very shaly horizon w/ many gastropods and pelecypods	Fairmount Member	Fairmount Member	gastropod shale		
			5'-7'			"D"	Miami town Shale	upper shingled
	first "fracta" zone	2'	"E"			middle shale		
	undescibed	5'	undesignated			middle shingled		
	second "fracta" zone	2'	2nd shingled			lower shale		
	undescibed	5'	undesignated			lower shingled		
third "fracta" zone	2'	3rd shingled	SE					

FIGURE 7-2.—History of stratigraphic nomenclature applied to the Miamitown-Bellevue interval.




FIGURE 7-3.—Reproduction of part of Desjardins' generalized section showing his Fairmount, Bellevue, and Corryville units and the clear early recognition of the Miamitown Shale or gastropod shale as a marker horizon. From Bucher, Caster, and Jones (1939, 1945).

jardins at the base of the first shingled zone, because all of the shingled zones were recognized as being fundamentally Bellevue-like. Nickles (1902), however, placed the base of the Bellevue above all three “*fracta*” or “shingled” zones, at the bottom of the “lower very massive shelly/rubbly horizon,” as did Hyde (1959) and Ford (1967).

Above the “lower very massive shelly/rubbly horizon” is a shaly horizon 3 meters (5 ft) thick characterized by large, “full” *Rafinesquina alternata*. Overlying this *Rafinesquina*-bearing shale is an “upper massive shelly/rubbly horizon,” the top of which is recognized as the top of the Bellevue.

The stratigraphy of Ford (1967) in this interval differs from that of Nickles (1902) in the important recognition of the Miami town Shale as a lithofacies that thickens to the northwest.

Instead of numbers or letters, a modified Desjardins nomenclature is used here to denote a set of nearly isochronous units which can be traced between outcrops. These modified, informal designations are listed in the right-hand column of figure 7-2; they are intended as convenient names to be used in discussing stratigraphic relationships. They are in no way intended as formal stratigraphic units.

CORRELATIONS

Figure 7-4 is a correlation chart of three of the outcrops

discussed in this paper: Rice and Gage Streets (OH-HA-0004, CHS in figure), Mt. Airy Forest (OH-HA-0039, MAF), and Miami town West (OH-HA-0008, MTN). Two additional outcrops, Wayne Road (KY-KE-0006, WYN) and Riedlin Road/Mason Road (KY-KE-0001, RDR) also are included. The Riedlin Road/Mason Road site is discussed by Diekmeyer in paper 3 in this volume. These correlations are based primarily on the matching of shale-percentage curves produced by a 1-meter running average of bed-by-bed lithologic data, a method similar to that used by Gray (1972). Details of correlation are corroborated by paleontological data, particularly the stratigraphically limited occurrence of *Heterorthina fairmountensis*, used as a datum in this diagram.

Note that these curves have similar shapes but that the actual percentage of shale at Miami town West is lower than that at Rice and Gage Streets. In other words, the patterns correlate across facies. The stippled patterns show this clearly; the denser pattern (70 to 100 percent shale) corresponds approximately to the Miami town Shale facies, which at Miami town West encompasses the interval between the lower shingled unit and the lower massive unit. At Rice and Gage Streets, the Miami town facies is limited to the gastropod shale.

The shingled intervals do not display the “shingled facies” everywhere; in some places a given shingled unit may




FIGURE 7-4.—Detailed stratigraphic cross section of the Fairview-Miami town-Bellevue interval from Miami town West through Cincinnati to Riedlin Road/Mason Road. This diagram shows how shale-percentage curves are used to make high-resolution stratigraphic correlations through a facies transition. The actual position of each section is represented by the left edge of each shale-percentage log. Superimposed on the facies diagram is the informal stratigraphic nomenclature used in this paper. Datum is the stratigraphically restricted *Heterorthina fairmountensis* occurrence. CHS = Rice and Gage Streets/Christ Hospital (OH-HA-0004), MAF = Mt. Airy Forest (OH-HA-0039), MTN = Miami town West (OH-HA-0008), RDR = Riedlin Road/Mason Road (KY-KE-0001), and WYN = Wayne Road (KY-KE-0006). See figure 7-1 for locations of sections.

contain only tabular limestones without edgewise-stacked *Rafinesquina*, fossiliferous shale, or wavy-bedded limestone that originally made the units noteworthy. This fact does not detract from their usefulness as stratigraphic markers; the upper-shingled-unit “spike” is visible at Miamitown West even though the actual limestones are thin and relatively scattered, and even though no coquina of edgewise *Rafinesquina* is present. The lateral consistency of the shale-percentage pattern probably resulted from rapid and widespread sea-level fluctuations. Each shingled unit, when combined with the underlying shale, may constitute a “cycle” (Tobin, 1982).

DEPOSITIONAL ENVIRONMENT

The stratigraphic position and unusual fauna of the Miamitown Shale make it an interesting study. Molluscan faunas in the Ordovician of the Appalachian Basin have been thought of as indicating relatively shallow water, and the Miamitown Shale occurs just below the Bellevue, which contains clear indicators of shallow-water deposition such as fragmented, abraded shells and robust specimens. On the other hand, the Miamitown is a fine-grained facies that must have been deposited in relatively calm water. Further work with benthic macrofossils may help in interpretation.

A regional stratigraphic cross section (fig. 7-5) illustrates how the Miamitown is a small part of a system of shales from the northwest intertonguing with limestones from the southeast. A simple interpretation of this pattern is that limestone-rich intervals represent shallower water, and shale-rich intervals represent deeper water; the intertonguing resulted from fluctuations in sea level. Notice that the upper part of the Bellevue reaches the farthest to the northwest, indicating that it is the top of a major shoaling-upward cycle. The Miamitown, positioned as it is toward the top of this cycle, is probably not comparable directly to the Kope Formation below. The shaly composition is the product of the




FIGURE 7-5.—Regional stratigraphic cross section of the Kope-Bellevue interval from Wells County, Indiana, to Hamilton County, Ohio. This diagram shows the major facies transitions as indicated by subsurface data.

rate of sea-level change rather than a reflection of absolute depth; it was probably deposited in shallower water than was most of the Kope.

CINCINNATI OUTCROPS: BELLEVUE HILL, EMMING STREET, AND RICE AND GAGE STREETS

SIGNIFICANCE

Outcrops around downtown Cincinnati have been more accessible to study historically than other outcrops; hence, a number of them are type sections. Three outcrops are of particular interest with respect to the Miamitown interval: (1) Bellevue Hill (OH-HA-0003), (2) Emming Street (OH-HA-0002), and (3) Rice and Gage Streets (OH-HA-0004) (fig. 7-6).

Bellevue Hill is one of the longest standing outcrops from which the Cincinnati section has been studied. It is the original type section of the Bellevue Limestone (Nickles, 1902), the neotype section of the Fairview Formation (Ford, 1967), and probably formed a basis for Desjardins’ generalized stratigraphic section. In addition to this, numerous studies have been made on its fauna and lithologic sequence.

Emming Street, located only a few blocks from Bellevue Hill, is more safely accessible for a close-up look. There are good exposures of the Miamitown/gastropod shale and the Bellevue Limestone, including the upper contact with the Corryville Formation.

Rice and Gage Streets is the neotype section of the Bellevue Limestone (Ford, 1967) and lies very close to the original type section on Bellevue Hill. It exposes the Fairview, Miamitown, and Bellevue units.




FIGURE 7-6.—Locations of outcrops in Cincinnati proper. Covington, Kentucky–Ohio, 7.5-minute quadrangle. BH = Bellevue Hill (OH-HA-0003), EMG = Emming Street (OH-HA-0002), and CHS = Rice and Gage Streets/Christ Hospital (OH-HA-0004).

LOCATIONS

The Bellevue Hill site (OH-HA-0003) lies just under Bellevue Hill Park to the east of Clifton Avenue, in sec. 13, City of Cincinnati, Hamilton County, Ohio, on the Covington, Kentucky–Ohio, 7.5-minute quadrangle (fig. 7-6). Parking and traffic are problems with all of the Cincinnati outcrops. It probably is unwise to visit Bellevue Hill from Clifton Avenue; fencing has been installed to prevent rock slabs from falling into Clifton Avenue; unfortunately, rocks dislodged by a climbing geologist commonly leap the fence and end up in the street, thereby endangering passing vehicles. If approached from Bellevue Hill Park, the Miamitown/gastropod shale through Corryville can be examined easily.

The Emming Street site (OH-HA-0002) is an old quarry face on the north side of Emming Street, in sec. 19, City of Cincinnati, Hamilton County, Ohio, on the Covington, Kentucky–Ohio, 7.5-minute quadrangle (fig. 7-6). Emming Street, only a few blocks from Bellevue Hill, is much more accessible. At the present time the exposure is limited to the interval between the upper shingled zone and the Bellevue-Corryville contact.

The Rice and Gage Streets site (OH-HA-0004) (fig. 7-7) is at the intersection of Rice and Gage Streets, on the north side of Gage Street, under the Christ Hospital heliport, in sec. 13, City of Cincinnati, Hamilton County, Ohio, on the Covington, Kentucky–Ohio, 7.5-minute quadrangle, 4332909 m N, 714976 m E, UTM zone 16 (fig. 7-6). This locality is the private property of Christ Hospital, but can be viewed easily from the streets themselves. If more intensive study or sampling is required, contact the Public Relations Department of Christ Hospital for permission.

LITHOLOGIC AND FAUNAL SEQUENCE

These Cincinnati sections, all rather close together, differ little from each other or from Desjardins' summary sections. Figure 7-8 illustrates some specific lithologic and faunal occurrences. All three shingled zones display a "coquinoid" facies consisting of shale with fragments of the brachiopods *Rafinesquina alternata*, *Hebertella occidentalis*, *Platystrophia* sp., and bryozoans. The fauna of the intervening shales is sparse, but the middle shale contains a molluscan fauna




FIGURE 7-7.—Photograph of Rice and Gage Streets locality, showing boundaries of stratigraphic units. ls = lower shingled; ms = middle shingled; us = upper shingled; lm = lower massive; Rs = *Rafinesquina* shale; um = upper massive.

consisting largely of the gastropod *Loxoplocus bowdeni*, the pelecypod *Modiolopsis*, and abundant brachiopods of the genus *Zygospira*. The gastropod shale contains nodular siltstones in which lingulid brachiopods are common. Near the top of this unit is the characteristic gastropod packstone (again, mostly *Loxoplocus bowdeni*).

The lower massive unit is particularly limestone rich; it contains wavy limestone beds 5 to 15 cm (2 to 6 inches) thick interbedded with subequal amounts of highly fossiliferous shale. Fossils are dominated by bryozoans, some colonies of which are "massive" or round and up to 15 cm (6 inches) across. *Platystrophia ponderosa*, other *Platystrophia*, *Hebertella occidentalis*, and *Rafinesquina alternata* also are common. The *Rafinesquina* shale contains large *Rafinesquina* concentrated in one or two beds. The upper massive unit contains a great abundance of large *Rafinesquina alternata* in highly fossiliferous shale with a few wavy-bedded limestones.

MT. AIRY FOREST INTERSTATE CUT

SIGNIFICANCE

This series of road cuts (OH-HA-0039) exposes Kope through Corryville strata. It is one of the largest and most continuous exposures of this interval in the area.

LOCATION

The Mt. Airy Forest site (OH-HA-0039) is a series of large road cuts on either side of I-74 between mile 15.6 and mile 16.8 where it passes through Mt. Airy Forest, sec. 4, Green Township, Hamilton County, Ohio, on the Cincinnati West, Ohio, 7.5-minute quadrangle, 438000 m N, 709000 m E, UTM zone 16 (fig. 7-9). Permission from the Ohio Department of Transportation is required for close examination and sampling of this interstate cut. It is also a steep cut (fig. 7-10), and climbing it may be risky both to the geologist and to motorists.

LITHOLOGIC SEQUENCE

This outcrop contains less limestone than the Cincinnati outcrops (compare figs. 7-8 and 7-11). The shingled units in particular are less well developed, and the intervening strata are very shaly. The fauna of these intervals is not markedly different from the fauna at Rice and Gage Streets, but the gastropod and pelecypod fauna of the gastropod shale is more diverse. Beds from this interval may contain *Modiolopsis*, *Ambonychia*, *Platystrophia*, *Zygospira*, as well as the abundant *Loxoplocus bowdeni*. "Float" specimens of small, well-preserved edrioasteroids probably also originate from this unit. The Miamitown Shale lithofacies extends down through the middle shale unit.

The lower massive unit, still in marked contrast to the underlying intervals, also contains less limestone than it does in Cincinnati. The *Rafinesquina* shale contains pelecypods at this locality. The upper massive unit probably is the source of large semi-articulated edrioasteroids that have been found in float material.

MIAMITOWN WEST

SIGNIFICANCE

This series of road cuts on the interchange between I-74




FIGURE 7-8.—Measured columnar section of the Rice and Gage Streets site (1:100), showing lithologies (see fig. 9-5 for patterns), Ford (1967) nomenclature, informal detailed stratigraphic nomenclature, and relative abundances of selected fossils, ordered according to occurrence as shown by cluster analysis. l.s. = lower shingled; m.s. = middle shingled; u.s. = upper shingled; g.s. = gastropod shale; mont. encr. = monticulated encrusting bryozoans.




FIGURE 7-9.—Locations of Mt. Airy Forest road cuts (OH-HA-0039). Cincinnati West, Ohio, 7.5-minute quadrangle.

and I-275 west of Miami town provides an expanded exposure of Ford's (1967) original type section for the Miami town Shale, the "Highway cuts west of Miami town." Because I-275 had not been constructed at the time, Ford's observations were limited to the cuts around the original I-74 (still well exposed). Now the northernmost cut, on the interchange ramp from I-74 west to I-275 south, provides an exposure of the upper massive and *Rafinesquina* shale units (fig. 7-12) not visible when Ford did his original work. Similarly, the southernmost cut exposes several additional meters of strata just below the original type section.

LOCATION

The Miami town West site (OH-HA-0008) is located at the interchange between I-74 and I-275 northwest of Miami town, Hamilton County, Ohio. The locality consists of six road cuts surrounding the various roadways of the interchange, in sec. 35, Whitewater Township, Hamilton County, Ohio, on the Addyston, Ohio-Kentucky, 7.5-minute quadrangle, 4343499 m N, 695446 m E, UTM zone 16 (fig. 7-13). As with all interstate highway cuts, permission from the Ohio Department of Transportation is required. Otherwise, all of the road cuts are generally accessible and relatively safe to the careful driver.

LITHOLOGIC AND FAUNAL SEQUENCE

A quick glance at figure 7-14 reveals the remarkable amount of shale in the shingled intervals. The upper shingled unit contains a thick grainstone and some fossiliferous shale, and the middle and lower shingled units are barely distinguishable. The Miami town lithofacies extends down to the top of the middle shingled unit. Faunally, the gastropod shale is still distinctive and contains the same gastropod-*Zygospira*-bivalve assemblage, but gastropods




FIGURE 7-10.—Photograph of Mt. Airy Forest road cut, showing stratigraphic contacts. ls = lower shingled; ms = middle shingled; us = upper shingled; lm = lower massive; Rs = *Rafinesquina* shale; and um = upper massive.

and pelecypods can be found throughout the Miami town lithofacies. The upper shingled interval contains a few unusual fossils; in or around this interval are localized deposits of small *Rafinesquina* (2-3 cm in length) and small edrioasteroids.

The lower massive unit here contains much less limestone than at the previous outcrops, and the fauna is different; *Rafinesquina*, *Zygospira*, and bryozoans (not large colonies) dominate. The *Rafinesquina* shale is relatively unfossiliferous; it contains a few *Rafinesquina* and some mollusk-rich siltstones. The upper massive unit, again less limestone rich than at Cincinnati, is dominated by *Rafinesquina* and contains large, poorly preserved edrioasteroids.

ACKNOWLEDGMENTS

This research was supported in part by the following grants: Cincinnati Museum of Natural History Research Grant 1989, Geological Society of America research grants 4402-90 and 4666-91, Paleontological Society/Margaret C. Wray Trust grant-in-assistance 1991, and Sigma Xi grant in aid of research 1991.

ADDITIONAL LOCALITIES FOR THE MIAMI TOWN SHALE

For details on locations, see Appendix A at the end of this volume.

Ashtree Drive and Hamilton Avenue (OH-HA-0041). Fairview through lower Bellevue exposed. Climbing gear required.

Congress Run (OH-HA-0016). Stream cuts along Congress Run and tributary to Congress Run extending 0.5 mile (0.8 km) north from intersection with Galbraith Road east of Winton Road. Section measured in detail by Ford and Osborne in 1963, Ohio Division of Geological Survey measured section no. 15373.

Crosby Road (OH-HA-0017). Section measured in detail by Ford in 1964, Ohio Division of Geological Survey measured section no. 15379.

Delhi Pike (OH-HA-0040). Contains Fairview and Miami town strata. Section measured in detail by Swinford and Vormelker in 1985, Ohio Division of Geological Survey measured section no. 16879.

Lawrenceburg (KY-BE-0003). Road cuts on either side of I-275 in Boone County, Kentucky, 0.4 mile (0.7 km) south of the Ohio River across from Lawrenceburg, Indiana.




FIGURE 7-11.—Measured columnar section of Mt. Airy Forest strata (1:100), showing lithologies (see fig. 9-5 for patterns), Ford (1967) nomenclature, informal detailed stratigraphic nomenclature, and relative abundances of selected fossils, ordered according to occurrence as shown by cluster analysis. l.s. = lower shingled; m.s. = middle shingled; u.s. = upper shingled; g.s. = gastropod shale; mont. encr. = monticulated encrusting bryozoans.




FIGURE 7-12.—Photograph of Miami town West road cut, showing stratigraphic contacts. ls = lower shingled; ms = middle shingled; us = upper shingled; lm = lower massive; Rs = *Rafinesquina* shale; and um = upper massive.




FIGURE 7-13.—Location of Miami town West road cuts (OH-HA-0008). T = approximate location of the original type section of the Miami town Shale. Addyston, Ohio-Kentucky, 7.5-minute quadrangle).




FIGURE 7-14.—Measured columnar section of strata at the Miami West locality (1:100), showing lithologies (see fig. 9-5 for patterns), Ford (1967) nomenclature, informal detailed stratigraphic nomenclature, and relative abundances of selected fossils, ordered according to occurrence as shown by cluster analysis. I. s. = lower shingled; m. s. = middle shingled; u. s. = upper shingled; g. s. = gastropod shale; mont. encr. = monticulated encrusting bryozoans.

Muddy Creek (OH-HA-0020). Upper Fairview, Miamitown, and Bellevue Formations are exposed in a series of stream cuts along Muddy Creek west of Beech Grove Drive. Sections measured in detail by Ford in 1963, Ohio Division of Geological Survey measured section nos. 15376, 15458, and 15459.

Riedlin Road/Mason Road (KY-KE-0001). See Diekmeyer (paper 3 in this volume).

Sheits Road (OH-HA-0023). Section measured in detail by Ford in 1964, Ohio Division of Geological Survey measured section no. 15380, and by Swinford and Vormelker in 1985, Ohio Division of Geological Survey measured section no. 16872.

Wayne Road (Devou Park) (KY-KE-0006). Exposures of upper Fairview, Miamitown, and lower Bellevue. Included on figure 7-4 cross section.

REFERENCES CITED

- Bucher, W. H., Caster, K. E., and Jones, Stewart, 1939, Elementary description of Cincinnati fossils and strata and plates of commoner fossils in the vicinity of Cincinnati, Ohio: University of Cincinnati, 13 p.
- _____, 1945, Elementary guide to the fossils and strata in the vicinity of Cincinnati: Cincinnati Museum of Natural History, 31 p.
- Dattilo, B. F., 1996, A quantitative paleoecological approach to high-resolution cyclic and event stratigraphy: the Upper Ordovician Miamitown Shale in the type Cincinnati: *Lethaia*, v. 29, p. 21-37.
- Desjardins, L. H., 1933, The pre-Illinoian glaciation of the Cincinnati region (abs.): *Kentucky Academy of Science Transactions*, v. 5, p. 30.
- _____, 1934, The preglacial physiography of the Cincinnati region: M.S. thesis (unpub.), University of Cincinnati, 43 p.
- _____, 1935, Physiography of the Cincinnati region: *Compass*, v. 15, p. 147-151.
- Ford, J. P., 1967, Cincinnati geology in southwest Hamilton County, Ohio: *American Association of Petroleum Geologists Bulletin*, v. 51, p. 918-936.
- Forsyth, J. L., 1946, The Eden and Maysville groups of the Cincinnati series at Cincinnati, Ohio: M.S. thesis (unpub.), University of Cincinnati, 122 p.
- Gray, H. H., 1972, Lithostratigraphy of the Maquoketa Group (Ordovician) in Indiana: *Indiana Geological Survey Special Report 7*, 31 p.
- Gutstadt, A. M., 1958, Upper Ordovician stratigraphy in eastern interior region: *American Association of Petroleum Geologists Bulletin*, v. 42, p. 513-547.
- Hyde, D. E., 1959, A structural and stratigraphic study of the Fairview-McMillan formational contact in the Cincinnati area: *Compass*, v. 36, p. 161-171.
- Jennette, D. C., 1986, Storm-dominated cyclic ramp deposits of the Kope-Fairview transition (Upper Ordovician), southwestern Ohio and northern Kentucky: M.S. thesis (unpub.), University of Cincinnati, 210 p.
- Nickles, J. M., 1902, The geology of Cincinnati: *Cincinnati Society of Natural History Journal*, v. 20, p. 49-100; reprint edition, p. 1-53.
- Tobin, R. C., 1982, A model for cyclic deposition in the Cincinnati Series of southwestern Ohio, northern Kentucky, and southeastern Indiana: Ph.D. dissertation, University of Cincinnati, 483 p.
- Ulrich, E. O., and Bassler, R. S., 1914, Report on the stratigraphy of the Cincinnati, Ohio, quadrangle: U.S. Geological Survey open-file report, 122 p. [Incomplete manuscript of text for the never-published Cincinnati Folio, apparently completed to its present state early in 1914; there is a copy made in 1951 in the Geology/Physics Library at the University of Cincinnati.]

Address of author: Benjamin F. Dattilo, Geoscience Department, Weber State University, 2507 University Circle, Ogden, UT 84408-2507 (formerly at the Department of Geology, University of Cincinnati).

8. THE BROOKVILLE DAM SPILLWAY—MIAMITOWN THROUGH WAYNESVILLE FORMATIONS (UPPER ORDOVICIAN, SOUTHEASTERN INDIANA)

by
Helen B. Hay and Roger J. Cuffey

SIGNIFICANCE

The Brookville Dam section (fig. 8-1), combined with the Bon Well Hill, Garr Hill, and South Gate Hill sections (see papers 10, 11, and 12 in this volume), form the Brookville composite section (fig. 8-2). Collectively they expose almost the entire Maysvillian and Richmondian succession in southeastern Indiana. “The importance of these sections lies not only in the stratigraphic thickness exposed and the high quality of the exposures, but also in their geographic location along the northwestern side of the Cincinnati Arch. The lower half of the section correlates with strata exposed in Ohio from Cincinnati northward to Middletown, Ohio, and the upper half correlates with outcrops around the margin of the arch. Therefore, the Brookville composite section is a key section in regional stratigraphic work” (Hay, 1977, p. I-6).

LOCATION

The Brookville Dam spillway (IN-FR-0002) is located at the western end of that dam, where the water flows over the dam and down to the river level below, roughly a mile north of Brookville, Franklin County, Indiana (fig. 8-1). From the junction of U.S. Route 52 and Indiana Route 101 near

the northern edge of Brookville, proceed 1.3 miles (2.1 km) northeast on Indiana Route 101, turn sharply northwest for 0.2 mile (0.3 km), and then drive 0.6 mile (1.0 km) west across the top of the dam. The road ends in a turn-around circle overlooking the spillway below and to the southwest. Alternatively, there is also convenient parking along Indiana Route 101 just west of the bridge crossing the Whitewater River, below the dam, only 0.6 mile (1.0 km) northeast of that junction. The spillway section lies in SW¹/₄SE¹/₄SE¹/₄ sec. 17 and NW¹/₄NE¹/₄NE¹/₄ sec. 20, T. 9 N., R. 2 W., Brookville, Indiana, 7.5-minute quadrangle, at 39°26'13"N latitude, 85°00'19"W longitude (4366972 m N, 671679 m E, UTM zone 16). Elevation of the base of the spillway is 623 ft (189.9 meters), at river level in the valley bottom out in front of the dam. From the traffic circle at the west end of the dam-top road, the spillway and the measured section are down to the left (southwest).

Across the spillway to the right (northwest) is a highwall exposing 60 meters (200 ft) of Ordovician bedrock, which correlates with the strata exposed at Garr Hill and the upper part of Bon Well Hill (see papers 10 and 11 in this volume). The height and steepness of the highwall make access very difficult, and, indeed, access is not normally permitted.




FIGURE 8-1.—Location of the Brookville Dam spillway (between the two arrows), on the Brookville (left) and Whitcomb (right), Indiana, 7.5-minute topographic quadrangles (note difficulties in matching contours and roads along their common border).




FIGURE 8-2.—Correlation of the four localities that collectively make up the Brookville composite section: Brookville Dam spillway (IN-FR-0002), Bon Well Hill (IN-FR-0001), Garr Hill/Brookville North (IN-FR-0003), and South Gate Hill/Indiana Route 1, labeled as Cedar Grove (IN-FR-0005).

For information about examining these exposures, contact the U.S. Army Corps of Engineers, Brookville Lake Office, P.O. Box 230, Brookville, Indiana, 47012.

version published some years ago (Hay, 1977, 1981). The lithofacies (fig. 8-4) and the stratigraphic nomenclature used here are discussed more fully in paper 17 of this volume (also see Hay, 1981, and Hay and others, 1981).

STRATIGRAPHY

ACKNOWLEDGMENTS

The stratigraphic section visible in the spillway is summarized in figure 8-3, which is updated from a preliminary

Susan Hovorka, Timothy Knisely, and John Webb helped




FIGURE 8-3.—Stratigraphic section exposed in the Brookville Dam spillway (modified from Hay, 1977, p. I-27, fig. I-11, and Hay, 1981, p. 151).




FIGURE 8-4.—Lithofacies characteristics (see Hay, paper 17 in this volume).

describe the spillway section initially. At the time they were students in Independent Study in Geology at Earlham College.

REFERENCES CITED

Hay, H. B., 1977, Field trip No. 1, Cincinnati stratigraphy from Richmond to Aurora, Indiana, in Pope, J. K., and Martin, W. D., eds., Field guidebook to the biostratigraphy and paleoenvironments of the Cincinnati Series in southeastern Indiana: Society of Economic Paleontologists and Mineralogists, Great Lakes Section, 7th Annual Field Conference, Miami University, p. 11-133 (reprinted as Pope and Martin, 1979; see Appendix B of this volume).

Hay, H. B., 1981, Lithofacies and formations of the Cincinnati Se-

ries (Upper Ordovician), southeastern Indiana and southwestern Ohio: Ph.D. dissertation (unpub.), Miami University, 236 p.

Hay, H. B., Pope, J. K., and Frey, R. C., 1981, Lithostratigraphy, cyclic sedimentation, and paleoecology of the Cincinnati Series in southwestern Ohio and southeastern Indiana, in Roberts, T. G., ed., GSA Cincinnati '81 Field Trip Guidebooks: American Geological Institute, v. 1, p. 73-86.

Addresses of authors: Helen B. Hay, deceased, formerly Geology Department, Earlham College, Richmond, IN 47374; and Roger J. Cuffey, Department of Geosciences (Deike Bldg.), Pennsylvania State University, University Park, PA 16802.

9. THE CORRYVILLE MEMBER OF THE GRANT LAKE FORMATION (UPPER ORDOVICIAN, SOUTHWESTERN OHIO)

by
Lawrence I. Goldman

INTRODUCTION

The Corryville Member of the Grant Lake Formation in southwestern Ohio is a thick-bedded, shale-rich (63 percent) facies that contains the most diverse open-marine fauna in the Cincinnati Series, including rare, articulated macro-invertebrate taxa. The unit is composed of planar-bedded, fossiliferous limestones and thick, platy- to flaggy-parted, sparsely fossiliferous shales. The Corryville Member has been interpreted as a transgressive facies at the base of a third-order shoaling-upward cycle (Tobin, 1982). Deposition was storm dominated and took place in a shallow epeiric sea just offshelf of a broad, gently sloping, west-facing carbonate ramp. The Corryville Member grades south and southwest from the Grant Lake Formation into the stratigraphically equivalent Grant Lake Limestone in Kentucky, a carbonate-rich unit that has been interpreted as having been deposited in a more proximal ramp setting. A specific type section for the Corryville does not exist, so this paper will discuss both the original "type area" and the proposed type section (Schumacher and others, 1991) (fig. 9-1).

DESCRIPTION

HISTORY OF STRATIGRAPHIC NOMENCLATURE

The Corryville Member initially was recognized by Nickles

(1902) from exposures in the Cincinnati area and was named for the community of Corryville, a northern Cincinnati neighborhood within which many of the unit exposures were situated. Nickles defined the unit primarily by the presence and abundance of the characteristic bryozoan *Chiloporella nicholsoni* (fig. 9-7.1) and by the absence of the bryozoan *Monticulipora molesta*, which delineated the underlying Bellevue Beds. Nickles described the Corryville Beds of his Lorraine Group as those in which "the limestones are thinner and less frequent than in the quarry beds and the shales are more yellowish. Blue shale also occurs." (Nickles, 1902, p. 83). The "quarry beds" are the underlying Fairview and Bellevue units. Nickles did not designate a type section for his Corryville Beds. In 1906, Bassler incorporated the Bellevue, Corryville, and the overlying Mt. Auburn Beds as members of the proposed McMillan Formation. Despite the fact that Bassler never published a complete description of the McMillan type section or of its unit boundaries, his McMillan Formation became accepted and uncontested in regional nomenclature for the next half century.

The 1960's and 1970's were a period of reawakened interest in the Cincinnati Series, marked by a general dissatisfaction with the established stratigraphy and an accompanying desire to redefine the entire Cincinnati Series along purely lithological criteria. The original nomenclature was universally rejected for its biostratigraphic underpinnings as well as its inability to account for the




FIGURE 9-1.—Index map showing the locations of site 1, McMillan Street (OH-HA-0027), and site 2, Stonelick Creek (OH-CT-0004).

gradational changes from limestone-dominated facies in northern Kentucky to the equivalent shale-dominated strata of southeastern Indiana. As a result, the name Corryville was abandoned, and its facies was incorporated into larger, undifferentiated lithostratigraphic formations, including the Grant Lake Limestone of Kentucky (Peck, 1966), the "unnamed beds" of Ohio (Ford, 1967), and the Dillsboro Formation (Brown and Lineback, 1966) and the "Brookville formation" (Hay, 1981; Hay and others, 1981), both of Indiana. This proliferation of names and boundaries was useful for the description of local rock units, but contributed little to an overall understanding of regional facies relationships. Because none of these studies directly addressed the stratigraphy in the type area, the original nomenclature continued to be used by those persons working within the Cincinnati area.

Research from the 1980's to the present has concentrated on the interpretation of the large-scale depositional dynamics of the Cincinnati Series and on the regional paleoenvironmental implications of small-scale, intraformational depositional cycles (Tobin, 1982; Jennette, 1986; Diekmeyer, 1990). Tobin (1982) completed the first modern detailed description of the Corryville throughout the southwestern Ohio, southeastern Indiana, and northern Kentucky area. He recognized the Corryville as a mappable unit representative of a distinct deeper water paleoenvironment, and he abandoned the McMillan and reinstated the Corryville as a formation. Tobin (1982, 1986) maintained that, although the original Nickles (1902) descriptions were outdated and lacking in detail, they did represent a valid biostratigraphic and lithostratigraphic identity and so had priority as rock-stratigraphic units.

Schumacher and others (1991) completed a regional documentation of the lithostratigraphic variations within the Bellevue, Corryville, and Mt. Auburn sequence. Through a combination of bedrock mapping, shale-percentage logs, geophysical logs, and mean percentage of shale, Schumacher and others recognized the continuity of these three units within the southwestern Ohio area, but disputed Tobin's (1982, 1986) nomenclature, primarily because the individual units could not be mapped easily at a map scale of 1:62,500 or smaller. In accordance with the guidelines set forth by the 1983 North American Stratigraphic Code (North American Commission on Stratigraphic Nomenclature, 1983), Schumacher and others reduced the Corryville to a member within the proposed Grant Lake Formation of Ohio, a modification of the stratigraphically equivalent, though lithologically disparate, Grant Lake Limestone. The nomenclature of Schumacher and others (1991) has been adopted for this paper.

LITHOLOGY

The Corryville, the middle member of the Grant Lake Formation, is composed of terrigenous shales and interbedded fossiliferous limestones and calcareous siltstones. Bedding contacts are primarily planar to lenticular, although there are minor amounts of irregular- to wavy-bedded limestones within the lower half of the unit. In southwestern Ohio, the Corryville Member averages 63 percent shale and 37 percent limestone. Mean shale percentages decrease to the south and southeast (<40 percent) and increase to the north and northwest (>70 percent) (Tobin, 1982; Schumacher and others, 1991). The Corryville attains a maximum thickness of more than 21 meters (68.9 ft) in central Warren County,

Ohio, and from there thins west, east, southeast, and south. At Maysville, Kentucky, where it is the middle member of the Grant Lake Limestone (Schumacher and others, 1991), its thickness is less than 6 meters (19.7 ft). The Grant Lake Limestone consists of predominantly irregular-bedded limestone (68 percent) and interbedded calcareous, fossiliferous shale (32 percent). Small, accessible stream exposures and road cuts of the Corryville Member of the Grant Lake Formation/Grant Lake Limestone are spread throughout southwestern Ohio, southeastern Indiana, and northern Kentucky (see locality list at the end of this paper and the locality appendix to the volume).

The following description of the Corryville is based mainly on exposures at the Stonelick Creek locality (OH-CT-0004). Carbonate rocks are here classified by texture according to Dunham (1962), as modified by Tobin (1982). Matrix-rich limestones, here referred to as "micstones" (the equivalent of Dunham's "mudstone"), are rare and account for less than 1 percent of all Corryville limestones. The most abundant limestone types are fossiliferous packstones and wackestones. These are poorly sorted, planar, mostly continuous, and in places exhibit upward fining of matrix and fossil content. Grainstones may be coarse, fine, or poorly washed (containing 6-15 percent mud matrix) and are generally more abundant within the lower half of the Corryville section. These limestones are poorly to well sorted, exhibit a moderate to high degree of fossil breakage, and rarely show evidence of bioturbation. Bed contacts are planar, but may be irregular or wavy. Both packstones and grainstones contain internal and external geopetal structures, and mud entrapment by shells is common. Mud clasts and intraclasts are rare. Most of the limestone beds are capped by a distinct, yellowish-gray siltstone, which is sparsely fossiliferous but which may contain trace fossils or articulated brachiopods and pelecypod molds. Bedding surfaces may contain symmetric and asymmetric ripples and current-oriented fossil material, including cephalopods, brachiopods, ramose bryozoans, and crinoids.

Calcsiltites and calcareous siltstones are abundant throughout the Corryville interval and are quite similar in their general appearance. Calcsiltites are defined as very well sorted grainstones with a mean grain size of >0.5 mm and tiny subhorizontally oriented fossil detritus (Tobin, 1982). Siltstones are calcareous and slightly dolomitic and contain small amounts of terrigenous quartz. Both calcsiltite and siltstone strata are thin, generally 1-4 cm (0.4-1.6 inches), but may achieve a thickness of 15 cm (6 inches). Planar lamination, hummocky cross-stratification, and graded bedding are common to both lithotypes. Basal bedding surfaces commonly are erosive and display groove marks, tool marks, flute marks, prod marks, gutter casts, and a lag of small, broken fossils. Fossil material in these beds is sparse, but articulated brachiopods and crinoid stalks of 12 cm (5 inches) or more in length have been found on bedding surfaces. Bioturbation and escape structures are common to both lithotypes, but most traces are restricted to the tops of beds (<2 cm depth) and represent less than 20 to 30 percent reworking of the bed area.

The term "shale" is herein used to describe any predominantly clay-rich bed. In general, Corryville shales are silty, calcareous, and sparsely fossiliferous and display platy to flaggy parting. They are gray to blue when fresh and weather to a yellow-gray color. The average Corryville shale is a dolomitic mudstone with a silt/clay ratio of 1.05 and a mean grain size of 28 micrometers (Tobin, 1982). Average

shale thickness is 7 cm (2.8 inches), with a range of 1 to 34 cm (0.4 to 13.4 inches). Evidence of bioturbation is very rare. A few fossiliferous, fissile-parted beds do exist, but in most instances where fossils are present, they are restricted to distinct horizontal layers within an otherwise sparsely fossiliferous shale bed. When traced laterally in section, these distinct layers generally merge into a discontinuous limestone bed, a strong indication that shale layers may represent more than one depositional event. Thin calcareous horizons and localized concentrations of silt are also abundant within the Corryville shale beds and can be best observed in core sections.

Storm beds and amalgamated beds are two types of strata commonly associated with Corryville deposition. Storm beds are tempestites that contain an erosive base and a basal fossiliferous limestone that grades upward into a calcisiltite and then into a planar- to cross-laminated calcareous siltstone (Brandt, 1986). An amalgamated bed is defined as a multiple-event bed that consists of two or more abruptly overlain siltstones or limestones within a single lithified unit. In most amalgamated beds, a basal fossiliferous packstone or grainstone is sharply overlain by a bioturbated, sparsely fossiliferous, calcareous siltstone. The base of the siltstone is commonly planar and may be erosional. Thicker amalgamated beds will contain several overlain packstones or grainstones that display varying degrees of winnowing, sorting, and breakage of fossils. The contacts between these amalgamated lithologies are planar to irregular and commonly contain thin (<1.0 cm/0.4 inch), discontinuous shale interbeds.

Both the upper and lower contacts of the Corryville Member in the southwestern Ohio area range from sharp to gradational and are differentiated by a change in the mean shale percentage, limestone bedding style, shale parting characteristics, and the fossil content of the shales. The underlying Bellevue Member contains a comparatively low mean shale percentage (17-47 percent), irregular- to wavy-bedded limestone, and fissile-parted, fossiliferous shale (Schumacher and others, 1991). The Bellevue-Corryville contact is sharp at the O'Bannon Creek (OH-CT-0008) and Stonelick Creek localities (OH-CT-0004), and has been placed at the lowest typical Corryville limestone or shale bed. Where the Bellevue-Corryville transition is gradational, the accepted procedure is to arbitrarily place the contact midway between the typical unit lithologies (Schumacher and others, 1991).

The overlying Mt. Auburn Member is dominantly fissile-parted, fossiliferous shale (50 percent) and nodular-bedded limestone (Schumacher and others, 1991). The shale contains small, fossiliferous, discontinuous limestone nodules and a very fine fossil detritus, which gives it a characteristically gritty texture. The Corryville-Mt. Auburn contact is placed at the base of the lowest typical Mt. Auburn shale bed or at the top of the highest typical Corryville limestone bed. Due to the high shale content of both the Corryville and the Mt. Auburn Members, sections containing the upper Corryville contact are rarely well exposed; the best examples of this interval can be seen in small stream exposures at the Hunts Creek (OH-BU-0004), Second Creek (OH-WA-0006), and Lick Run (OH-WA-0004) localities. The large, globose, and thick-shelled brachiopod *Platystrophia ponderosa auburnensis*, known locally as the "double-headed Dutchman," is restricted to the Mt. Auburn, and its initial appearance can assist in the placement of the upper Corryville contact.

SITE DESCRIPTIONS

SITE 1: THE CORRYVILLE "TYPE AREA"

About 1900, when the original stratigraphic nomenclature for the Cincinnati area was established, the most convenient outcrops for study were those exposures located within the neighborhoods bordering downtown. Many of the larger local exposures, such as the one at Fairview Park (OH-HA-0001), were actively quarried for limestone, which was used in the construction of local buildings, walls, and public walkways (Hannibal and Davis, 1992).

Nickles (1902) found abundant outcrops of his Corryville Beds in the Corryville, Clifton, Avondale, Walnut Hills, Fairview Heights, and Price Hill communities (fig. 9-2), apparently enough exposures upon which to recognize the unique lithology and paleontology of the interval, as well as its general stratigraphic position. However, Nickles' (1902) original Corryville description did not designate a specific type section for the unit; hence, these Cincinnati communities, and especially Corryville, have been adopted as an informal "type area" for the unit. This "type area" corresponds to an east-west-trending, present-day topographic high, the result of downcutting by past and present drainage systems. This portion of Cincinnati is represented in portions of four 7.5-minute quadrangles: Cincinnati West (Ohio), Cincinnati East (Ohio), Newport (Ky.-Ohio), and Covington (Ky.-Ohio).

Over the years, Nickles' "type area" has become thoroughly urbanized and, with a few exceptions, all of the oldest Cincinnati exposures have been either demolished or buried. None of the original Corryville exposures is intact. Records from the fossil collections of the University of Cincinnati place previously existing Corryville localities along the southeast side of Highland Avenue, at the top of Fairview Heights, on McMillan Street east of Reading Road, and behind the Seminole Apartments southwest of the intersection of Ravine and McMillan Streets. Loose slabs of Corryville limestone still can be found within the Corryville area; several possibly in situ limestone beds are exposed in the grass-covered bank on the southern end of University Plaza in the Corryville area. New Corryville exposures occasionally appear as a result of building construction or sewer maintenance, but these are usually of temporary duration.

At the time of this writing, only one fairly clean Corryville exposure is present in the "type area": in the empty lot west of 16 McMillan Street (OH-HA-0027), approximately 66 meters (200 ft) from the intersection of Vine and McMillan Streets in the Corryville area (fig. 9-3). This outcrop is located in a small, paved parking space, and the settled appearance of this site does not suggest immediate plans for construction. The exposure has become fairly weathered in the past few years, but the individual limestone, siltstone, and shale beds are still apparent. The base of this 3-meter (12-ft) exposure lies at an approximate elevation of 845 ft (257.6 meters) above sea level; the section most likely represents the middle to upper portions of the Corryville. An excellent example of an amalgamated bed can be found approximately 0.5 meter (1.6 ft) from the base of the exposure. The basal half of this 12-cm-thick (5-inch-thick) bed is a coarse, fossiliferous grainstone. The upper portion of the bed consists of a calcisiltite with an erosive base that fines upward into a cross-laminated and slightly bioturbated siltstone.

The contact between the Bellevue and the Corryville Mem-




FIGURE 9-2.—Location of the Corryville “type area” in Cincinnati, Ohio. The line pattern indicates the area in which the Corryville Member should be the surface bedrock. This area corresponds to elevations between 820 and 865 ft (249.9 and 263.7 meters) above sea level. The small black rectangle marks the location of the McMillan Street exposure (OH-HA-0027) in figure 9-3. The Avondale neighborhood is north of Corryville. The Price Hill neighborhood is southwest of the area shown on this map.



FIGURE 9-3.—A “type” Corryville exposure in the empty, south-facing lot just west of 16 McMillan Street in Cincinnati, Ohio (OH-HA-0027).

bers can be seen in two partially buried exposures, one atop the cliff at the southwestern corner of Bellevue Hill Park (OH-HA-0003), and the other at the top of the small Bellevue exposure just below the transmitting tower above Emming Street (OH-HA-0002). This contact corresponds roughly to an elevation of 820 ft (250.0 meters) above sea level. Although the shales at these sites are buried or recessed, there is an obvious transition from irregular-bedded, brachiopod-rich limestones that contain randomly oriented fossils (Bellevue) to planar-bedded limestones that incorporate subhorizontally oriented fossils that each have a silt cap containing convex-up *Rafinesquina* (Corryville).

The Mt. Auburn contact is not exposed in Cincinnati. Given an approximate thickness of 44 ft (15 meters) in the type area (Caster and others, 1955), the expected elevation for this contact is about 865 feet (264 meters) above sea level. This estimate is corroborated by the comparatively higher elevations of previously existing Mt. Auburn exposures. Given boundary elevations and past and present outcrop sites, a bedrock map can be constructed for the prediction of potential Corryville exposures within the type area (fig. 9-2).

SITE 2: STONELICK CREEK, THE PROPOSED "TYPE SECTION"

Location

Stonelick Creek (OH-CT-0004; fig. 9-4) contains a composite section of the entire Corryville and has been proposed as the official type section for the Corryville Member of the Grant Lake Formation (Schumacher and others, 1991). The Corryville is present in a series of discontinuous stream cuts upstream and downstream of the Ohio Route 131 bridge (39°10'40"N latitude, 84°06'43"W longitude; Newtonsville, Ohio, 7.5-minute quadrangle), Stonelick Township, Cler-

mont County, Ohio, about 1.3 miles (2.2 km) west of the center of Newtonsville, Ohio, about 9.3 miles (15.5 km) east of the city limits of Milford. Corryville exposures range in height from 0.6 meter (2.0 ft) to 18.0 meters (59.0 ft) and occur along both sides of the creek from 1,000 ft (305 meters) north to 1.4 miles (2.3 km) south of the bridge. Two measured sections from Stonelick Creek are on file at the Ohio Geological Survey; measured section no. 16836 represents the lower half of the Corryville, and measured section no. 16837 contains the upper two-thirds of the Corryville Member. Contacts with the underlying Bellevue and the overlying Mt. Auburn Members of the Grant Lake Formation are present, although the latter contact is more gradational and is not exposed well in any single outcrop. Permission to traverse the creek south of the Ohio Route 131 bridge should first be obtained from the landowner at the farm located at 6028 S. Manila Road, just west of the creek. A private lane runs from the landowner's property south along the creek and provides ready access to the lowermost Corryville sections and to the contact with the Bellevue Limestone.

Discussion

Stratigraphic variations within the Corryville section are best understood from cores and from stream outcrops located outside of the Cincinnati area. Stonelick Creek is the most complete and best known Corryville exposure in southwestern Ohio. Most of the rare macrofossil and ichnofossil species attributed to the Corryville have been collected at this site, either in situ from the creek banks or from the loose material lying in the creek bed. Periodic flooding in the spring and fall maintains clean exposures, and new fossil material is continually weathering out of the rock. Water levels at Stonelick Creek remain low throughout most of the summer, fall, and early winter.

The entire Corryville section is exposed at Stonelick




FIGURE 9-4.—Location of Stonelick Creek (OH-CT-0004) in Clermont County, Ohio. Corryville exposures occur along the creek from 1,000 feet (0.3 km) north to 1.4 miles (2.3 km) south of the Ohio Route 131 bridge.

Creek. The lateral continuity of stream cuts facilitates the correlation of meter-scale or larger depositional trends and of smaller lithologically and paleontologically distinct strata. The mean shale percentage at Stonelick Creek is 59 percent, slightly less than the regional average (63 percent). The Corryville Member is approximately 19.4 meters (63.6 ft) thick at this locality (fig. 9-5). Previous estimations of overall unit thickness (Hinterlong, 1981; Schumacher, 1984) differ slightly from this measurement, which is ultimately dependent upon the placement of the gradational upper contact. The contact with the underlying Bellevue Member is also gradational and has been arbitrarily placed at the lowest thick, planar-bedded fossiliferous grainstone. This bed tends to form ledges and is best seen in the southern-most prominent east-facing exposure along the creek (fig. 9-6), about 1.4 miles (2.3 km) south of the Ohio Route 131 bridge, where the creek makes a sharp southward bend. The Fairview Formation below the Bellevue Member is exposed farther downstream, at the Belfast-Owensville Road bridge. The upper contact with the Mt. Auburn Member occurs north of the Ohio Route 131 bridge or near the tops of the highest creek exposures. This contact is gradational and cannot be seen in any single creek-level exposure.

The Corryville Member can be subdivided into four distinct vertical "zones" (fig. 9-5) distinguished by variations in mean shale percentage and shale and limestone lithology. The lowest zone, "Zone 1," is approximately 8.9 meters (29.2 ft) thick and extends upward from the lower contact with the Bellevue. It can be examined best in the first wide, west-facing exposure upstream of the Bellevue contact. This zone is dominated by thick (6 cm/2.4 inches average), continuous to discontinuous packstones, grainstones, and calcisiltites. Shales are slightly fossiliferous, average 6 cm (2.4 inches) in thickness, and account for 54-59 percent of this zone. The brachiopods *Platystrophia* and *Rafinesquina nasuta* are primarily restricted to this zone. The trace fossils *Diplocraterion*, *Trichophycus*, and *Chondrites* type B (Osgood, 1970) are abundant in the calcisiltites and in the silty tops of the limestones. Bedding contacts are mostly planar, with the exception of three separate, distinct, laterally persistent, irregular-bedded packstones. These strata contain abundant randomly oriented shells (*Platystrophia*, *Rafinesquina*) and large, broken ramose bryozoans and are interbedded with fissile, fossiliferous shales that contain small, fossiliferous limestone nodules. The largest of the intervals is 0.5 meter (1.6 ft) thick and is located approximately 5.5 meters (18.0 ft) above the Bellevue contact.

"Zone 2" extends from 8.9 to 10.0 meters (29.2 to 32.8 ft) above the Bellevue contact and is characterized by thick, sparsely fossiliferous shales (8 cm/3 inches average thickness) and interbedded thin calcisiltites, siltstones, and wackestones (4 cm/1.6 inches average thickness). This zone is shale rich (65-72 percent) and is capped by a continuous 12- to 19-cm-thick (4.7- to 7.5-inch-thick) limestone bed that varies lithologically from a poorly washed grainstone to a packstone. Shales may exceed a thickness of 34 cm (13.4 inches). Soft, yellow, discontinuous shales, commonly referred to as "butter shales" or "trilobite shales" (Brandt, 1980; Velbel, 1985), occur in this zone; these fine-grained layers are the source of abundant extended and enrolled specimens of *Flexicalymene* and *Isotelus* (fig. 9-7). Enrollment may have been an instinctive response to deep and abrupt burial, after which the organisms, unable to escape, were preserved articulated and in situ (Brandt Velbel, 1985). Rock of "Zone 2" can be seen in a 3-meter-high (10-ft-high)

north-facing exposure 600 meters (1,970 ft) south of the Ohio Route 131 bridge.

"Zone 3" extends from 10.1 to 13.5 meters (33.1 to 44.3 ft) above the Bellevue contact and is characterized by a lower mean shale percentage (55 percent), thinner shale beds (6 cm/2.4 inches average), and an increased number of packstones and amalgamated beds. Amalgamated beds are abundant in the upper half of this zone and may account for up to 45 percent of all beds in this interval. The lower 1.0 meter (3.3 ft) of this zone is dominated by calcisiltites and calcareous siltstones, and "trilobite shales" are present. Trace fossils are abundant in the siltstones and silty bed tops of strata within this zone. "Zone 3" can be examined at the first north-facing exposure south from the Ohio Route 131 bridge and just beyond the point where the creek bends 90° to the east.

"Zone 4" extends from 13.6 to 19.4 meters (44.6 to 63.6 ft) above the Bellevue contact (to the Mt. Auburn contact), has a mean shale percentage of 63-67 percent, and is thicker and shalier than is "Zone 3." Shales average 9 cm (3.5 inches) in thickness and are sparsely fossiliferous, although fossil content does increase with proximity to the contact with the Mt. Auburn Member. "Zone 4" can be subdivided into four stacked bedding sequences that are similar to "Zone 2" in size and lithology (fig. 9-5). Each shale-rich sequence ranges from 1 to 2 meters (3 to 6 ft) in thickness and contains thin (3 cm/1.2 inches average) and continuous calcisiltites, siltstones, and fossiliferous packstones. The calcisiltites and siltstones are planar- to wavy-bedded and commonly are laminated and display sole marks. The top of each sequence is delineated by a prominent and thick (15 to 41 cm/5.9 to 16.1 inches) amalgamated bed composed of two to four irregular- to planar-bedded grainstones and packstones with two or more thin, discontinuous shale interlayers. "Zone 4" can be seen complete in the higher cliffs along the creek, specifically the steep west-facing exposure at the first 180° creek bend south of the Ohio Route 131 bridge.

PALEONTOLOGY

The Corryville is widely known and appreciated for its abundant and well-preserved open-marine fossil assemblage, the most diverse of any stratigraphic unit of the Cincinnati Series. The limestones in the stratigraphic center of the Corryville Member have the highest generic diversity (number of genera/square meter) (Hinterlong, 1981). The macroinvertebrate fauna includes articulate and inarticulate brachiopods, bryozoans, pelecypods, gastropods, trilobites, orthoconic nautiloids, crinoids, and other echinoderms (table 9-1). Most abundant are trilobite fragments (*Flexicalymene*, *Isotelus*); crinoid columnals; thin, smooth ramose bryozoans; and the brachiopods *Rafinesquina* and *Zygospira*. Pelecypods are commonly preserved as molds. Less common Corryville taxa include ostracodes, conodonts, graptolites, scolecodonts, conularids, and *Cornulites*. Rare fossils in the Corryville include edrioasteroids (fig. 9-7.8), asteroids, the carpoid *Enoploura popei* (Caster, 1952) (fig. 9-7.3), and *Neostrabops martini* (Caster and Macke, 1952), the only aglaspid known from the Cincinnati (fig. 9-7.2). *Enoploura popei* and *N. martini* are known only from the Corryville, *N. martini* only from the Stonelick Creek locality.

Escape structures and other ichnofossils are present and abundant throughout the Corryville (Osgood, 1970; Meyer and others, 1981). These trace fossils are almost always re-




FIGURE 9-5.—Composite stratigraphic column of the Corryville Member of the Grant Lake Formation (Upper Ordovician) from Stonelick Creek (OH-CT-0004) in Clermont County, Ohio.



FIGURE 9-6.—Exposure of the Bellevue-Corryville contact at Stonelick Creek in Clermont County, Ohio (OH-CT-0004). The base of the Corryville is placed at the thick, ledge-forming grainstone near the center of the exposure.

TABLE 9-1.—*Faunal list of the Corryville Member of the Grant Lake Formation*

In the absence of a modern taxonomic listing for the unit, the following informal faunal list has been compiled. All taxa were verified from previous publications, from personal examination of fossiliferous limestone and shale beds, or both. The previous list of Dalvé (1948) is outdated and in need of serious revision.

Taxa	References
	Cnidaria (?)
<i>conulariids</i>	Caster (1942)
	Brachiopoda, Articulata
<i>Hebertella occidentalis</i>	Caster and others (1955), Hinterlong (1981)
<i>Onniella</i>	Tobin (1982)
<i>Plectorthis</i>	Caster and others (1955)
<i>Platystrophia</i> spp. (incl. <i>P. laticosta</i> and <i>P. ponderosa</i>)	Nickles (1902), Braun (1916), Caster and others (1955), Gutstadt (1958), Carpenter and Ory (1961), Ford (1967), Hinterlong (1981), Tobin (1982)
<i>Rafinesquina alternata</i>	Nickles (1902), Van Fossen (1951), Hinterlong (1981), Schumacher (1984), Meyer (1990)
<i>Rafinesquina nasuta</i>	Nickles (1902), Van Fossen (1951), Caster and others (1955), Carpenter and Ory (1961)
<i>Zygospira modesta</i>	Meyer and others (1981), Hinterlong (1981), Tobin (1982), Schumacher (1984)
	Brachiopoda, Inarticulata
<i>Schizocrania filosa</i>	Nickles (1902), Caster and others (1955)
<i>Trematis millepunctata</i>	Nickles (1902), Caster and others (1955), Schumacher (1984)

TABLE 9-1.—*continued*

Taxa	References
	Bryozoa
<i>Amplexopora</i>	Hinterlong (1981)
<i>Batostomella gracilis</i>	Caster and others (1955), Tobin (1982)
<i>Chiloporella nicholsoni</i>	Nickles (1902), Braun (1916), Caster and others (1955), Gutstadt (1958), Hinterlong (1981), Tobin (1982)
<i>Dekayia</i>	Nickles (1902), Caster and others (1955), Hinterlong (1981), Tobin (1982)
<i>Heterotrypa</i>	Nickles (1902), Gutstadt (1958), Hinterlong (1981)
<i>Homotrypa</i>	Caster and others (1955)
<i>Monticulopora</i>	Nickles (1902), Caster and others (1955), Hinterlong (1981), Tobin (1982)
<i>Parvohalloporella ramosa</i>	Caster and others (1955), Gutstadt (1958)
	Annelida
scolecodonts	Schumacher (1984)
	Arthropoda, Aglaspida
<i>Neostrabops martini</i>	Caster and Macke (1952), Hinterlong (1981)
	Arthropoda, Ostracoda
ostracodes	Dalvé (1948), Tobin (1982)
	Arthropoda, Trilobita
<i>Acidaspis</i>	Hinterlong (1981)
<i>Amphilichas halli</i>	Dalvé (1948)
<i>Ceraurus milleranus</i>	Nickles (1902), Caster and others (1955)
<i>Flexicalymene</i> (incl. <i>F. meeki</i>)	Caster and others (1955), Hinterlong (1981), Tobin (1982), Schumacher (1984), Frey (1987b)
<i>Isotelus</i>	Caster and others (1955), Hinterlong (1981), Schumacher (1984), Frey (1987b)
	Mollusca, Cephalopoda
<i>Camerocheras</i>	Dalvé (1948 as <i>Endocheiras</i>), Frey (1987b)
<i>Oncoceras</i>	Frey (1987b)
<i>Treptoceras</i>	Dalvé (1948), Frey (1987b)
	Mollusca, Gastropoda
<i>Bellerophon</i>	Nickles (1902)
<i>Cyclonema</i>	Nickles (1902), Tobin (1982), Schumacher (1984)
<i>Cyclora</i>	Tobin (1982)
<i>Loxoplocus bowdeni</i>	Caster and others (1955), Tobin (1982)
	Mollusca, Monoplacophora
<i>Cyrtolites</i>	Dalvé (1948)
	Mollusca, Pelecypoda
<i>Ambonychia</i> (incl. <i>Byssonychia</i>)	Nickles (1902), Braun (1916), Pojeta (1962), Tobin (1982), Schumacher (1984), Frey (1987a, b)
<i>Anomalodonta</i>	Braun (1916), Frey (1987a)
<i>Caritodens demissa</i>	Pojeta (1971), Schumacher (1984), Frey (1987a)
<i>Cleidophorus</i>	Frey (1987a)
<i>Corallidomus</i>	Frey (1987a, b)
<i>Ctenodonta</i>	Frey (1987a)
<i>Cuneamya</i>	Frey (1987a)
<i>Cycloconcha</i>	Frey (1987a)
<i>Cymatonota</i>	Frey (1987a)
<i>Deceptrix</i>	Frey (1987a)
<i>Lyrodesma</i>	Frey (1987a)
<i>Modiolopsis</i>	Nickles (1902), Schumacher (1984), Frey (1987a)
<i>Orthodesma</i>	Frey (1987a, b)
<i>Pholadomorpha</i>	Frey (1987a)
<i>Psiloconcha</i>	Frey (1987a)
<i>Rhytima</i>	Frey (1987a)
	Echinodermata, Asteroidea
<i>Promopalaeaster</i>	Schumacher (1984)

Taxa	References
<i>Enoploura popei</i>	<p style="text-align: center;">Echinodermata, "Carpoidea"</p> Caster (1952), Hinterlong (1981)
<i>Anomalocrinus</i> <i>Cincinnatiocrinus</i> <i>Iocrinus</i> (incl. <i>I. subcrassus</i>) "Lichenocrinus" <i>Ohioocrinus</i> <i>Pycnoocrinus</i> (Including what is commonly called <i>Glyptocrinus dyeri</i>)	<p style="text-align: center;">Echinodermata, Crinoidea</p> Nickles (1902) Dalvé (1948) Hinterlong (1981), Meyer and others (1981), Schumacher (1984), Frey (1987b) Nickles (1902), Caster and others (1955) Nickles (1902) Nickles (1902), Caster and others (1955), Hinterlong (1981), Meyer and others (1981), Schumacher (1984), Frey (1987b)
<i>Carneyella pilea</i> <i>Isorophus cincinnatensis</i> <i>Streptaster vorticellatus</i>	<p style="text-align: center;">Echinodermata, Edrioasteroidea</p> Caster and others (1955), Bell (1972, 1976), Meyer and others (1981), Meyer (1990) Foerste (1914), Caster and others (1955), Kesling and Mintz (1960), Bell (1972, 1976), Meyer and others (1981), Meyer (1990) Bell (1972), Meyer and others (1981), Meyer (1990)
graptolites	<p style="text-align: center;">Hemichordata, Graptolithina</p> Schumacher (1984), Bergström and Mitchell (1986)
<i>Asteriacites stelliforme</i> <i>Chondrites</i> type B <i>Diplocraterion biclavatum</i> <i>Fascifodina floweri</i> <i>Lockeia siliquaria</i> <i>Palaeophycus</i> type B <i>Palaeoscia floweri</i> <i>Phycodes flabellum</i> <i>Rusophycus carleyi</i> <i>Rusophycus pudicum</i> <i>Trichophycus venosum</i> "escape structures"	<p style="text-align: center;">Trace fossils</p> Osgood (1970), Meyer and others (1981) Osgood (1970), Hinterlong (1981), Meyer and others (1981), Tobin (1982, 1986) Osgood (1970), Hinterlong (1981), Meyer and others (1981), Tobin (1982, 1986) Osgood (1970), Meyer and others (1981) Osgood (1970), Meyer and others (1981) Osgood (1970), Hinterlong (1981), Meyer and others (1981), Tobin (1982, 1986) Caster (1942), Osgood (1970), Hinterlong (1981) Osgood (1970), Meyer and others (1981) Osgood (1970), Meyer and others (1981) Osgood (1970), Hinterlong (1981), Meyer and others (1981) Osgood (1970), Hinterlong (1981), Meyer and others (1981), Tobin (1982, 1986) Tobin (1982)
conodonts <i>Cornulites corrugatus</i> <i>Lepidocoleus</i> (machaeridian)	<p style="text-align: center;">Other</p> Bergström and Mitchell (1986) Schumacher (1984) Dalvé (1948), Schumacher (1984)


stricted to the siltstones, calcisiltites, silty tops of beds, and the more silty portions of amalgamated beds. Common trace fossils include *Chondrites* type B (fig. 9-7.6), *Trichophycus venosum* ("turkey tracks") (fig. 9-7.5), and *Diplocraterion biclavatum* (the "dumbbell" U-shaped burrow); these particular ichnofossils commonly occur in association in the thicker calcareous siltstone layers. Less common ichnofossils are *Lockeia siliquaria*, *Asteriacites stelliforme* (an impression of an asteroid echinoderm), *Phycodes flabellum*, *Paleophycus* type B, *Rusophycus carleyi*, and *Rusophycus pudicum*. *Rusophycus carleyi* and *R. pudicum* have been interpreted as the resting and feeding traces of *Isotelus* and *Flexicalymene*, respectively (Osgood, 1970). All three known specimens wherein a *Flexicalymene* was found in situ within a *R. pudicum* burrow were discovered from the Corryville section of Stonelick Creek (Osgood, 1970). Unique to the Corryville are *Fascifodina floweri*, a vertical bundle of tunnel systems set in a semi-elliptical pattern, and *Palaeoscia floweri*, a series of regular to irregular concentric circles marked by a central pore (Osgood, 1970). *Conostichus*, a conical, convex

hyporelief believed to be a sea-anemone burrow (Pemberton and others, 1988), is present in calcisiltites and siltstones within the 2 meters (6.6 ft) of section below the upper Corryville contact.

Corryville fossils exhibit a variety of preservational modes, from original shell material to molds. Individual brachiopod shells range in appearance from pristine to fragmented to heavily encrusted and bored. Disarticulated trilobite and crinoid fragments are common, but show little to no abrasion and corrosion, implying little transport and a parautochthonous origin. Fossil orientation within and at the surface of beds is dominantly subhorizontal, but may in places become imbricated or shingled or may display no preferential orientation at all. The upper bedding surfaces of many of the packstones and grainstones exhibit abundant whole or disarticulated convex-up pelecypod and brachiopod shells, commonly referred to as shell "pavements." The shells of one *Rafinesquina* pavement display a strongly bimodal NE-SW orientation that has been attributed to a preburial storm surge in the same direction (Meyer, 1990). Brachiopods, with

FIGURE 9-7.—Fossils of the Corryville Member of the Grant Lake Formation. UCGM = University of Cincinnati Geology Museum.

- 1 *Chiloporella nicholsoni*, UCGM uncatalogued specimen, X1. Nickles (1902) gave the authorship of this species as "(James)," but Bassler (1915) gave it as Nickles and Bassler, 1900, and specifically did not recognize *Ceramopora nicholsoni* U. P. James, 1875.
- 2 *Neostrabops martini* Caster and Macke, 1952, dorsal view of holotype, UCGM 25569, X1.
- 3 *Enoploura popei* Caster, 1952, convex (? dorsal) side of carapace, holotype, UCGM 25993, X1.
- 4 *Flexicalymene meeki* (Foerste, 1910), enrolled specimen from Stonelick Creek, Clermont County, Ohio, X1.
- 5 *Trichophycus venosum* (Miller), three specimens on siltstone, X0.2.
- 6 *Chondrites* type B Osgood, 1970, X0.25.
- 7 *Iocrinus subcrassus* (Meek & Worthen, 1865), uppermost Corryville, UCGM 44362, X0.25.
- 8 *Isorophus cincinnatiensis* (Roemer), on valve of brachiopod shell, X1.



the exception of *Platystrophia*, are dominantly thin shelled and are commonly preserved as unbroken valves with little to no corrosion or abrasion. Byrozoan morphology is dominated by small- to medium-sized ramose colonies, but foliate and encrusting varieties are plentiful as well. Massive foliate and ramose bryozoan colonies over 30 cm (1 ft) in diameter can be found intact in limestones and shales; most of these are overturned and exhibit multiple surface borings.

PALEOENVIRONMENTAL INTERPRETATION

Certain taphonomically unique Corryville beds have been ascribed directly to storm-related, catastrophic burial (Brandt, 1980; Meyer and others, 1981; Brandt Velbel, 1985; Meyer, 1990). Examination of these deposits has produced a wealth of paleoecological data concerning the population composition, size-frequency distribution, preferred life mode, and substratum preference of Late Ordovician invertebrates. In addition, paleoenvironmental data, including current orientation, amount of preburial transport, surface exposure time, and burial timing and conditions, can be interpreted from these deposits. These beds have been dually categorized as "event beds" (strata attributed to rapid and episodic sedimentation) and "Fossil-Lagerstätten" (unique, well-preserved fossil assemblages that yield an unusual amount of paleontological and taphonomic information). Examples of such "event beds" include deposits containing abundant, articulated *Iocrinus subcrassus* (fig. 9-7.7) collected from siltstones just below the Corryville-Mt. Auburn contact (Schumacher, 1984) and the aforementioned "trilobite shales." "Fossil-Lagerstätten" is an appropriate description for the abundant, articulated edrioasteroids (*Isorophus*, *Streptaster*, and *Carneyella*) discovered intact on *Rafinesquina* shells atop two upper Corryville brachiopod shell pavements (Meyer and others, 1981; Meyer, 1990) (fig. 9-7.8). Rassman (1981), in a study of one of these pavements, concluded that it was an incipient hardground formed by early diagenetic marine cementation.

The Corryville Member of the Grant Lake Formation has been interpreted as having been deposited in an offshore environment situated along a gently sloping, west-facing carbonate ramp in a shallow epeiric sea (Tobin, 1982). The Corryville represents an offshore facies situated at the base of a third-order shoaling-upward cycle; its lower contact with the Bellevue Member marks the beginning of an abrupt transgression from a more shallow, turbulent depositional environment to a comparatively deeper water environment (Tobin, 1982; Tobin and Pryor, 1985). The relative abundance of siltstones and shales, the relative thickness of the shale beds, the excellent preservation of trace fossils and sedimentary structures, and the abundance of whole, unabraded, thin-shelled and delicate invertebrate macrofossils all indicate low-energy deposition, probably below wave base, in a protected, quiet-water environment (Tobin, 1982). Depositional trends related to the eastward shallowing of the ramp include a decrease in Corryville shale content and average shale thickness toward the south and east and an increase in the abundance of grainstones to the south and east (Tobin, 1982; Schumacher and others, 1991).

Corryville deposition was dominated by episodic storm events (Tobin, 1982). The large numbers of storm beds, amalgamated beds, and current-oriented, storm-related sedimentary structures support this interpretation. Limestones are most likely parautochthonous tempestites; grainstones are indicative of proximal deposition, and siltstones and

calcisiltites are representative of distal deposition (Tobin, 1982). Taphonomic evidence from smothered hardgrounds and the "trilobite shales" indicates that shale deposition was, at least at times, both rapid and episodic (Brandt, 1980; Meyer and others, 1981; Brandt Velbel, 1985; Meyer, 1990) and that some shale beds record more than one depositional event. The four zones evident at Stonelick Creek are distinct lithologically and paleontologically and can be correlated in field exposures and core sections in southwestern Ohio. The existence of such traceable zones implies temporal variations in depositional environment on a regional scale. Small-scale bedding cycles in the Kope and Fairview Formations, lower in the Cincinnati, have been attributed to fluctuations in absolute sea level (Jennette, 1986). Such fluctuations also may be the cause of depositional variations in the Corryville Member.

ADDITIONAL CORRYVILLE LOCALITIES

The following list is divided into two parts, one for the Grant Lake Formation and the second for the Grant Lake Limestone, as used by the Ohio Division of Geological Survey. For more information on a given site, see Appendix A at the end of this volume.

CORRYVILLE MEMBER OF GRANT LAKE FORMATION

Bellevue Hill (OH-HA-0003). Partial exposure at top of cliff at the southwestern corner of park. The Bellevue-Corryville contact is present at this site and is set at a prominent limestone bed which forms a small ledge near the cliff edge. Corryville limestones display a distinctive silt covering on the bed top and convex-up brachiopods. Most shales are recessed or buried.

Blue Rock Road (OH-HA-0005). Between 1.0 meter (3.3 ft) and 3.9 meters (12.8 ft) of Corryville is present, but mostly buried, at the base of the cut. The upper 7.6 meters (25.0 ft) of section is composed of the Mt. Auburn Member and the Sunset Formation. Both a shell pavement with abundant edrioasteroids and a horizon containing abundant specimens of the crinoid *Iocrinus subcrassus* have been excavated from the Corryville section at this site (Meyer, 1990; Schumacher, 1984).

"Big 4" railroad cut (Maud cut) (OH-BU-0002). Most of this exposure is Mt. Auburn, but Corryville limestones can be found at the base of two cliffs along the tracks.

Dornbusch (OH-HA-0026). This section contains 11.2 meters (36.7 ft) of the Corryville Member.

Elk Creek (OH-BU-0003). This stream cut exposes 5.5 meters (18.0 ft) of the Corryville interval.

Emming Street (OH-HA-0002). The Bellevue-Corryville contact is present, but partially buried. Corryville limestones are exposed in the small clearing on top of the small west-facing cliff exposure.

Hunts Creek (OH-BU-0004). This locality contains the upper 7.4 meters (24.2 ft) of the Corryville Member, the Corryville-Mt. Auburn contact, and 1.3 meters (4.3 ft) of the Mt. Auburn Member.

Lick Run (OH-WA-0004). The contact with the overlying Mt. Auburn is present and sharp; the contact is best seen in the westernmost north-facing exposure.

Middletown (OH-BU-0005). The Corryville is present in the lower 5.0 meters (16.4 ft) of the road cut. The lower half of the section, including the Corryville-Mt. Auburn contact,

is partially buried underneath a layer of talus. This locality is the type section for Hay's (1981) "Excello member" of the "Brookville formation," referred to as the "Excello" locality.

Muddy Creek (OH-HA-0020). Most of this exposure in the stream cut just south of Muddy Creek Road bridge is the Bellevue Member, but the Bellevue-Corryville contact occurs near the top of the exposure.

O'Bannon Creek (OH-CT-0008). The Bellevue-Corryville contact is present about 4 meters above the stream bed and is sharp. This Corryville exposure contains nearly all of the first "zone." The 0.5-meter (1.6-ft) irregular-bedded horizon is located 5.5 meters (18.0 ft) above the Bellevue-Corryville contact, the same stratigraphic position as at Stonelick Creek.

Ravine Street (OH-HA-0028). A few small Corryville slabs are present and loose in this poor, mostly buried exposure behind the Seminole Apartments southwest of the intersection of Ravine and McMillan Streets in Cincinnati.

Rumpke Landfill (OH-HA-0029). Inaccessible exposure in a field east of U.S. Route 27 N (Colerain Avenue). The almost 14 meters (46 ft) of Corryville and the Corryville-Mt. Auburn contact once were exposed here, but have been buried as a result of landfill activity.

Second Creek (OH-WA-0006). The uppermost Corryville interval and the Corryville-Mt. Auburn contact are present at a small waterfall close to the Cozzandale Road bridge. The Mt. Auburn is well exposed northeast of the bridge just around the first stream meander.

Sharonville (OH-HA-0030). The Corryville can be seen best at the easternmost side of the field just above the highest zone of light-gray, sparsely fossiliferous Miami town Shale.

Sheits Road (OH-HA-0023). This locality exposes 12.8 meters (42 ft) of the Bellevue Member, the Bellevue-Corryville contact, and 5.6 meters (18.4 ft) of the lower Corryville.

Todd Run (OH-CT-0009). 6.4 meters (21.0 ft) of the Corryville Member are exposed in the stream cut.

West Middletown (OH-BU-0006). Exposed here are 5.0 meters (16.4 ft) of Corryville section, overlain by 5.2 meters (17.1 ft) of the Mt. Auburn Member.

Westwood-Northern Boulevard (OH-HA-0031). Many of the beds in this exposure are thick calcisiltites; this interval may represent the upper Corryville.

Wynbrook Apartments (OH-HA-0025). Exposed here are rocks of the Miami town Shale and the Bellevue, Corryville, and Mt. Auburn Members of the Grant Lake Formation. The Corryville begins at the first bench up from the parking-lot level.

CORRYVILLE MEMBER OF GRANT LAKE LIMESTONE

Eagle Creek (OH-BR-0005). This section begins at stream level approximately 795 ft (242.3 meters) above sea level. The exposure contains 3.4 meters (11.2 ft) of the Corryville Member, the contact of the Corryville and the overlying Straight Creek Member (Mt. Auburn equivalent), and 5.5 meters (18.0 ft) of the Straight Creek Member.

Georgetown (OH-BR-0006). The Grant Lake Limestone is 28.8 meters (94.4 ft) thick at this locality and complete; the contact with the overlying Bull Fork Formation is gradational, and the contact with the underlying Fairview Formation is sharp (Lee, 1974).

Goose Run (OH-BR-0007). This series of stream cuts exposes the uppermost 4.0 meters (13.1 ft) of the Corryville

Member, the contact of the Corryville and overlying Straight Creek Member (Mt. Auburn equivalent), and 5.5 meters (18.0 ft) of the Straight Creek Member.

Manchester (OH-AD-0003). Exposed in this road cut is the contact with the overlying Bull Fork Formation (Lee, 1974); this site is within a few hundred meters of the Isaacs Creek exposure of Sweet (1979).

Ripley #2 (OH-BR-0008). This exposure reportedly exposes a sharp and obvious contact between the even-bedded limestones and shales of the underlying Fairview Formation and the overlying 19.7 meters (64.6 ft) of irregular-bedded Grant Lake Limestone (Lee, 1974).

Sleepy Hollow (KY-MS-0005). This is the type section of the Grant Lake Limestone and is reported to be 33.4 meters (109.6 ft) thick. The upper contact with the Bull Fork Formation and the lower contact with the Fairview Formation are gradational within a 3-meter (10-ft) interval (Lee, 1974).

Thomas More Parkway (KY-KE-0002). This site was exposed in 1989 and consists of three broad benches bounded by several 2- to 3-meter (6- to 10-ft), west-facing exposures. The Bellevue may be present in the lowest exposure.

White Oak Creek (OH-BR-0009). This stream cut exposes 6.5 meters (21.3 ft) of the Corryville Member and begins at stream level, approximately 860 ft (262 meters) above sea level.

REFERENCES CITED

- Bassler, R. S., 1906, A study of the James types of Ordovician and Silurian Bryozoa: Proceedings of the U.S. National Museum, v. 30, no. 1442, p. 1-66.
- _____, 1915, Bibliographic index of American Ordovician and Silurian fossils: U.S. National Museum Bulletin 92, 2 v., 1,521 p.
- Bell, B. M. 1972, A study of North American Edrioasteroidea: Ph.D. dissertation (unpub.), University of Cincinnati, 1,022 p.
- _____, 1976, A study of North American Edrioasteroidea: New York State Museum and Science Service Memoir 21, 447 p.
- Bergström, S. M., and Mitchell, C. E., 1986, The graptolite correlation of the North American Upper Ordovician Standard: *Lethaia*, v. 19, p. 247-266.
- Brandt, D. S., 1980, Phenotypic variation and paleoecology of *Flexicalymene* (Arthropoda: Trilobita) in the Cincinnati Series (Upper Ordovician) near Cincinnati, Ohio: M.S. thesis (unpub.), University of Cincinnati, 148 p.
- _____, 1986, Preservation of event beds through time: *Palaeos*, v. 1, p. 92-96.
- Brandt Velbel, Danita, 1985, Ichnologic, taphonomic, and sedimentologic clues to the deposition of Cincinnati shales (Upper Ordovician), Ohio, in Curran, H. A., ed., Biogenic structures: their use in interpreting depositional environment: Society of Economic Paleontologists and Mineralogists Special Publication 35, p. 299-307.
- Braun, E. L., 1916, The Cincinnati Series and its brachiopods in the vicinity of Cincinnati: Cincinnati Society of Natural History Journal, v. 22, p. 18-44.
- Brown, G. D., Jr., and Lineback, J. A., 1966, Lithostratigraphy of the Cincinnati Series (Upper Ordovician) in southeastern Indiana: American Association of Petroleum Geologists Bulletin, v. 50, p. 1018-1023.
- Carpenter, J. W., and Ory, T. R., 1961, The American Upper Ordovician Standard. VI. The Covington Sequence at Maysville, Kentucky: Ohio Journal of Science, v. 61, p. 372-378.
- Caster, K. E., 1942, Two siphonophores from the Paleozoic: *Palaeontographica Americana*, v. 3, no. 14, 34 p.
- _____, 1952, Concerning *Enoploura* of the Upper Ordovician and its relation to other carpod echinoderms: *Bulletins of American Paleontology*, v. 34, no. 141, 56 p.
- Caster, K. E., Dalvé, E. A., and Pope, J. K., 1955, Elementary guide to the fossils and strata of the Ordovician in the vicinity of Cincinnati, Ohio: Cincinnati Museum of Natural History, 47 p.

- Caster, K. E., and Macke, W. B., 1952, An aglaspid merostome from the Upper Ordovician of Ohio: *Journal of Paleontology*, v. 26, p. 753-757.
- Dalvé, Elizabeth, compiler, 1948, The fossil fauna of the Ordovician in the Cincinnati region: University Museum, Department of Geology and Geography, University of Cincinnati, 56 p.
- Diekmeyer, S. C., 1990, Quantitative analysis of cyclical faunal patterns in the Upper Ordovician Kope through Bellevue sequence, Cincinnati, Ohio: M.S. thesis (unpub.), University of Cincinnati, 168 p.
- Dunham, R. J., 1962, Classification of carbonate rocks according to depositional texture, in Ham, W. E., ed., *Classification of carbonate rocks: American Association of Petroleum Geologists Memoir 1*, p. 108-121.
- Foerste, A. F., 1914, Notes on Agelacrinidae and Lepadocystinae, with descriptions of *Thresherodiscus* and *Broccocystis*: *Scientific Laboratories of Denison University Bulletin*, v. 17, art. 14, p. 399-487.
- Ford, J. P., 1967, Cincinnati geology in southwest Hamilton County, Ohio: *American Association of Petroleum Geologists Bulletin*, v. 51, p. 918-936.
- Frey, R. C., 1987a, The occurrence of pelecypods in early Paleozoic epeiric-sea environments, Late Ordovician of the Cincinnati, Ohio area: *Palaios*, v. 2, p. 3-23.
- _____, 1987b, The paleoecology of a Late Ordovician shale unit from southwest Ohio and southeastern Indiana: *Journal of Paleontology*, v. 61, p. 242-267.
- Gutstadt, A. M., 1958, Upper Ordovician stratigraphy in eastern interior region: *American Association of Petroleum Geologists Bulletin*, v. 42, p. 513-547.
- Hannibal, J. T., and Davis, R. A., 1992, Guide to the building stones of downtown Cincinnati: a walking tour: *Ohio Division of Geological Survey Guidebook 7*, 44 p.
- Hay, H. B., 1981, Lithofacies and formations of the Cincinnati Series (Upper Ordovician), southeastern Indiana and southwestern Ohio: Ph.D. dissertation (unpub.), Miami University, 236 p.
- Hay, H. B., Pope, J. K., and Frey, R. C., 1981, Lithostratigraphy, cyclic sedimentation, and paleoecology of the Cincinnati Series in southwestern Ohio and southeastern Indiana, in Roberts, T. G., ed., *GSA Cincinnati '81 Field Trip Guidebooks: American Geological Institute*, v. 1, p. 73-86.
- Hinterlong, G. D., 1981, Paleoecologic succession in the Corryville Member (McMillan Formation, Upper Ordovician) Stonelick Creek, Clermont County, Ohio: M.S. thesis (unpub.), Miami University, 119 p.
- Jennette, D. C., 1986, Storm-dominated cyclic ramp deposits of the Kope-Fairview transition (Upper Ordovician), southwestern Ohio and northern Kentucky: M.S. thesis (unpub.), University of Cincinnati, 210 p.
- Kesling, R. V., and Mintz, L. W., 1960, Internal structures in two edrioasteroid species, *Isorophus cincinnatiensis* (Roemer) and *Carneyella pilea* (Hall): *University of Michigan Museum of Paleontology Contributions*, v. 15, no. 14, p. 315-348.
- Lee, G. B., 1974, Lithostratigraphy of the Cincinnati Series (Upper Ordovician) from Maysville, Kentucky to Dayton, Ohio: M.S. thesis (unpub.), Miami University, 127 p.
- Meyer, D. L., 1990, Population paleoecology and comparative taphonomy of two edrioasteroid (Echinodermata) pavements: Upper Ordovician of Kentucky and Ohio: *Historical Biology*, v. 4, no. 3, p. 155-178.
- Meyer, D. L., Tobin, R. C., Pryor, W. A., Harrison, W. B., Osgood, R. G., Hinterlong, G. D., Krumpolz, B. J., and Mahan, T. K., 1981, Stratigraphy, sedimentology, and paleoecology of the Cincinnati Series (Upper Ordovician) in the vicinity of Cincinnati, Ohio, in Roberts, T. G., ed., *GSA Cincinnati '81 Field Trip Guidebooks: American Geological Institute*, v. 1, p. 31-72.
- Nickles, J. M., 1902, The geology of Cincinnati: *Cincinnati Society of Natural History Journal*, v. 20, no. 2, p. 49-100; reprint edition, p. 1-53.
- North American Commission on Stratigraphic Nomenclature, 1983, *North American Stratigraphic Code: American Association of Petroleum Geologists Bulletin*, v. 67, p. 841-875.
- Osgood, R. G., Jr., 1970, Trace fossils of the Cincinnati area: *Palaeontographica Americana*, v. 6, no. 41, p. 277-444.
- Peck, J. H., 1966, Upper Ordovician formations in the Maysville area, Kentucky: *U.S. Geological Survey Bulletin 1244-B*, 30 p.
- Pemberton, S. G., Frey, R. W., and Bromley, R. G., 1988, The ichnotaxonomy of *Conostichus* and other plug-shaped ichnofossils: *Canadian Journal of Earth Sciences*, v. 25, p. 866-892.
- Pojeta, John, Jr., 1962, The pelecypod genus *Byssonychia* as it occurs in the Cincinnati at Cincinnati, Ohio: *Palaeontographica Americana*, v. 4, no. 30, p. 166-216.
- _____, 1971, Review of Ordovician pelecypods: *U.S. Geological Survey Professional Paper 695*, 46 p.
- Rassman, Barbara, 1981, A sedimentologic comparison of two dissimilar Ordovician edrioasteroid pavements (hardgrounds): M.S. thesis (unpub.), University of Cincinnati, 224 p.
- Schumacher, G. A., 1984, Sedimentology, biostratigraphy, and paleoecology of Cincinnati (Upper Ordovician) *Iocrinus subcrassus* crinoid (Echinodermata) assemblages: M.S. thesis (unpub.), University of Cincinnati, 210 p.
- Schumacher, G. A., Swinford, E. M., and Shrake, D. L., 1991, Lithostratigraphy of the Grant Lake Limestone and Grant Lake Formation (Upper Ordovician) in southwestern Ohio: *Ohio Journal of Science*, v. 91, p. 56-68.
- Tobin, R. C., 1982, A model for cyclic deposition in the Cincinnati Series of southwestern Ohio, northern Kentucky, and southeastern Indiana: Ph.D. dissertation (unpub.), University of Cincinnati, 483 p.
- Tobin, R. C., 1986, An assessment of the lithostratigraphic and interpretive value of the traditional "biostratigraphy" of the type Upper Ordovician of North America: *American Journal of Science*, v. 286, p. 673-701.
- Tobin, R. C., and Pryor, W. A., 1985, Cincinnati Series—model for cyclic and episodic deposition of carbonates and shales on a storm-dominated ramp (abs.): *American Association of Petroleum Geologists Bulletin*, v. 69, p. 311-312.
- Van Fossen, J. D., 1951, A study of the Rafinesquinae of the Middle Maysville (Upper Ordovician): M.S. thesis (unpub.), University of Cincinnati, 98 p.

Address of author: Lawrence I. Goldman, MINEsoft, Ltd., 165 S. Union Blvd., #510, Lakewood, CO 80228.

10. THE “BROOKVILLE FORMATION” (“EXCELLO,” WAYNESVILLE, AND LIBERTY MEMBERS) AT BON WELL HILL NEAR BROOKVILLE (UPPER ORDOVICIAN, SOUTHEASTERN INDIANA)

by
Helen B. Hay and Roger J. Cuffey

SIGNIFICANCE

The Bon Well Hill road cut (fig. 10-1) exposes the middle portion of the Brookville composite section (fig. 10-2; Hay, 1977, p. I-5 to I-6). In particular, the disconformable contact between the “Excello” and the Waynesville is well exposed here, as well as the middle portion of the “Brookville Formation” of Hay (1981; Hay and others, 1981).

LOCATION

The Bon Well Hill locality (IN-FR-0001) is the long road cut on Indiana Route 101, 1.3-1.7 miles (2.1-2.7 km) north-east of its junction with U.S. Route 52 inside the north edge of Brookville, Franklin County, Indiana (fig. 10-1). The base of the section, at elevation 747 ft (227.7 meters), extends northwestward along the entry road leading to the top of the Brookville Dam. The road cut is in S¹/₂SW¹/₄SE¹/₄ sec. 16, T. 9 N., R. 2 W., Whitcomb, Indiana, 7.5-minute quadrangle, and is centered at 39°26'15"N, 84°59'25"W (4367024 m N, 672966 m E, UTM zone 16).

STRATIGRAPHY AND PALEONTOLOGY

The Bon Well Hill locality was described initially by Hay (1977, p. I-23 to I-26), and later revised and updated by Hay and others (1981, p. 79, 82-84, as Stop 3). The following is taken almost completely from the latter work.

Limestone and shale strata of the “Excello member” of the

“Brookville formation” are exposed along the driveway to the dam and at the base of the long section facing Indiana Route 101. Blue-gray shale of the Waynesville Shale Member of the “Brookville formation” occupies most of the section along Route 101. The Liberty Member of the “Brookville formation” occurs at the top of the hill.

The general stratigraphic description and overall faunal chart are provided in figures 10-3 and 10-4, respectively. Discussion of the lithofacies recognized here (fig. 10-5) and stratigraphic nomenclature can be found in Hay (paper 17 in this volume), as well as in Hay (1981). There is a stratigraphic gap of about 9 meters (30 ft) between the top of this road cut and the base of the Garr Hill section (IN-FR-0003) on the basis of study of the continuous exposure at South Gate Hill (IN-FR-0005; see paper 12 in this volume).

Most of the “Excello member” at this outcrop is facies 2b, although the shales are less limy and the limestones less rubbly than they are in the average occurrence of this facies. The other two facies in the “Excello” are 1b with more than 70 percent shale compared with 55-70 percent for facies 2b, and a thin band of facies 3a, cross-bedded calcarenites interbedded with sandy-textured, brown, phosphatic shales at the top of the member. The Waynesville Shale Member contains two intervals of facies 1a separated by a medial band of facies 2a. There is a gradual increase in limestone in the upper half of the outcrop, and the base of the Liberty Member is placed where the shale percentage drops below 70 percent. As figure 10-4 indicates, there are several faunal zones in this outcrop which are related to the lithofacies.

The most interesting question about this section is the nature of the contact between the “Excello” and Waynesville Shale Members and the cause of the abrupt and widespread facies change across the contact. The lower part of the Waynesville Shale Member is nearly devoid of fossils and the few indurated beds in the shales are calcareous siltstones rather than the normal bioclastic limestones. This barren shale overlies the cross-bedded calcarenites at the top of the “Excello,” not only at this locality, but throughout most of the northern two-thirds of the outcrop area in southeastern Indiana and southwestern Ohio. This contact is easy to recognize both in cores and in outcrop. Cumings and Galloway (1913) recognized a significant faunal as well as lithologic change across this boundary. It is possible that there is a disconformity between the calcarenites and the basal Waynesville shales, or, alternatively, the disconformity may be just under the calcarenites. In the former case the calcarenites could represent the culmination of the regression, and in the latter case, they could be a basal transgressive “sandstone.” If there is a disconformity, then the Waynesville Shale Member is a transgressive unit in which the clastics were derived from erosion of the exposed land, and the absence of fossils is probably due to turbid conditions or unfavorable chemical conditions, perhaps brackish water. The upper half of the Waynesville is much more fossiliferous than the lower half. It contains a diverse molluscan assemblage although brachiopod diversity is low. If this interpretation of the Waynesville Shale Member is correct, then facies 1a in this case represents a different




FIGURE 10-1.—Location of the Bon Well Hill road cut (BWH; ends indicated by arrows). Whitcomb, Indiana, 7.5-minute quadrangle.




FIGURE 10-2.—Correlation of the four localities that collectively make up the Brookville composite section: Bon Well Hill (IN-FR-0001), Brookville Dam spillway (IN-FR-0002), Garr Hill/Brookville North (IN-FR-0003), and South Gate Hill/Indiana Route 1, labeled as Cedar Grove (IN-FR-0005).

environment than is interpreted for the same facies in the Kope. This “Excello”-Waynesville Shale Member boundary is interpreted as the end of the second transgressive-regressive cycle and the beginning of the third, which culminated in deposition of the sediments of the Whitewater and Saluda Formations, which are exposed in the outcrop at Garr Hill (IN-FR-0003).

Beyond the data provided in the overall faunal chart (fig. 10-3), the faunas of the Bon Well Hill section, especially the mollusks, were discussed by Frey (*in* Hay and others, 1981); most of the next several paragraphs is taken from his discussion:

The well-known Upper Ordovician fauna characteristic of




FIGURE 10-3.—Stratigraphic section exposed at Bon Well Hill (from Hay and others, 1981, p. 82, fig. 7; with permission).

the classical Cincinnati area is best known for its well-preserved articulate brachiopods and trepostome ectoprocts. A closer examination of the Upper Ordovician section exposed at Bon Well Hill, however, indicates that some lithologic units, especially the more argillaceous limestones and claystones, have faunas dominated by mollusks, particularly pelecypods. Some of these argillaceous strata have large, diverse faunas of mollusks, trilobites, and echinoderms. One such unit is the "trilobite shale," which is located in the Waynesville Shale Member of the "Brookville formation," exposed at the 15-meter (49-ft) interval at this locality (fig. 10-3). Fifty species of invertebrates have been collected and identified from this 1.5-meter-thick (5-ft-thick) claystone unit; 13 of these are pelecypods. Pelecypods dominated brachiopods in Paleozoic shallow-marine environments where local conditions were such that mobility was an asset and nutrient supply was plentiful. The "trilobite shale" was probably deposited under such conditions, as articulate brachiopods and trepostomes are uncommon, whereas the endobysate suspension-feeding pelecypods *Ambonychia suberecta* (Ulrich), *Corallidomus concentricus* (Hall and

Whitfield), and *Pholadomorpha pholadiformis* (Hall) are abundant (densities up to 160 individuals/m²) and reach large size (up to 75 mm in length). Brachiopods become plentiful only at the very top of the shale where bedding planes are covered with valves of the ubiquitous "pioneer" genera *Rafinesquina* and *Onniella*.

In contrast to the richness of the fauna of the "trilobite shale," the lowermost shales of the Waynesville Shale Member have a sparse fauna of linguloid brachiopods, small paleotaxodont pelecypods, and occasional trilobites. This portion of the Waynesville Shale Member is thought to represent very shallow water conditions and a more stressful environment than that of the "trilobite shale" unit. Other molluscan-rich assemblages include a 2-meter (6.6-ft) band of rubbly calcareous mudstone near the top of the "Excello member," which is exposed at the lower (southwest) end of the Bon Well Hill exposure. This unit contains specimens of the ambonychiid pelecypods *Ambonychia alata* Meek and *Anomalodonta gigantea* S. A. Miller plus the cylindrical pelecypod *Orthodesma recta* Hall and Whitfield, which is often preserved in its presumed life position, with the posterior end of the shell directed up, perpendicular to bedding planes. The platycerid gastropod *Cyclonema humerosum* Ulrich is also abundant in this unit.

The strata at the base of the Bon Well Hill section would be referred to the upper Arnheim or Oregonia of the traditional Cincinnati units.

REFERENCES CITED

- Amsden, T. W., 1983, The Late Ordovician brachiopod genera *Lepidocyclus* and *Hiscobeccus*: Oklahoma Geological Survey Bulletin 132, p. 36-44.
- Cumings, E. R., and Galloway, J. J., 1913, The stratigraphy and paleontology of the Tanner's Creek section of the Cincinnati Series in Indiana: Indiana Department of Geology and Natural Resources, Thirty-seventh Annual Report for 1912, p. 353-479.
- Hay, H. B., 1977, Field trip No. 1, Cincinnati stratigraphy from Richmond to Aurora, Indiana, in Pope, J. K., and Martin, W. D., eds., Field guidebook to the biostratigraphy and paleoenvironments of the Cincinnati Series in southeastern Indiana: Society of Economic Paleontologists and Mineralogists, Great Lakes Section, 7th Annual Field Conference, Miami University, p. 11-133 (reprinted in Pope and Martin, 1979; see Appendix B of this volume).
- Hay, H. B., 1981, Lithofacies and formations of the Cincinnati Series (Upper Ordovician), southeastern Indiana and southwestern Ohio: Ph.D. dissertation (unpub.), Miami University, 236 p.
- Hay, H. B., Pope, J. K., and Frey, R. C., 1981, Lithostratigraphy, cyclic sedimentation, and paleoecology of the Cincinnati Series in southwestern Ohio and southeastern Indiana, in Roberts, T. G., ed., GSA Cincinnati '81 Field Trip Guidebooks: American Geological Institute, v. 1, p. 73-86.

Addresses of authors: Helen B. Hay, deceased, formerly Geology Department, Earlham College, Richmond, IN 47374; and Roger J. Cuffey, Department of Geosciences (Deike Bldg.), Pennsylvania State University, University Park, PA 16802.

Principal diagnostic characteristics of lithofacies	
Group 1 >70% shale	1a Well-bedded, mostly thin limestones and medium to thick, fissile to blocky shales. Includes massive units of 100% shale. Low-diversity fauna.
	1b Rubbly, shaly, nodular limestones; medium to thick shales with limy stringers and layers of fossils in shale matrix. May have some well-bedded limestones as in 1a. Low-diversity fauna. Most easily seen in cores.
Group 2 55-70% shale	2a Like 1a except has high diversity of fauna. Intermediate in shale percentage between 1a and 3a.
	2b Well-bedded limestones, rubbly nodular limestones, and limy shales. Allochems tend to be more finely broken than 2a. High-diversity fauna. Intermediate between 3a and 3c.
Group 3 <55% shale	3a Well-bedded, thin to medium limestones and thin to medium, fissile to blocky shales. High-diversity fauna.
	3b Well-bedded, thin to medium limestones with or without rubbly or nodular shales. High-diversity fauna. Intermediate between 3a and 3c.
	3c Thin-bedded, rubbly weathering, argillaceous limestones, very thin limy shales. High-diversity fauna.
	3d Limestones and shales as in 3a except >20% of limestones are fine-grained laminated burrowed, barren of large fossils; trails on tops. Grade to siltstones.
Group 4 Miscellaneous	4a High-angle cross-bedded calcarenites, bimodal dip. Little or no shale.
	4b Granule-size fossils in thick to massive beds; little or no shale. May be like 4a if seen in similar outcrops.
	4c Massive <i>Loxoplocus</i> limestone bed. Marble Hill Bed.
	4d Bioturbated light-gray limestone with low-diversity fauna of ostracodes, algae, snails, and <i>Tetradium</i> . Related to Saluda facies complex.




FIGURE 10-5.—Lithofacies characteristics (see Hay, paper 17 in this volume).

11. LIBERTY, LOWER AND UPPER WHITEWATER, AND SALUDA STRATA AT GARR HILL/BROOKVILLE NORTH (UPPER ORDOVICIAN, SOUTHEASTERN INDIANA)

by
Helen B. Hay and Roger J. Cuffey

SIGNIFICANCE

The Garr Hill road cut (fig. 11-1) exposes the upper part of the Brookville composite section (fig. 11-2), which is discussed and correlated in Hay and Cuffey's discussion on the Brookville Dam spillway locality (paper 8 in this volume; see also Hay, 1977, p. I-5 to I-6, and Hay and others, 1981). Particularly noteworthy here are the abundant, diverse fossils at the base of the section and the *Tetradium* coral zone and Saluda Formation near the top.

LOCATION

The Garr Hill/Brookville North locality (IN-FR-0003) is the long road cut on both sides of Indiana Route 101, 6.0-6.4 miles (9.6-10.2 km) northeast of the junction of U.S. Route 52 and Indiana Route 101 at the north edge of Brookville,

Franklin County, Indiana (fig. 11-1). The road cut is 0.8-1.2 miles (1.3-1.9 km) south from the crossroads at the office of the Mounds State Recreation Area, which is about 10 miles (16 km) south of Liberty, Indiana. Elevation of the base of the measured section is 900 ft (274.3 meters). The road cut occupies the E¹/₂SE¹/₄NE¹/₄ and E¹/₂NE¹/₄SE¹/₄ sec. 35, T. 10 N., R. 2 W., Whitcomb, Indiana, 7.5-minute quadrangle, at 39°29'11" to 39°29'29"N, 84°56'58"W (4372556-4373108 m N, 676410 m E, UTM zone 16). The name "Garr Hill" appears on the topographic map about a mile southwest of the road cut, but applies to the entire complex of hills in the surrounding vicinity.

STRATIGRAPHY AND PALEONTOLOGY

Hay (1977, p. I-18 to I-22) presented a preliminary version of the Garr Hill/Brookville North section and later updated it (Hay and others, 1981, p. 80, 84-86). Much of the following discussion is taken from Hay and others (1981).

The Garr Hill section includes, from bottom to top, the Liberty Member of the "Brookville formation," the lower part of the Whitewater Formation, a thin, coral-bearing Saluda unit, and the basal beds of the upper part of the Whitewater Formation (fig. 11-3). Local ranges of important fossil species are given in figure 11-4. Hay (1981), Hay and others (1981), and Hay (paper 17 in this volume) discuss the lithofacies identified here (fig. 11-5) and the stratigraphic classification applied to these rocks.

The base of the road cut exposes fossiliferous gray limestone and blue- to light-gray shale of the Liberty Member of "Brookville formation." The Liberty is overlain by lenticular to rubbly limestones and shales of the Whitewater Formation. A relatively thin interval of the Saluda Formation, consisting of calcisiltite strata including zones of tabulate corals (*Tetradium*) is intercalated in the Whitewater strata.

This outcrop is of interest not only because it exposes several different facies and faunal assemblages (figs. 11-3, 11-4), but also because it shows a stratigraphic sequence of facies interpreted as the regressive phase of the third transgressive-regressive cycle in the Cincinnati. The transgressive phase of the third cycle probably began with deposition of the muds of the Waynesville Shale Member (Bon Well Hill; see paper 10 in this volume), and the peak of the transgression is represented by facies 2a at the base of this locality, Garr Hill. There is a gap of 9 meters (30 ft) between these two outcrops, based on comparison with the continuous exposure at South Gate Hill (see paper 12 in this volume).

Facies 2a at the base of the Garr Hill/Brookville North section has an abundant and extremely diverse fauna. The abundant and well-preserved brachiopods represent more genera than occur together at any other stratigraphic position in the Cincinnati of this area. Horn corals first appear in Indiana and Ohio at this level; bryozoans, trilobites, echinoderms, and mollusks are also abundant. This faunal assemblage suggests deposition on a stable open shelf, in comparison to the less stable, more rigorous conditions indicated by the lower diversity and the lack of fossils near the base of the underlying Waynesville Shale Member. The abraded fossils, some jumbled together in coquinalike beds,




FIGURE 11-1.—Location of the Garr Hill/Brookville North road cut (extent marked by arrows). Whitcomb, Indiana, 7.5-minute quadrangle.




FIGURE 11-2.—Correlation of the four localities that collectively make up the Brookville composite section: Garr Hill/Brookville North (IN-FR-0003), Bon Well Hill (IN-FR-0001), Brookville Dam Spillway (IN-FR-0002), and Indiana Route 1/South Gate Hill, labeled as Cedar Grove (IN-FR-0005).

indicate periodic high-energy conditions, which would be expected on a shallow open shelf.

The increase in limestone percentage upward in this outcrop, from facies 2a to 3a to 3d, is interpreted as an indication of shoaling, so that sediments of the later two facies were deposited at or near wave base. The contact between the well-bedded rocks of the Liberty Member and the rubbly rock (facies 3c) of the Whitewater Formation is gradational

here, as it is at many other localities. The gradational strata, assigned to facies 3b, are included in the Whitewater Formation. The rubbly rock of facies 3c is well exposed at the top of the outcrop, where it is somewhat weathered.

The *Tetradium* coral layer and associated facies 4d of the Saluda Formation is well exposed in this outcrop. Hatfield (1968) interpreted the massive *Tetradium* colonies as serving the function of a wave baffle marginal to a shallow




FIGURE 11-3.—Stratigraphic section at Garr Hill/Brookville North (from Hay and others, 1981, p. 84, fig. 9; with permission).

lagoonal setting that produced the silty dolomites that are characteristic of the Saluda Formation farther south in Indiana. At this locality, however, only the coral zone and the burrowed micritic limestones of facies 4d occur. These burrowed limestones of facies 4d have an unusual fauna dominated by algae and small mollusks and probably formed around the edge of the Saluda lagoon just inside the *Tetradium* wave-baffle zone. The Saluda sediments bearing mud cracks undoubtedly were deposited in very shallow

water and were intermittently exposed to the atmosphere.

Facies 3c, the characteristic facies of the Whitewater Formation, also represents very shallow conditions on the basis of its stratigraphic association with the Saluda and its unique lithologic characteristics. Facies 3c may be envisioned as representing the relatively quiet interior of a shallow platform. Toward the margin of the platform, sediments of facies 3b were deposited, and at the platform margin, facies 3a. In slightly deeper water the shallier sediment of facies 2a accumulated. This scenario is supported by analysis of facies sequences, characteristics of individual facies, and the geometry of these facies bodies throughout the region. Superposition of these facies, as in this outcrop, indicates regression.

Correlation of the Garr Hill/Brookville North locality with the Richmond composite section to the north is discussed in paper 15 in this volume.

REFERENCES CITED

- Amsden, T. W., 1983, The Late Ordovician brachiopod genera *Lepidocyclus* and *Hiscobeccus*: Oklahoma Geological Survey Bulletin 132, p. 36-44.
- Hatfield, C. B., 1968, Stratigraphy and paleoecology of the Saluda Formation (Cincinnatian) in Indiana, Ohio, and Kentucky: Geological Society of America Special Paper 95, 34 p.
- Hay, H. B., 1977, Field trip No. 1, Cincinnatian stratigraphy from Richmond to Aurora, Indiana, in Pope, J. K., and Martin, W. D., eds., Field guidebook to the biostratigraphy and paleoenvironments of the Cincinnatian Series in southeastern Indiana: Society of Economic Paleontologists and Mineralogists, Great Lakes Section, 7th Annual Field Conference, Miami University, p. 11-133 (reprinted in Pope and Martin, 1979; see Appendix B of this volume).
- Hay, H. B., 1981, Lithofacies and formations of the Cincinnatian Series (Upper Ordovician), southeastern Indiana and southwestern Ohio: Ph.D. dissertation (unpub.), Miami University, 236 p.
- Hay, H. B., Pope, J. K., and Frey, R. C., 1981, Lithostratigraphy, cyclic sedimentation, and paleoecology of the Cincinnatian Series in southwestern Ohio and southeastern Indiana, in Roberts, T. G., ed., GSA Cincinnati '81 Field Trip Guidebooks: American Geological Institute, v. 1, p. 73-86.

Addresses of authors: Helen B. Hay, deceased, formerly Geology Department, Earlham College, Richmond, IN 47374; and Roger J. Cuffey, Department of Geosciences (Deike Bldg.), Pennsylvania State University, University Park, PA 16802.




FIGURE 11-4.—Fauna of the Garr Hill/Brookville North section (from Hay and others, 1981, p. 85, fig. 10; with permission). *Lep.* = *Leptaena*; *S. planumbona* = *Strophomena planumbona*; *Tet.* = *Tetradium*. Note that *Lepidocyclus capax* (Conrad, 1842) was assigned to the genus *Hiscobecus* by Amsden in 1983.

Principal diagnostic characteristics of lithofacies	
Group 1 >70% shale	1a Well-bedded, mostly thin limestones and medium to thick, fissile to blocky shales. Includes massive units of 100% shale. Low-diversity fauna.
	1b Rubbly, shaly, nodular limestones; medium to thick shales with limy stringers and layers of fossils in shale matrix. May have some well-bedded limestones as in 1a. Low-diversity fauna. Most easily seen in cores.
Group 2 55-70% shale	2a Like 1a except has high diversity of fauna. Intermediate in shale percentage between 1a and 3a.
	2b Well-bedded limestones, rubbly nodular limestones, and limy shales. Allochems tend to be more finely broken than 2a. High-diversity fauna. Intermediate between 3a and 3c.
Group 3 <55% shale	3a Well-bedded, thin to medium limestones and thin to medium, fissile to blocky shales. High-diversity fauna.
	3b Well-bedded, thin to medium limestones with or without rubbly or nodular shales. High-diversity fauna. Intermediate between 3a and 3c.
	3c Thin-bedded, rubbly weathering, argillaceous limestones, very thin limy shales. High-diversity fauna.
	3d Limestones and shales as in 3a except >20% of limestones are fine-grained laminated burrowed, barren of large fossils; trails on tops. Grade to siltstones.
Group 4 Miscellaneous	4a High-angle cross-bedded calcarenites, bimodal dip. Little or no shale.
	4b Granule-size fossils in thick to massive beds; little or no shale. May be like 4a if seen in similar outcrops.
	4c Massive <i>Loxoplocus</i> limestone bed. Marble Hill Bed.
	4d Bioturbated light-gray limestone with low-diversity fauna of ostracodes, algae, snails, and <i>Tetradium</i> . Related to Saluda facies complex.




FIGURE 11-5.—Lithofacies characteristics (see Hay, paper 17 in this volume).

12. “EXCELLO” (ARNHEIM) TO BASAL SALUDA STRATA ON INDIANA ROUTE 1 AT SOUTH GATE HILL (UPPER ORDOVICIAN, SOUTHEASTERN INDIANA)

by
Helen B. Hay, Brian Kirchner, and Roger J. Cuffey

SIGNIFICANCE

The spectacular road cuts on the long slope down South Gate Hill (fig. 12-1) expose 61 meters (200 ft) of section, including much of the Richmondian Stage in southeastern Indiana. When added to the exposure at the Brookville Dam Spillway (IN-FR-0002), a total stratigraphic interval of 98 meters (310 ft) is exposed within a distance of about 6 miles, extending from the Maysvillian Miami town Shale at the base to the Richmondian Saluda Formation at the top (fig. 12-2). Included in this interval are the Bellevue Formation and all of the “Brookville formation.” Also important is the geographic location of these outcrops north and west along the Cincinnati Arch; the Brookville Dam spillway provides the northernmost known exposure of Maysvillian rocks in Indiana. Therefore, the four outcrops in the vicinity of Brookville—the Brookville Dam spillway (IN-FR-0002), Bon Well Hill (IN-FR-0001), Garr Hill (IN-FR-0003), and this locality, South Gate Hill—are significant for regional correlation with cores and other outcrops in Indiana and Ohio. The other Brookville localities are each described in other papers in this volume (8, 10, and 11, respectively). This locality also correlates with the Madison/U.S. Route 421 sections (IN-JE-0001, IN-JE-0002, and IN-JE-0003) described in paper 6 in this volume.

LOCATION

South Gate Hill (IN-FR-0005), also called the Indiana Route 1 or the Cedar Grove locality, consists of road cuts on both sides of Indiana Route 1, 1.0-1.5 miles (1.6-2.4 km) north of the crossroads at the village of South Gate and 1.9-2.4 miles (3.0-3.8 km) south of the junction with U.S. Route 52 west of Cedar Grove, Franklin County, Indiana (fig. 12-1). These cuts are in the $W\frac{1}{2}W\frac{1}{2}SE\frac{1}{4}$ sec. 23, T. 8 N., R. 2 W., Cedar Grove, Indiana, 7.5-minute quadrangle, at $39^{\circ}20'08''$ to $39^{\circ}20'30''N$ and $84^{\circ}57'20''$ to $84^{\circ}57'24''W$ (4355853-4356469 m N, 676098-676194 m E, UTM zone 16). Elevation of the base of this section is about 750 ft (228.6 meters). The highway has recently been greatly improved, but its roadbed apparently remains in almost the same position as on the published topographic map. Recently, signs forbidding digging here have appeared, and so appropriate caution or conservation should be observed.

STRATIGRAPHIC DESCRIPTION

Exposed in the South Gate Hill road cuts are strata of the Saluda Formation, the lower part of the Whitewater Formation, and the “Brookville formation”; the “Brookville” includes the Liberty Member, the Waynesville Shale Member, and the “Excello member” (fig. 12-3). Initial field descriptions were reported by Kirchner (1991); they are updated in this present paper, although recording of the local ranges of fossils is not yet finalized and, hence, such are not included. Lithofacies recognized here (fig. 12-4) are discussed more fully by Hay (1981), Hay and others (1981), and Hay (paper 17 in this volume).

SALUDA FORMATION

At this location 9+ ft (2.7+ meters) of Saluda is exposed. The facies is 4d and consists of thoroughly burrowed, micritic, light-gray limestone containing *Tetradium* in some beds. There is a large *Tetradium* colony in float at the top of the section. No thin sections are available from this location; thin sections from Garr Hill show that the fossils present are algae with structure preserved, ostracodes, and small snails; very few are normal-marine fossils of the Cincinnati. No dolomite was found either at Garr Hill or South Gate Hill, although facies 4d is associated with Saluda dolomitic facies farther west and southwest, giving confidence to the assignment of this interval to the Saluda.

WHITEWATER FORMATION

At this locality 5 ft (1.5 meters) of the lower or sub-Saluda part of the Whitewater Formation is present. The rocks consist primarily of facies 3b—planar or rubbly limestones and limy shales; limestone is >50 percent. There are some beds of facies 4d (characteristic of the Saluda); thus, it possibly could be included in the Saluda. Overall, this unit is transitional between Whitewater and Saluda lithologies.

“BROOKVILLE FORMATION”

The informal name “Brookville formation” was proposed by Hay and others (1981, p. 76) for the rocks between the top of the Bellevue Limestone and the bottom of the Whitewater Formation. (It is not the same rock unit as the Brookville Clay of Pennsylvania nor the Brookville Terrane of Kansas.) The “Brookville formation,” as used here, is the same as much of the upper part of the Dillsboro Formation, as recognized by the Indiana Geological Survey (Brown and Lineback, 1966; Gray, 1972; Hay, paper 17 in this volume). It likewise is equivalent to the Corryville through Liberty interval of Caster, Dalvé, and Pope (1955), and the Corryville Member of the Grant Lake Formation through the Liberty Formation of the Ohio Division of Geological Survey (Schumacher and others, 1991).

LIBERTY MEMBER

The Liberty Member here has 115 ft (35.1 meters) present. The section is described in five parts.

(1) The upper 35.5 ft (10.8 meters) is mostly facies 3a; <55 percent shale with planar limestones; a majority of the shales lack limy lenses, stringers, or nodules (not limy shales). Facies 3d at the top of the interval is like 3a except that >20 percent of the limestones are composed of sand- and silt-size allochems and are laminated and burrowed. Facies 3d also occurs within the Liberty at Garr Hill.

(2) The upper interval of facies 1a is 8 ft (2.4 meters) thick and is >70 percent shale; this unit is a marker unit in the outcrop (fig. 12-3, 150-foot level). Hay surmised that the top of this shale may be where the Ohio Geological Survey might put the contact between the Waynesville and the




FIGURE 12-1.—Location of the South Gate Hill road cuts (SGH; arrows indicate extent of exposures) along Indiana Route 1, Cedar Grove, Indiana, 7.5-minute quadrangle.

Liberty; if this interpretation is correct, the Liberty at this locality would be 10.8 meters (35.5 ft) thick, rather than 35.1 meters (115 ft) as stated here, and the Waynesville would be 36.7 meters (120.5 ft) thick, rather than 12.5 meters (41.0 ft). For discussion of this suggested modification of the Waynesville-Liberty boundary, see Hay (paper

17 in this volume).

(3) Facies 2a is 20 ft (6.1 meters) thick and is like facies 3a above except that the shale percentage is 55-70 percent. The *Thaerodonta* peak zone is either within this interval or within the 3a zone just above the 150-foot level.

(4) Facies sequence 2b, 3b, 2b is 36.5 ft (11.1 meters) thick.




FIGURE 12-2.—Correlation of the South Gate Hill section (IN-FR-0005), here labeled Indiana Rte. 1 Cedar Grove, with the other three localities that collectively make up the Brookville composite section: Bon Well Hill (IN-FR-0001), Brookville Dam spillway (IN-FR-0002), and Garr Hill/Brookville North (IN-FR-0003).

Facies 2b is 55-70 percent shale with limy shales and planar or rubbly limestones; facies 3b is the same as 2b except that it contains <55 percent shale. Faunal diversity increases from the lower part of this interval upward. The *Glyptorthis* interval is tentatively located within the 3b unit, the 2b unit above it, or both; this interval correlates faunally with the

base of the Garr Hill section.

(5) The lower interval of facies 2a is 14.5 ft (4.4 meters) thick and is the same as (3) above. The boundary between the Liberty and the Waynesville (at the base of the unit) is placed to include in the Liberty the strata in which the shale percentage is predominantly <70 percent. The shale




FIGURE 12-3.—Stratigraphic section exposed at South Gate Hill. Profile shows topographic benches, faces, and road signs along the west side of Indiana Route 1 as of late 1991. Although the position of the disconformity is placed above the interval of facies 3a, it is possible that the disconformity lies below this interval. See figure 12-4 for explanation of the lithologic patterns and facies.




FIGURE 12-4.—Lithofacies characteristics (see Hay, paper 17 in this volume).

percentage decreases upward to <55 percent in the upper part of the Liberty, as noted above.

WAYNESVILLE SHALE MEMBER

At this locality 41 ft (12.5 meters) of the Waynesville is present, consisting of mostly facies 1a (>70 percent shale), except for 9 ft (2.7 meters) of facies 2a in the middle. The shales are not limy and the limestones are planar rather than rubbly. The lowest part of the member, just above the top of the "Excello member," is nearly barren of fossils; indurated beds are siltstones rather than fossiliferous limestones. Fossil content increases upward; the brachiopod fauna is dominated by *Onniella* and *Rafinesquina*. The Waynesville/"Excello" boundary is interpreted to be a disconformity (Hay, 1981; Hay and others, 1981; Hay, paper 17 in this volume).

"Excello member"

At this locality, 22+ ft (6.7+ meters) of "Excello" (informal name) is exposed; it is described below in two parts. The portion of the "Excello" exposed here is the upper part of

the Arnheim of Ohio usage (Schumacher and others, 1991). (Note that this is not the Excello Shale of Oklahoma.)

(1) Facies 3a is 8.5 ft (2.6 meters) thick and is <55 percent shale. Limestones are rather well-sorted calcarenites separated by brown shales with phosphatized steinkerns of small brachiopods, mollusks, and other fossils, which give these beds a sandy texture compared to most Cincinnati gray shales. It is not certain whether this unit of facies 3a is just above or just below the disconformity.

(2) Facies 2b is 13.5+ ft (4.1+ meters) thick and is 55-70 percent shale. Shales are limy, and limestones are rubbly or planar. The limestones tend to be lighter gray than those higher in the section and contain a large fraction of sand-size allochems, including many crinoid ossicles.

PALEOENVIRONMENTAL INTERPRETATION

This facies sequence and the characteristics of the superimposed facies suggest an interpretation of changing environments through time that can be summarized as follows: (1) Regression, or at least a very shallow water environment, existed at the time the "Excello" sediments were deposited, with emergence producing a disconformity between them

and the Waynesville Shale Member. Facies 3a at the top of the "Excello" may represent a beach in the regressive phase or the basal unit of the Waynesville transgression; in the latter case, this unit should be included in the Waynesville rather than in the "Excello." (2) The scarcity of fossils in the lowest part of the Waynesville followed upward by an increase in fossil abundance and diversity represents a transgressive phase that probably culminated in facies 2a in the Liberty at water depth below but near storm wave base. Therefore, the environment in which the sediments of the Waynesville Shale Member were deposited is best interpreted as very shallow water rather than relatively deep water. (3) Following the peak of the transgression in the basal facies 2a of the Liberty, there may have been a period of slight shoaling that resulted in deposition of the facies 2b-3b sequence, followed by a renewed minor transgression marked by the symmetrical overlying sequence of 2b-2a. In other words, the upward facies sequences of 2a, 2b, 3b, 2b, 2a probably represents a minor regressive phase followed by a minor transgressive phase. (4) The increase in limestone relative to shale (facies of group 3) near the top of the Liberty suggests a final shoaling phase that culminated in the environment responsible for the Saluda facies, 4d, which was probably deposited in extremely shallow water of somewhat higher than normal salinity, based on its association with dolomite and *Tetradium* at other localities (Hatfield, 1968; Hay, paper 17 in this volume). The depositional environments of the Saluda and the "Excello"/Waynesville disconformity are two of the significant anchors to which this interpretation is tied.

The changes in water depth described here are probably only slight; all of these sediments indicate depositional environments above or, more commonly, not far below, storm wave base. The role of storms in final deposition of the carbonate sediments was probably very significant. Although extended discussion and justification for these interpretations is not feasible here, it is available elsewhere (Hay, 1981; paper 17 in this volume).

REFERENCES CITED

- Brown, G. D., Jr., and Lineback, J. A., 1966, Lithostratigraphy of the Cincinnati Series (Upper Ordovician) in southeastern Indiana: American Association of Petroleum Geologists Bulletin, v. 50, p. 1018-1023.
- Caster, K. E., Dalvé, E. A., and Pope, J. K., 1955, Elementary guide to the fossils and strata of the Ordovician in the vicinity of Cincinnati: Cincinnati Museum of Natural History, 47 p.
- Gray, H. H., 1972, Lithostratigraphy of the Maquoketa Group (Ordovician) in Indiana: Indiana Geological Survey Special Report 7, 31 p.
- Hatfield, C. B., 1968, Stratigraphy and paleoecology of the Saluda Formation (Cincinnati) in Indiana, Ohio, and Kentucky: Geological Society of America Special Paper 95, 34 p.
- Hay, H. B., 1981, Lithofacies and formations of the Cincinnati Series (Upper Ordovician), southeastern Indiana and southwestern Ohio: Ph.D. dissertation (unpub.), Miami University, 236 p.
- Hay, H. B., Pope, J. K., and Frey, R. C., 1981, Lithostratigraphy, cyclic sedimentation, and paleoecology of the Cincinnati Series in southwestern Ohio and southeastern Indiana, in Roberts, T. G., ed., GSA Cincinnati '81 Field Trip Guidebooks: American Geological Institute, v. 1, p. 73-86.
- Kirchner, Brian, 1991, Description, correlation, and interpretation of Cincinnati Series lithofacies in the Indiana State Route 1 roadcut near Brookville, Indiana: Independent study in geology, Earlham College, 42 p.
- Schumacher, G. A., Swinford, E. M., and Shrake, D. L., 1991, Lithostratigraphy of the Grant Lake Limestone and Grant Lake Formation (Upper Ordovician) in southwestern Ohio: Ohio Journal of Science, v. 91, p. 56-68.

Addresses of authors: Helen B. Hay, deceased, and Brian Kirchner, formerly Geology Department, Earlham College, Richmond, IN 47374; and Roger J. Cuffey, Department of Geosciences (Deike Bldg.), Pennsylvania State University, University Park, PA 16802.

13. UPPER ARNHEIM THROUGH LOWER WHITEWATER STRATA AT CAESAR CREEK LAKE (UPPER ORDOVICIAN, SOUTHWESTERN OHIO)

by
Gregory A. Schumacher, Douglas L. Shrake, E. Mac Swinford, Lynn A. Rockwell, and Roger J. Cuffey

SIGNIFICANCE

Exposures in the emergency spillway and below the dam of Caesar Creek Lake permit close examination of the abundantly fossiliferous, later Cincinnati strata of the northeastern side of the Cincinnati Arch. These exposures can be profitably compared to those on the northwestern/western side of the arch (see papers in this volume on localities near Richmond, Brookville, and Madison, Indiana), as well as farther southward. Moreover, although the entire type-Cincinnati is rightly regarded as fossiliferous, the strata in the Caesar Creek spillway are exceptionally so, to the point that the State of Ohio and the U.S. Army Corps of Engineers have taken the unusual step of permitting fossil collecting within this state park.

LOCATION

The Caesar Creek emergency spillway and dam are located about 3 miles (5 km) southeast of Waynesville, Warren County, Ohio (figs. 13-1, 13-2). They are situated at the southwestern end of Caesar Creek Lake/Reservoir, constructed and maintained by the U.S. Army Corps of Engineers. Caesar Creek State Park surrounds the lake.

From the junction of U.S. Route 42 and Ohio Route 73 at the south corner of Waynesville, proceed 1.1 miles (1.7 km) southeast on Route 73, turn south (at the Corps of Engineers sign) on Clarksville Road, and drive 3.2 miles (5.1 km) to the parking area in the middle of the spillway (fig. 13-2). En route, 2.3 miles (3.7 km) from Route 73, the Caesar Creek Visitors Center lies 0.2 mile (0.3 km) to the left (southeast). Continuing 0.2 mile (0.3 km) farther, a road to the right (northwest) descends into the gorge below the dam; and 0.3 mile (0.5 km) still farther, the main road crosses the top of the dam.

The parking area along Clarksville Road in the center of the emergency overflow spillway (OH-WA-0001) is located at 39°28'49"N, 84°03'25"W (4373976 m N, 753156 m E, UTM zone 16), on the Oregonia, Ohio, 7.5-minute quadrangle. (This area has not been surveyed into townships and ranges.) The topmost exposed strata lie at an elevation of 925 ft (281.9 meters). Clarksville Road crosses the floor of the emergency spillway at an elevation in the center of 883 ft (269.1 meters) above sea level.

Much of the following discussion is taken from Schumacher and others (1987), and Shrake and others (1988), and Shrake (1992). Paleoenvironmental interpretations are from Schumacher and Ausich (1983).

The emergency spillway is 2,667 ft (813.1 meters) long and 450 ft (137.2 meters) wide, giving the spillway an area of over 1.2 million square feet (112,000 square meters), which is equivalent to more than 27 acres (11 hectares).

The exposure below the dam, called Caesar Creek gorge (OH-WA-0002), is reached via the winding road 0.8 mile (1.3 km) long descending to a parking lot in the bottom of the deep valley of Caesar Creek itself; the outcrop is at the lefthand (north) end of the bottom of the dam, 0.1 mile (0.2 km) from that lot. The shale bank there is at 39°29'05"N, 84°03'49"W (4374458 m N, 752555 m E, UTM zone 16). Its




FIGURE 13-1.—Location of the Caesar Creek sites. The spillway is indicated by the southern arrow, the shale bank below the dam by the northern arrow, and the Visitors Center by the solid triangle. Oregonia, Ohio, 7.5-minute quadrangle.

base is at an elevation of 755 ft (230.1 meters).

As mentioned above, even though this locality is within a state park, the general public is permitted to collect fossils from the emergency spillway, although excavation is not allowed. The only requirement is to sign a permit form at the Visitors Center each time one wishes to collect. For further information and current rules, write or call:

Caesar Creek Lake Visitors Center
U.S. Army Corps of Engineers
4020 N. Clarksville Road
Waynesville, Ohio, 45068
513-897-1050




FIGURE 13-2.—Diagrammatic road net in vicinity of the Caesar Creek spillway and dam (from Shrake, 1992).

Caesar Creek Lake Ranger Station
(same address)
513-897-1738

Caesar Creek State Park Office
Ohio Department of Natural Resources
8570 E. State Route 73
Waynesville, Ohio, 45068
513-897-3055.

STRATIGRAPHY

The Caesar Creek localities have been described by Schumacher and Ausich (1983), Schumacher and others (1987), Shrake and others (1988), and Shrake (1992). Further discussion of the formations exposed here can be found in Schumacher (paper 16 in this volume). The stratigraphic units exposed are illustrated in figure 13-3.

At the base of the dam approximately 10 ft (3.3 meters) of

the Arnheim Formation and about 40 ft (12.2 meters) of the Waynesville Formation are exposed. This outcrop is excellent for studying the Arnheim Formation's nodular-bedded, rubbly weathering nature and its contact with the overlying Waynesville Formation.

The Arnheim Formation is traditionally separated into two members, the basal Sunset and overlying Oregonia. Only about 10 ft (3.3 meters) of the Oregonia is exposed at the dam. The Oregonia Member averages 60 percent shale and 40 percent limestone. The limestone in the Oregonia Member occurs as thin (1.5 inches/3.8 cm), discontinuous, wavy beds and as nodules (1 to 3 inches/2.5 to 7.6 cm in diameter), intermixed and interbedded with fissile shale layers. The member was named for exposures near Oregonia, Ohio, 3 miles (4.8 km) southwest of the park. Field mapping in Warren County indicates the uppermost portion of the Oregonia Member is a consistent waterfall former and tends to have a rubbly weathering pattern.

Approximately 40 ft (12.2 meters) of the basal Waynes-




FIGURE 13-3.—Stratigraphic sections exposed at Caesar Creek dam. Vertical numbers indicate feet above sea level. 1 = Waynesville Formation poorly exposed above bench from about 784 ft (239.0 meters) to the top of the outcrop at about 811 ft (247.2 meters); 2 = Clarksville Road at approximately 883 ft (269.1 meters) (modified from Shrake and others, 1988, p. 42).

ville Formation overlies the Arnheim Formation at the dam section (fig. 13-3). In northeastern Warren County, the Waynesville averages about 110 ft (34 meters) in thickness and is 74 percent shale and 26 percent limestone. Thick (5 inches/12.7 cm), platy-parted shale layers are interbedded with thin (2 inches/5 cm), planar, moderately continuous limestone layers. The type area for the Waynesville Formation includes the outcrops of the unit in the vicinity of Waynesville, Ohio.

At the emergency spillway, the floor and walls expose the sequence from the top of the Waynesville Formation to the basal portion of the Whitewater Formation. The formations are distinguished by the ratio of limestone to shale, the style of bedding, and the thickness of the individual layers. Examination of the bedding styles visible in the spillway walls is encouraged; however, climbing on the vertical walls is strictly prohibited by the Corps of Engineers.

The upper 20 ft (6 meters) of the Waynesville is exposed along the southwestern end of the emergency spillway but is covered by colluvium. These strata are predominantly calcareous blue-gray shales with fossiliferous, gray limestone lenses (packstones). This portion of the spillway section is generally interpreted to have been deposited in a lower energy environment below normal wave base with periodic disturbances by storms. Supporting evidence for a lower energy environment consists of a lack of wave-rippled limestones, shingled brachiopods, broken and abraded fossils, and cross-bedded limestones. However, the presence of graded beds, partly winnowed limestones, and well-preserved echinoderms indicates that high-energy sedimentologic events such as storms influenced the unit periodically. Kreisa and others (1981) reported evidence of storms in Caesar Creek limestones.

The Liberty Formation is exposed in the emergency spillway floor and walls (fig. 13-3). The formation averages 27 ft (8.2 meters) thick and 57 percent shale and 43 percent limestone (packstones and grainstones). The Waynesville Formation and the Liberty Formation are distinguished by the percentage of limestone (26 vs. 43 percent) and by thickness of the limestone layers (2 inches/5 cm vs. 4 inches/10 cm).

The Liberty Formation represents a transitional zone between the lower energy Waynesville below and the higher energy Whitewater above. This interpretation is based on the increased number of limestone beds that have been winnowed of most, if not all, of the silt- and clay-size material, the presence of rare wave-rippled limestones and shingled brachiopods, and an increase in abraded and broken fossils from the Waynesville.

Although the Whitewater Formation averages 66 ft (20 meters) in thickness in northeastern Warren County, only the basal 30 ft (9.1 meters) is exposed along the emergency-spillway wall and the overlying slope (fig. 13-3). The Whitewater averages 52 percent shale and 48 percent limestone in northeastern Warren County. However, the basal section exposed in the park is 45 percent shale and 55 percent limestone. Limestone layers in the Whitewater Formation are thin (2 inches/5 cm), wavy bedded, discontinuous, and argillaceous. Shale layers are thin and fissile.

Higher energy conditions are inferred for the Whitewater on the basis of abundant completely winnowed limestones (grainstones), lower faunal diversity, and an increase in encrusting bryozoans, cross-laminations, and abraded fossils.

No Saluda dolomite has been mapped in Warren County. Its approximate horizon presumably would be at the top of

the spillway section, based on extrapolations of thickness trends from the west.

PALEONTOLOGY

Some of the more common fossils at Caesar Creek spillway include the rugose coral *Grewingia canadensis*; the bryozoans *Parvohallopora subnodosa* and *Constellaria polystomella*; the brachiopods *Platystrophia*, *Hebertella*, *Onniella meeki*, *Rafinesquina alternata*, *Hiscobeccus capax* (formerly referred to *Lepidocyclus*), *Strophomena*, *Zygospira modesta*, and *Leptaena richmondensis*; the pelecypods *Ambonychia radiata* and *Cymatonota typicalis*; the nautiloid cephalopod *Treptoceras duseri*; and the gastropods *Cyclonema bilix* and *Trochonema*. Less common are the trilobites *Flexicalymene meeki*, *Isotelus*, *Acidaspis onealli*, *Chasmops breviceps*, and *Ceraurinus icarus* and the echinoderms *Cupulocrinus*, *Cincinnatiocrinus*, *Dendrocrinus*, *Gaurocrinus*, *Xenocrinus*, *Lichenocrinus*, and *Petraster*. In addition to providing a classic locality to collect body fossils, the emergency spillway provides an excellent opportunity to study many of the trace fossils of the local Cincinnati rocks, such as *Rusophycus*, *Chondrites*, and *Diplocraterion*. Examples of these fossils can be seen on the weathered limestone slabs in the talus along the southern wall of the spillway. Well-preserved echinoderms from several horizons in the spillway have been described by Schumacher and Ausich (1983).

The Whitewater limestones and shales here yield abundant robust branching bryozoans, especially at the top of the main road cut on the southeastern side of the spillway. The Whitewater is generally interpreted to be of very shallow water origin (Cuffey, paper 2 in this volume, fig. 2-4). A modern analog for abundant ramose bryozoans on extremely shallow carbonate bottoms can be seen off South Cat Cay in the Bahamas (Hoffmeister and others, 1967, p. 180, 183, 185; Cuffey and Fonda, 1976); a comparable example on terrigenous sediments was examined in 1977 by Cuffey off St. Theresa Beach in the panhandle of Florida. Branching bryozoans can also be important in deeper waters, at mid-shelf depths, as documented by Lagaaij and Gautier (1965, p. 52, 54) and Schopf (1969, p. 237, 240), but such occurrences are clearly different from these Whitewater deposits. Abundant branching bryozoans thus are not simplistically indicative of deeper waters (as commonly seems to be presumed), but can be found in certain shallow habitats as well.

Such great bryozoan abundance warrants investigation; hence, approximately 200 colony fragments were examined and identified (Rockwell and Cuffey, 1996). Thirty-seven species were found in the highest exposed Whitewater strata here, a comparatively high biodiversity for this phylum at a single locality within the type-Cincinnati. Bryozoans identified (* denotes species constituting 5 percent or more of this bryozoan fauna) included many trepostomes: the monotypid *Cyphotrypa madisonensis*; the monticuliporids *Gortanipora bassleri*, *Homotrypa alta*, *H. austini*, *H. cincinnatiensis*, *H. communis*, *H. creditensis*, *H. cylindrica*, *H. nitida*, *H. nodulosa*, *H. richmondensis*, *H. wortheni*, *Homotrypella rustica*, and *Peronopora pachymura*; the amplexoporids *Amplexopora variabile* and *Rhombotrypa quadrata**; the heterotrypids *Heterotrypa cystata*, *H. inflecta*, *Dekayia semipilaris*, and *Leptotrypa minima*; the halloporids *Hallopora congrua**, *H. dalei*, *H. elegantula*, *H. nodulosa*, *Parvohallopora ramosa**, *P. rugosa*, *P. subnodosa**, *P. subplana*, and *P. wesenbergiana*; the trematopod

*Batostoma varians**; and the batostomellids *Batostomella meeki*, *Bythopora dendrina**, and *Eridotrypa mutabilis*. Three bifoliate cryptostomes (*Arthropora cleavelandi**, *Crateriopora lineata*, and *Dicranopora emacerata*) and one paleotubuliporine cyclostome (*Cuffeyella arachnoidea*) are also present (Taylor and Wilson, 1996). (*Parvohallopora ramosa* and *P. rugosa* may be conspecific; their distributions do overlap, but they do not coincide. It is important to record each variant wherever it turns up, because that eventually may tell us what each was. For example, perhaps the variants are a reflection of local variation in microhabitat.)

Before the Caesar Creek spillway was excavated, the bryozoans of the Whitewater Formation had been studied by Cumings (1908) and by Utgaard and Perry (1964). Thus, this fauna has been sampled extensively, but independently three times. Approximately two-thirds of the species found in each study appear in one or both of the others, indicating substantial but not complete overlap among these three investigations. Moreover, in order to assess possible sampling effects upon the understanding of the local faunal composition in these fossils, the Caesar Creek bryozoans were collected as two separate equal-sized subsamples. Again, about two-thirds of Caesar Creek's total of 37 species occur in each subsample, but only half are present in both; however, if the numbers of specimens identified are counted instead of just the species, a much greater overlap (about 80 percent) is seen. The seven most common Caesar Creek species all appear in both subsamples, most in Cumings (1908), and half in Utgaard and Perry (1964).

As noted above, the bryozoan-bearing beds at Caesar Creek are chronologically correlative with the Saluda Dolomite to the west, the bryozoans of which also have been analyzed recently (Butler and Cuffey, 1994, 1996; papers 6 and 14 in this volume). There is a total of 42 bryozoan species at this horizon across the Cincinnati region. The high diversity at Caesar Creek, as contrasted with the lower diversity in the Saluda itself, nicely fits with the paleoenvironmental interpretation of a shallow, open-marine shelf for this locality (Cuffey, Butler, and Rockwell, 1996).

REFERENCES CITED

- Butler, K. L., and Cuffey, R. J., 1994, Bryodiversity and paleostress in the Saluda dolomitic shelf lagoon (latest Ordovician, southeastern Indiana) (abs.): Geological Society of America Abstracts with Programs, v. 26, no. 5, p. 7.
- Butler, K. L., and Cuffey, R. J., 1996, Reduced bryozoan diversity and paleoenvironmental stress in the Saluda Dolomite (uppermost Ordovician, southeastern Indiana), in Gordon, D. P., Smith, A. M., and Grant-Mackie, J. A., eds., Bryozoans in space and time: Wellington, New Zealand, Proceedings of the 10th International Bryozoology Conference (1995), National Institute of Water and Atmospheric Research, Ltd., p. 55-61.
- Cuffey, R. J., Butler, K. L., and Rockwell, L. A., 1996, Bryozoan diversity and salinity in the uppermost Ordovician (Saluda and Whitewater Formations) across southeastern Indiana and southwestern Ohio (abs.): Geological Society of America Abstracts with Programs, v. 28, no. 6, p. 34-35.
- Cuffey, R. J., and Fonda, S. S., 1976, "Giant" *Schizoporella*—sedimentologically important cheilostome bryozoans in Pleistocene and Recent carbonate environments of Bermuda, the Bahamas, and Florida (abs.): Geological Society of America Abstracts with Programs, v. 8, no. 4, p. 474-475.
- Cumings, E. R., 1908, The stratigraphy and paleontology of the Cincinnati Series of Indiana: Indiana Department of Geology and Natural Resources, Thirty-second Annual Report for 1907, p. 605-1189.
- Hoffmeister, J. E., Stockman, K. W., and Multer, H. G., 1967, Miami Limestone of Florida and its Recent Bahamian counterpart: Geological Society of America Bulletin, v. 78, p. 175-190.
- Kreisa, R. D., Dorobek, S. L., Accoti, P. J., and Ginger, E. P., 1981, Recognition of storm-generated deposits in the Cincinnati Series, Ohio (abs.): Geological Society of America Abstracts with Programs, v. 13, no. 6, p. 285.
- Lagaaij, R., and Gautier, Y. V., 1965, Bryozoan assemblages from marine sediments of the Rhône delta, France: Micropaleontology, v. 11, p. 39-58.
- Rockwell, L. A., and Cuffey, R. J., 1996, Great bryozoan diversity and sampling in the uppermost Ordovician (Whitewater Formation) at Caesar Creek (southwestern Ohio) (abs.): Geological Society of America Abstracts with Programs, v. 28, no. 6, p. 62.
- Schopf, T. J. M., 1969, Paleocology of ectoprocts (bryozoans): Journal of Paleontology, v. 43, p. 234-244.
- Schumacher, G. A., and Ausich, W. I., 1983, New Upper Ordovician echinoderm site: Bull Fork Formation, Caesar Creek Reservoir (Warren County, Ohio): Ohio Journal of Science, v. 83, p. 60-64.
- Schumacher, G. A., Shrake, D. L., Swinford, E. M., Brockman, C. S., and Wickstrom, L. H., 1987, Stratigraphy and depositional environments of the Cincinnati Group of southwestern Ohio: Field trip guidebook for 16th Eastern Section meeting of American Association of Petroleum Geologists, Ohio Geological Society, 73 p.
- Shrake, D. L., 1992, Excursion to Caesar Creek State Park in Warren County, Ohio: a classic Upper Ordovician fossil-collecting locality: Ohio Division of Geological Survey Guidebook 12, 18 p.
- Shrake, D. L., Schumacher, G. A., and Swinford, E. M., 1988, Field guidebook to the stratigraphy, sedimentology, and paleontology of the Upper Ordovician rocks in southwestern Ohio: Guidebook for Fifth Midyear Meeting of Society of Economic Paleontologists and Mineralogists, Ohio Division of Geological Survey, 81 p.
- Taylor, P. D., and Wilson, M. A., 1996, *Cuffeyella*, a new bryozoan genus from the Late Ordovician of North America, and its bearing on the origin of the post-Paleozoic cyclostomes, in Gordon, D. P., Smith, A. M., and Grant-Mackie, J. A., eds., Bryozoans in space and time: Wellington, New Zealand, Proceedings of the 10th International Bryozoology Conference (1995), National Institute of Water and Atmospheric Research, Ltd., p. 351-360.
- Utgaard, J. E., and Perry, T. G., 1964, Trepostomatous bryozoan fauna of the upper part of the Whitewater Formation (Cincinnati) of eastern Indiana and western Ohio: Indiana Geological Survey Bulletin 33, 111 p.

Addresses of authors: Gregory A. Schumacher, Douglas L. Shrake, and E. Mac Swinford, Ohio Department of Natural Resources, Division of Geological Survey, 4383 Fountain Square Drive, Columbus, OH 43224-1362; and Lynn A. Rockwell and Roger J. Cuffey, Department of Geosciences (Deike Bldg.), Pennsylvania State University, University Park, PA 16802.

14. ARNHEIM THROUGH UPPER WHITEWATER STRATA AT THE MADISON-INDIANA ROUTE 56 ROAD CUTS (UPPER ORDOVICIAN, SOUTHEASTERN INDIANA)

by
Donald E. Hattin and Roger J. Cuffey

SIGNIFICANCE

The road cuts along Indiana Route 56, where the road ascends the valley wall up from the Ohio River just west of Madison, expose the longest continuous section of Ordovician rocks in Indiana (Hattin and others, 1961, p. 328). Although one or two more recent cuts now may exceed this section somewhat in length, it is still well exposed and abundantly fossiliferous and continues to be an important locality for examining later Cincinnati formations in the region. This section can be compared profitably with that exposed in the relatively recent cuts along U.S. Route 421 northeast of Madison (IN-JE-0001, IN-JE-0002, and IN-JE-0003; see Hay, Totten, and Cuffey, paper 6 in this volume).

LOCATION

This locality (IN-JE-0004) consists of four closely spaced road cuts along Indiana Route 56 (formerly also Indiana Route 62) at various distances east or downgrade from its junction with the present Indiana Route 62 about 4 miles (6.4 km) west of downtown historic Madison, Jefferson County, Indiana (fig. 14-1):

1. 0.3-0.7 mile (0.5-1.1 km) east of intersection,
2. 0.8 mile (1.3 km) east of intersection,
3. 0.9 mile (1.4 km) east of intersection, and
4. 1.0-1.2 miles (1.6-1.9 km) east of intersection.

The most conspicuous cut is the high wall exposing the Saluda Limestone at the sharp curve 0.5-0.6 mile (0.8-1.0 km) east of that junction. The cuts occupy the NE corner sec. 6 and N $\frac{1}{2}$ N $\frac{1}{2}$ NW $\frac{1}{4}$ sec. 5, T. 3 N., R. 10 E., and S $\frac{1}{2}$ SW $\frac{1}{4}$ SE $\frac{1}{4}$ sec. 31, T. 4 N., R. 10 E., on the Madison West, Indiana-

Kentucky, 7.5-minute quadrangle. The high cut through the Saluda is situated at 38°44'12"N and 85°26'28"W (4288480 m N, 635461 m E, UTM zone 16).

STRATIGRAPHIC SECTION

The following section (fig. 14-2) was measured in 1957 by Donald E. Hattin and published (as Stop 8 of field trip 9) in Hattin and others (1961, p. 343-347). Only the most abundant fossils or fossil groups are listed, and names of taxa have been updated, where appropriate (R. A. Davis, 1992, written commun.).

Middle and Lower Silurian units, 52.6 ft (16.0 meters) exposed

UPPER ORDOVICIAN	feet (meters)
<u>(Upper) Whitewater Formation</u>	
dd. Limestone, dark gray, fine to medium grained, with brachiopods, gastropods, cephalopods, ostracodes, and stromatoporoids, and shale, dark gray, calcareous, laminated (several stromatoporoid heads are erosionally truncated)	5.2 (1.6)
Total thickness of Whitewater	5.2 ft (1.6 m)

Saluda Limestone

cc. Dolomitic limestone, light gray to greenish gray, fine to coarse grained, local geodes, many calcite-filled worm? burrows, contains ramose bryozoans (all those sectioned being *Batostomella*)




FIGURE 14-1.—Location of the Indiana Route 56 road cuts; the western and eastern arrows indicate the extent of the cuts and the middle arrow indicates the high cut through the Saluda Limestone. Madison West, Indiana-Kentucky, 7.5-minute quadrangle.




FIGURE 14-2.—Graphic section of strata exposed in the Indiana Route 56 road cuts west of Madison, Indiana (from Hattin and others, 1961, p. 328, fig. 11); current Indiana Geological Survey usage would label beds A-R as the Dillsboro Formation.

articulate brachiopods, ramose and massive bryozoans, and pelecypods	5.9	(1.8)	h. Clay shale, weathered moderate yellowish brown, moderately calcareous, with four thin limestones, light brownish gray, fine to medium grained, biofragmental; unit contains gastropods, <i>Onniella</i> , other articulate brachiopods, and ramose bryozoans	1.7	(0.5)
p. Limestone, medium gray, medium to coarse grained, biofragmental, and shale, greenish gray, very calcareous; unit contains gastropods, pelecypods, articulate brachiopods, ramose bryozoans, and <i>Grewingkia</i>	10.6	(3.2)	g. Limestone, medium gray, medium to coarse grained, biofragmental, with thin shale partings; unit contains <i>Onniella</i> , other brachiopods, and gastropods	5.1	(1.6)
o. Clay shale, olive gray, gritty, very calcareous, with thin layers of calcareous claystone; unit contains brachiopods and bryozoans	2.5	(0.8)	f. Clay shale, light olive gray, moderately calcareous, abundant fossil fragments, with beds of limestone like those in unit "g"; unit contains <i>Onniella</i> , other brachiopods, pelecypods, gastropods, trilobites, and ramose bryozoans	7.2	(2.2)
n. Limestone, medium gray, fine to coarse grained, biofragmental, and lime mudstone, fine grained, with few fossils; unit has thin olive-gray clay-shale interbeds and numerous brachiopods, ramose and massive bryozoans, and <i>Grewingkia</i>	3.4	(1.0)	e. Clay shale, dusky yellow, moderately calcareous, few fossils, with nodular claystone in lower part and thin limestone beds as in unit "g"	8.6	(2.6)
Total thickness of Liberty	42.4 ft	(12.9 m)	d. Limestone, medium to dark gray, coarse grained, biofragmental, with interbeds of calcareous, gritty, clay shale	2.8	(0.9)
<u>Waynesville Formation</u> - upper limit indefinite			c. Clay shale, weathered dusky yellow, blocky, moderately calcareous, with two thin limestones and a few lime-mudstone nodules; unit contains crinoid columnals	7.7	(2.3)
m. Clay shale, medium gray to light olive gray, very calcareous, and limestone, light brownish gray, medium to coarse grained, biofragmental; unit contains <i>Grewingkia</i> , <i>Isotelus</i> , massive and ramose bryozoans, articulate brachiopods, and gastropods	10.3	(3.1)	Total thickness of Waynesville	68.9 ft	(21.0 m)
l. Clay shale, olive gray to greenish gray, gritty, very calcareous, completely transitional to argillaceous, highly fossiliferous limestone; unit contains some hard, medium-gray biofragmental limestone; wide variety of fossils litter slope, abundant <i>Onniella</i>	12.2	(3.7)	<u>Arnheim Formation</u> - upper limit indefinite		
k. Limestone, medium dark gray, fine to coarse grained; unit contains articulate brachiopods and ramose and massive bryozoans	0.7	(0.2)	b. Limestone, medium gray to light brownish gray, coarse grained, biofragmental, with nearly equal quantity of clay shale, greenish gray, highly calcareous; unit contains profusion of <i>Rafinesquina</i> , other brachiopods, trilobites, bryozoans, crinoid columnals, pelecypods, <i>Cyclonema</i> , and other gastropods	17.2	(5.2)
j. Clay shale, greenish gray, somewhat gritty, very calcareous, with numerous layers of limestone, medium gray, medium to coarse grained, biofragmental; unit contains abundant <i>Onniella</i> , other articulate brachiopods, ramose and massive bryozoans, gastropods, and <i>Grewingkia</i>	9.5	(2.9)	a. Clay shale, greenish gray, moderately calcareous, with two thick beds of limestone, light to dark gray, coarse grained, biofragmental, with numerous thin layers and nodules of limestone, argillaceous, fine to medium grained; unit not so fossiliferous as unit "b"; unit contains articulate brachiopods, pelecypods, and ramose bryozoans	10.6	(3.2)
i. Limestone, medium dark gray, medium to coarse grained, biofragmental, with thin calcareous shale breaks; unit contains <i>Onniella</i> , other articulate brachiopods, pelecypods, and massive bryozoans	3.1	(0.9)	Total thickness of exposed Arnheim	28.8 ft	(8.8 m)
			Total thickness of measured section	253.9 ft	(77.4 m)

STRATIGRAPHIC NOTES

The following comments on Richmondian stratigraphy in southeastern Indiana are mainly those written by Hattin for a field trip for the 1961 Cincinnati meeting of the Geological Society of America (Hattin and others, 1961, p. 328-331, Stop 8); taxonomic names have been updated (R. A. Davis, 1992, written commun.).

The lowest part of the section, separated from the upper part by a 25-foot (7.6-meter) covered interval, is exposed at the foot of the hill beside a small cafe on the north side of the road. Upper Maysvillian beds at the cafe are not included in the measured section. Strata of the lower part of the Richmond group consist of a sequence of alternating fossiliferous limestones and generally less fossiliferous shales. These beds have been classed, in upward order, as Arnheim, Waynesville, and Liberty formations wholly on paleontological criteria and are not formations in the restricted sense of the term. Arnheim guide fossils include *Homotrypa bassleri* and *Retrosirostra carleyi* in combination with conspicuous numbers of *Cyclonema bilix* and *Flexicalymene meeki*. The Arnheim contains a proportionately greater quantity of limestone than the overlying Waynesville. Chief guide fossil of the Waynesville Formation is *Onniella meeki*, which abounds on some bedding planes. Species that occur in conspicuous numbers in the formation are *Leptaena richmondensis*, *Zygospira modesta*, and *Thaerodonta clarksvillensis*. *Glyptorthis insculpta* is common in a zone at the top of the formation that was classed as Liberty by Cumings (1908). Waynesville strata are generally more shaly than beds above or below, but sharp contacts cannot be drawn on a lithologic basis.

In the Liberty Formation, *Rhombotrypa quadrata*, a rare bryozoan, is an abundant and characteristic species (Cuffey and Perry, 1964). *Hiscobeccus capax* (formerly referred to the genus *Lepidocyclus*), *Strophomena planumbona*, and *Grewingia canadensis* are more abundant in the Liberty than in adjacent strata.

Ordovician formations above the Liberty can be distinguished readily on lithologic character alone. In the Madison, Indiana, area, the Saluda is characteristically dolomitic, is chiefly medium bedded to very thick bedded, and contains abundant quartz silt that dominates insoluble residues of most beds. The only characteristic Saluda fossils are *Favistina stellata* and *Tetradium approximatum*. The former generally is confined to the lower, and the latter to the upper, of two biostromes that lie near the base of the formation. For many years the contact between the Liberty and Saluda formations of the Madison area has been defined as the base of the lower of the two coral biostromes. However, such a contact segregates as Liberty much dolomitic rock that is related genetically to the Saluda environment of deposition. Although the biostromes provide a convenient and natural base for the Saluda north of Madison, southward from Madison these coral beds terminate laterally well above the base of a section of thick-bedded dolomitic limestone that is typical of the Saluda. The contact between the Saluda is diachronous and descends the section southward from Madison. If the Saluda is to be considered a lithogenetic unit, then it is necessary to place the contact at the base of these dolomitic strata, for this is the most reliable and logical boundary between Liberty and Saluda beds in the southern part of the Indiana outcrop. A thin, widespread, dark-gray glauconitic shale (unit Y; fig. 14-2) serves as a key marker bed in demonstrating the contact relationship between

the two formations and serves as the datum for the cross section in figure 14-3. At Marble Hill, at the south end of the cross section, 34 ft (10.4 meters) of thick-bedded, not visibly fossiliferous dolomitic limestone lies below the shale marker bed. At Madison (3, Clifty Power Plant, on fig. 14-3), 25.8 ft (7.9 meters) of fossiliferous dolomitic limestone and mudstone lies beneath the key bed. At Versailles, only 8.1 ft (2.5 meters) of dolomitic limestone, constituting the *Tetradium* biostrome, lies beneath the key bed. From south to north the base of the Saluda grades into progressively higher nondolomitic and highly fossiliferous beds of the upper Liberty formation.

Lying above the key shale bed are laminated, thick-bedded to very thick bedded dolomitic limestones that thin progressively northward from Jefferson County, Indiana. Ripple marks and mud cracks are common in the laminated beds, but Hattin has seen no fossils in these beds. The lower part of the laminated beds passes northward from Madison into a thin sequence of alternating shales and medium- to thick-bedded limestones. Highest of the widespread Saluda marker beds is a resistant bed, or group of beds, that forms the lip of many high waterfalls in the area between Bethlehem and Versailles, Indiana. Vertical cylinders of calcite in this bed are probably spar-filled burrows of worms. The *Tetradium* biostrome, laminated beds, and "falls lip" bed of the Madison area constitute a bundle of rock units that can be traced farther northward than any other Saluda beds. Above the "falls lip" limestone bed is 15.5 ft (4.7 meters) of thick-bedded to very thick bedded dolomitic limestone that passes northward, by lateral gradation, into strata typical of the Whitewater formation. The Whitewater-Saluda contact, like the Liberty-Saluda contact, is diachronous, transcending time lines from north to south.

The Saluda is mostly a poorly fossiliferous, chiefly dolomitic limestone wedge that lies between, and is partly gradational with, highly fossiliferous, nondolomitic strata. North of Versailles, beds assigned to the Saluda constitute an essentially nondolomitic facies that can be traced as far north as Hamburg, beyond which the formation is unrecognizable. Near Laurel, Indiana, 9 miles (14.5 km) north of Hamburg, beds in the position of the Saluda are a highly fossiliferous, thin-bedded shale and limestone sequence that contains an abundance of stromatoporoids and compound corals.

Saluda features such as laminated bedding, oscillation ripple marks of short length and low amplitude, mud cracks, and general lack of fossils suggest that the formation is the product of quiet, shallow-water deposition, possibly in a stagnant or a highly saline environment. In contrast, biofragmental limestones above and below the Saluda represent high-energy, well-oxygenated environments where normal-marine faunas flourished. The association of biostromal units with lower Saluda beds through much of the Indiana outcrop, and abundance of biostromes in the vicinity of the northern terminus of the formation, suggest a relationship between the biostromes and the environment in which the Saluda facies was deposited. Although much more field, subsurface, and laboratory work needs to be accomplished, we tentatively suggest that Saluda deposition was linked with lagoonal conditions that existed behind low but extensive shoal areas on which banks of *Favistina*, *Tetradium*, and stromatoporoids grew extensively. (See Hatfield, 1968, and Larabee, 1994, for further elaboration of these suggestions.)

The lower strata of the Saluda here contain numerous




FIGURE 14-3.—South-north cross section in southeastern Indiana showing the stratigraphic relations of the Saluda Limestone (from Hattin and others, 1961, p. 330, fig. 12). The Madison-Indiana Route 56 outcrop is equivalent to 3, Clifty Power Plant.

bryozoans, but, in contrast to the diversity in the rest of the type-Cincinnatian, only three species are represented here, *Homotrypa cincinnatiensis*, *Rhombotrypa quadrata*, and *Heterotrypa subfrondosa* (Butler and Cuffey, 1994, 1996). The Saluda correlates with the bryozoan-rich mid-Whitewater beds at Caesar Creek (Rockwell and Cuffey, 1996; paper 13 in this volume). The Saluda's reduced bryozoan diversity, in contrast to the high diversity at Caesar Creek, reflects well the hypersaline shelf-lagoon paleoenvironment inferred for the Madison area at this horizon (Cuffey, Butler, and Rockwell, 1996).

Whitewater strata are generally thin and very irregularly bedded as compared with underlying Saluda beds. Although *Rhynchotrema dentatum* has been cited as a characteristic Whitewater fossil, specimens are very rare south of Hamburg, Indiana. *Platystrophia acutilirata* is a common species in the northern part of the Indiana outcrop. The stromatoporoid *Labechia huronesis* is very common in the Whitewater and forms local biostromes. Algal biostromes are well developed in the formation near Richmond, Indiana. At most exposures, the Whitewater contains large numbers of the smooth-shelled ostracode *Leperditia*. The Whitewater thins southward along the Indiana outcrop by progressive gradation of the lower beds into the upper Saluda and seemingly by regional erosional truncation of

the upper beds. The unit cannot be recognized in Indiana sections south and west of Harts Falls (fig. 14-3). The name "Hitz bed" (Foerste, 1903) has been applied to the thin but highly fossiliferous Whitewater section in the area around Madison, Indiana.

Northward from northern Ripley County, the Elkhorn formation, youngest of the Richmondian formations in the type area, lies between the Whitewater and the Brassfield (Silurian) formations. We tentatively conclude from field studies that the Elkhorn terminates southward by lateral gradation into the upper Whitewater formation and by erosional truncation of the upper beds. Little is known or published concerning the Elkhorn fauna, but the entire section is exposed in several places south and southwest of Richmond, Indiana.

REFERENCES CITED

- Butler, K. L., and Cuffey, R. J., 1994, Bryodiversity and paleostress in the Saluda dolomitic shelf lagoon (latest Ordovician, southeastern Indiana) (abs.): Geological Society of America Abstracts with Programs, v. 26, no. 5, p. 7.
- Butler, K. L., and Cuffey, R. J., 1996, Reduced bryozoan diversity and paleoenvironmental stress in the Saluda Dolomite (uppermost Ordovician, southeastern Indiana), in Gordon, D.

- P., Smith, A. M., and Grant-Mackie, J. A., eds., Bryozoans in space and time: Wellington, New Zealand, Proceedings of the 10th International Bryozoology Conference (1995), National Institute of Water and Atmospheric Research, Ltd., p. 55-61.
- Cuffey, R. J., Butler, K. L., and Rockwell, L. A., 1996, Bryozoan diversity and salinity in the uppermost Ordovician (Saluda and Whitewater Formations) across southeastern Indiana and southwestern Ohio (abs.): Geological Society of America Abstracts with Programs, v. 28, no. 6, p. 34-35.
- Cumings, E. R., 1908, The stratigraphy and paleontology of the Cincinnati Series of Indiana: Indiana Department of Geology and Natural Resources, Thirty-second Annual Report for 1907, p. 605-1189.
- Foerste, A. F. 1903, The Richmond Group along the western side of the Cincinnati Anticline in Indiana and Kentucky: American Geologist, v. 31, p. 333-361.
- Hatfield, C. B., 1968, Stratigraphy and paleoecology of the Saluda Formation (Cincinnati) in Indiana, Ohio, and Kentucky: Geological Society of America Special Paper 95, 34 p.
- Hattin, D. E., Nosow, Edmund, Perkins, R. D., Stumm, E. C., Mound, M. C., and Utgaard, John, 1961, Field excursion to the Falls of the Ohio, field trip 9: Guidebook for Field Trips, Cincinnati Meeting, 1961, Geological Society of America, p. 295-350.
- Larabee, J. H., 1994, Stratigraphy and paleoecology of coralline biostromes and patch reefs, Upper Ordovician of southeastern Indiana (abs.): Geological Society of America Abstracts with Programs, v. 26, no. 5, p. 49.
- Rockwell, L. A., and Cuffey, R. J., 1996, Great bryozoan diversity and sampling in the uppermost Ordovician (Whitewater Formation) at Caesar Creek (southwestern Ohio) (abs.): Geological Society of America Abstracts with Programs, v. 28, no. 6, p. 62.

Addresses of authors: Donald E. Hattin, Department of Geological Sciences, Indiana University, 1005 East 10th Street, Bloomington, IN 47405; and Roger J. Cuffey, Department of Geosciences (Deike Bldg.), Pennsylvania State University, University Park, PA 16802.

15. THE WHITEWATER FORMATION ON U.S. ROUTE 27 NEAR RICHMOND (UPPERMOST ORDOVICIAN, EAST-CENTRAL INDIANA)

by
Helen B. Hay and Roger J. Cuffey

SIGNIFICANCE

The part of the Whitewater Formation exposed in the U.S. Route 27 road cut about 1 mile (1.6 km) south of Richmond, Indiana (fig. 15-1), is significant because it provides a good and readily accessible exposure of the rubbly, thin-bedded facies that is most characteristic of the formation and because it furnishes some of the youngest, richly fossiliferous strata available within the type-Cincinnatian. Although the Whitewater Formation contains other facies (fig. 15-2), it is this particular lithology that constitutes the greatest volume of the formation and by which it is readily identified at other localities around the margin of the outcrop area. This part of the formation also is exposed and accessible along Sim Hodgkin Parkway (IN-WY-0003) in the Whitewater Gorge through Richmond, at Thistlethwaite Falls (IN-WY-0006) at the northern termination of the gorge on the West Fork of the East Fork of the Whitewater River, and in the Short Creek road cut (IN-WY-0002) just east of the Richmond-U.S. Route 27 cut.

The strata at this road cut are laterally equivalent to the Saluda Formation at Garr Hill (IN-FR-0003) (see Hay and Cuffey, paper 11 in this volume); the Saluda pinches out between Garr Hill and Richmond (Hay, 1981). Because the Saluda dolomite constitutes a lens within the Whitewater (see figs. 2-3 and 2-4 in Cuffey, paper 2 in this volume), there has been a tendency to divide the Whitewater into "lower" and "upper" parts, depending on whether the strata are below or above the Saluda or the Saluda horizon. As noted later in this paper, the Richmond-U.S. Route 27 road cut therefore includes rock of both the lower and upper portions of the Whitewater. Complicating this terminology, however, is recent recognition of additional Ordovician strata above the Elkhorn Shale (fig. 15-2; also see Schumacher, paper 16 in this volume), which was long considered the topmost Ordovician in the region.

The remainder of the Whitewater Formation is exposed at the other localities indicated on figures 15-1 and 15-2; the most significant localities are scattered outcrops along Elkhorn Creek between Straight Line Road (Pike) and Indiana Route 227. The creek is on private property and is not readily accessible; moreover, it generally is also heavily overgrown. The Ordovician/Silurian contact is exposed at the base of Elkhorn Falls (IN-WY-0004) just west of where Elkhorn Creek intersects Indiana Route 227; this site also is on private property. Relatively unfossiliferous shale, presumably but not certainly the Elkhorn, can be seen underneath the Silurian Brassfield Limestone forming the lip of Elkhorn Falls, but only when the creek waters are low.

All these localities mentioned constitute the Richmond composite section, which thus extends from the Liberty Member of the "Brookville formation" up through the Whitewater Formation—the uppermost Ordovician of this area—to the Silurian contact (fig. 15-2).

LOCATION

Lower and upper Whitewater strata are exposed in the road cut on U.S. Route 27, 1.2-1.3 miles (1.9-2.1 km) south-

west of its junction with Indiana Route 227 on the south edge of Richmond, Wayne County, Indiana (IN-WY-0001) (fig. 15-1). This road cut is located in NW¹/₄SE¹/₄SW¹/₄ sec. 17, T. 13 N., R. 1 W., on the Richmond, Indiana, 7.5-minute quadrangle, at 39°47'13"N and 84°54'10"W (4406000 m N, 679610 m E, UTM zone 16).

STRATIGRAPHY AND INTERPRETATION OF THE RICHMOND COMPOSITE SECTION

Figure 15-2 presents the stratigraphic units and lithofacies of the Richmond composite section and places the U.S. Route 27 road cut within that context. Lithofacies (fig. 15-3) and formation names are discussed by Hay (1981, and paper 17 in this volume). The Whitewater Formation, as defined here, includes all strata between the top of the Liberty Member of the "Brookville formation" and the Ordovician/Silurian boundary; the Elkhorn Shale here is considered to be a member of the Whitewater Formation. It makes sense to include strata above the Elkhorn Shale in the Whitewater Formation because facies 3c, the most typical Whitewater facies, occurs both above and below the shale. Except for the fact that the upper occurrence of facies 3c is cemented with dolomite, it is identical to that in the lower part of the Whitewater at the U.S. Route 27 locality. Strata that are lithologically transitional between Liberty and Whitewater are included in the Whitewater. The total thickness of the Whitewater Formation here is about 34 meters (113 ft). Descriptions of each locality in the composite section are provided by Hay (1981).

The Richmond composite section correlates with Garr Hill near Brookville, Indiana (see Hay and Cuffey, paper 11 in this volume). The *Thaerodonta* peak zone at about 7 meters (23 ft) above the base at Garr Hill correlates both faunally and lithologically with the base of the Richmond composite section. The Saluda Formation at Garr Hill should be equivalent to about the lower or middle part of the U.S. Route 27 road cut, although no Saluda facies occur at Richmond.

THE U.S. ROUTE 27 ROAD CUT

Figure 15-2 presents the stratigraphic units and lithofacies along U.S. Route 27 and figure 15-4 indicates the distribution of fossils in partially equivalent strata at Thistlethwaite Falls (IN-WY-0006) at the north edge of Richmond. Two facies, 3b and 3c (see fig. 15-3) are exposed at the U.S. Route 27 road cut. Facies 3c is the typical rubbly, thin-bedded argillaceous limestone containing irregular lumps of limestone separated by thin seams of limy terrigenous clay and silt. It can be difficult to tell limestones from shales, and accurate bed-by-bed measurements are virtually impossible. The overall insoluble content is about 33 percent, on the basis of acid treatment of bulk samples, and the carbonate content of "shales" is as much as 56 percent. It is this poor separation of siliciclastics and carbonates that is the distinctive characteristic of this facies and causes its properties. The rubbly weathering of the facies is caused by clay seams that coat and separate irregular limestone nodules. Very thin, irregular bedding is evident when the



FIGURE 15-1.—Location of the Richmond-U.S. Route 27 road cut (**US27** and arrow). Richmond, Indiana, and New Paris, Ohio—Kentucky, 7.5-minute quadrangles. Other localities (stars) that constitute the Richmond composite section are: **EC**, Elkhorn Creek (IN-WY-0008); **EF**, Elkhorn Falls (IN-WY-0004); **MF**, Middlefork Reservoir (IN-WY-0007); **SC**, Short Creek (IN-WY-0002); **SR**, Sub Run (IN-WY-0005); and **TF**, Thistlethwaite Falls (IN-WY-0006).

RICHMOND COMPOSITE




FIGURE 15-2.—Stratigraphy of the Richmond-U.S. Route 27 road cut in the context of the entire Richmond composite section: Elkhorn Creek (IN-WY-0008), Middlefork Reservoir (IN-WY-0007), Sub Run (IN-WY-0005), and Thistlethwaite Falls (IN-WY-0006). See Hay (1981) for detailed descriptions of these other sites. See figure 15-3 for description of lithofacies.

outcrop is viewed from a distance.

Thin sections of facies 3c show that these rocks have a higher proportion of whole fossils and large fossil fragments and a smaller percentage of sand- and silt-size allochems compared with facies such as 3a or 2a that have well-segregated limestone and shale beds. The thin sections also show abundant evidence of algae and micritic matrix compared with other facies.

Facies 3b at the top of the outcrop is best seen along the east side of the road. It is like facies 3c except that some of the limestones are thicker, more planar, and less argillaceous than those of facies 3c. Some of the limestones of this facies contain abundant algal oncolites.

These two facies also constitute most of the Bellevue Member of the Grant Lake Formation (Schumacher and others, 1991) in southern Indiana and Ohio and northern Kentucky. Although both facies are present, facies 3b is more common in the Bellevue, whereas facies 3c is more common in the Whitewater.

ENVIRONMENTS

The characteristics of facies 3c, in general, suggest that it probably formed in very shallow, but not strongly agitated water, perhaps the interior of a broad shoal. This interpretation is consistent with a number of properties of the rock: the poor separation of limestones and shales into discrete beds, the high proportion of large allochems, the low proportion of fine allochems, the abundant micrite, and the presence of algae. Facies 3b probably formed in a slightly higher energy environment marginal to the platform, which would explain the thicker, more planar, less clayey limestones produced by storm segregation of coarser allochems and fine siliciclastics into separate limestone and shale beds. This environment of slightly higher energy than facies 3c is consistent with

the presence of oncolites in facies 3b. The interpretation of a very shallow environment also is consistent with the stratigraphic relation of this facies to the Saluda Formation (Hay, paper 17 in this volume).

Both facies contain an abundant and diverse normal-marine fauna (fig. 15-4). Large trepostome bryozoans constitute the greatest volume of the rock and may have helped protect the sediment from reworking by waves and currents. Mollusks are probably next in abundance, particularly clams and snails that occur as steinkerns. The brachiopods are commonly articulated, which again points to a relatively quiet environment. Corals also are common.

ADDITIONAL LOCALITIES

Other localities that constitute the Richmond composite section (figs. 15-1 and 15-2; see Appendix A for descriptions) include:

Elkhorn Creek (IN-WY-0008).

Elkhorn Falls (IN-WY-0004).—Unfossiliferous gray shale (0.3 meter/1 ft) is exposed under low waterfalls just west of the Indiana Route 227 bridge over Elkhorn Creek. The falls are formed by the Silurian-age Brassfield Limestone. This shale was long thought to represent the Elkhorn Shale, but possibly may be an unnamed shale higher up, in the Brassfield Limestone. Otherwise, about 3 meters (10 ft) of Brassfield Limestone is exposed. Private property.

Middlefork Reservoir (IN-WY-0007).

Short Creek/Straight Line Pike (IN-WY-0002).—The same strata present at the Richmond-U.S. Route 27 locality also are exposed 0.7 mile (1.1 km) to the east, in the small road cut where Straight Line Pike (Road) crosses Short Creek. The road cut has yielded unusually large colonies, up to 15 cm (6 inches) across by 7 cm (3 inches) high, of the trepostome bryozoan *Monticulipora epidermata* Ulrich and Bassler, 1904, which are being studied by G. Hillmer and R. J. Cuffey.

Sim Hodgkin Parkway (IN-WY-0003).

Sub Run (IN-WY-0005).

Thistlethwaite Falls (IN-WY-0006).—Figure 15-4 shows the stratigraphic ranges of important fossils in this outcrop.

REFERENCES CITED

- Amsden, T. W., 1983, The Late Ordovician brachiopod genera *Lepidocyclus* and *Hiscobeccus*: Oklahoma Geological Survey Bulletin 132, p. 36-44.
- Hay, H. B., 1977, Field trip No. 1, Cincinnati stratigraphy from Richmond to Aurora, Indiana, in Pope, J. K., and Martin, W. D., eds., Field guidebook to the biostratigraphy and paleoenvironments of the Cincinnati Series in southeastern Indiana: Society of Economic Paleontologists and Mineralogists, Great Lakes Section, 7th Annual Field Conference, Miami University, p. 11-133 (reprinted in Pope and Martin, 1979; see Appendix B of this volume).
- Hay, H. B., 1981, Lithofacies and formations of the Cincinnati Series (Upper Ordovician), southeastern Indiana and southwestern Ohio: Ph.D. dissertation (unpub.), Miami University, 236 p.
- Schumacher, G. A., Swinford, E. M., and Shrake, D. L., 1991, Lithostratigraphy of the Grant Lake Limestone and Grant Lake Formation (Upper Ordovician) in southwestern Ohio: Ohio Journal of Science, v. 91, p. 56-68.

Addresses of authors: Helen B. Hay, deceased, formerly Department of Geology, Earlham College, Richmond, IN 47374; and Roger J. Cuffey, Department of Geosciences (Deike Bldg.), Pennsylvania State University, University Park, PA 16802.

Principal diagnostic characteristics of lithofacies	
Group 1 >70% shale	1a Well-bedded, mostly thin limestones and medium to thick, fissile to blocky shales. Includes massive units of 100% shale. Low-diversity fauna.
	1b Rubbly, shaly, nodular limestones; medium to thick shales with limy stringers and layers of fossils in shale matrix. May have some well-bedded limestones as in 1a. Low-diversity fauna. Most easily seen in cores.
Group 2 55-70% shale	2a Like 1a except has high diversity of fauna. Intermediate in shale percentage between 1a and 3a.
	2b Well-bedded limestones, rubbly nodular limestones, and limy shales. Allochems tend to be more finely broken than 2a. High-diversity fauna. Intermediate between 3a and 3c.
Group 3 <55% shale	3a Well-bedded, thin to medium limestones and thin to medium, fissile to blocky shales. High-diversity fauna.
	3b Well-bedded, thin to medium limestones with or without rubbly or nodular shales. High-diversity fauna. Intermediate between 3a and 3c.
	3c Thin-bedded, rubbly weathering, argillaceous limestones, very thin limy shales. High-diversity fauna.
	3d Limestones and shales as in 3a except >20% of limestones are fine-grained laminated burrowed, barren of large fossils; trails on tops. Grade to siltstones.
Group 4 Miscellaneous	4a High-angle cross-bedded calcarenites, bimodal dip. Little or no shale.
	4b Granule-size fossils in thick to massive beds; little or no shale. May be like 4a if seen in similar outcrops.
	4c Massive <i>Loxoplocus</i> limestone bed. Marble Hill Bed.
	4d Bioturbated light-gray limestone with low-diversity fauna of ostracodes, algae, snails, and <i>Tetradium</i> . Related to Saluda facies complex.




FIGURE 15-3.—Lithofacies characteristics. (see Hay, paper 17 in this volume).




FIGURE 15-4.—Fauna of Thistlethwaite Falls (IN-WY-0006), which is partially equivalent to the Richmond-U.S. Route 27 site (modified from Hay, 1977, p. I-16, fig. I-6). Note that *Lepidocyclus capax* was assigned to the genus *Hiscobeccus* by Amsden (1983).

16. A NEW LOOK AT THE CINCIANNIAN SERIES FROM A MAPPING PERSPECTIVE

by
Gregory A. Schumacher

INTRODUCTION

The interpretation of the geology and paleontology of the Upper Ordovician Cincinnati Series has been evolving for over a century and a half. Drake (1825) was the first to publish an account of geology in the Ohio River valley of southwestern Ohio. In the 170+ years following this publication, hundreds of investigations have been carried out on various geological and paleontological aspects of the rocks of the Cincinnati Series (see Appendix B at the end of this volume).

Most of these studies focused on a local geographical area or described and discussed the taxonomy of a group of fossils (for example, James, 1886; Nickles, 1902; Foerste, 1903; Caster and Kjellesvig-Waering, 1964; Ford, 1967; Warn and Strimple, 1977). In the last decade or two, a number of geological investigations have examined the regional geology of the Cincinnati Series using measured stratigraphic sections and lithofacies analysis (Hay, 1981; Tobin, 1982) or bedrock geologic mapping supplemented by measured stratigraphic sections and continuous cores (Weir and others, 1984; Schumacher, Swinford, and Shrake, 1991), or graptolite and conodont biostratigraphy (Bergström and Mitchell, 1990a, 1990b; Mitchell and Bergström, 1991).

This body of geological and paleontological literature provides an excellent framework to build upon in taking a new look at the geology of the Cincinnati Series using modern mapping and geological concepts. The Ohio Department of Natural Resources, Division of Geological Survey, in conjunction with The Ohio State University, have been re-examining the Cincinnati Series of southwestern Ohio. Geologists with the Ohio Division of Geological Survey have been conducting detailed bedrock geologic mapping, measuring and describing stratigraphic sections and cores, interpreting geophysical logs, and studying conodont biostratigraphy. Researchers at The Ohio State University have examined the graptolite and conodont biostratigraphy. Our efforts have resulted in: (1) the mapping of the Cincinnati Series in Ohio at a scale of 1:24,000, (2) the measurement and description of 100 stratigraphic sections, (3) the drilling, measurement, and description of 32 continuous cores, (4) the interpretation of 95 suites of geophysical logs, and (5) the development of biostratigraphic control on portions of nine cores. These data serve as the basis for our ongoing revision and refinement of the understanding of the surficial and subsurface lithostratigraphy, graptolite and conodont biostratigraphy, and depositional setting of the Cincinnati Series of southwestern Ohio.

LITHOSTRATIGRAPHY

HISTORICAL BACKGROUND

The lithostratigraphic nomenclature of the Cincinnati Series has been, and continues to be, a point of debate. This debate centers on whether the units originally described by Orton (1873), Nickles (1902, 1903), Foerste (1903, 1905, 1909), Bassler (1906), and Cumings (1908) are interpreted as biostratigraphic units or as having a major lithostrati-

graphic component as well as a biostratigraphic component. The original stratigraphic units are collectively termed traditional stratigraphic units. In figure 16-1, the stratigraphic column of Caster and others (1955) illustrates the traditional units.

The traditional stratigraphic units were accepted and used to describe the stratigraphy of the Cincinnati, Ohio, region until the 1960's. Beginning in the middle 1960's, some stratigraphers (for example, Brown and Lineback, 1966; Peck, 1966; Hatfield, 1968; Gray, 1972) abandoned the traditional stratigraphic units because they concluded that these units were defined as biostratigraphic units (fig. 16-1). They abandoned the traditional units in accordance with Article 6 of the Code of Stratigraphic Nomenclature (American Commission on Stratigraphic Nomenclature, 1961), which states that formal lithostratigraphic units should be based on lithology not on fossil content or fossil ranges.

On the other hand, a second group of stratigraphers believed the traditional units were defined in both a lithologic and biostratigraphic context and chose to retain these units redefined to the standards of the Code of Stratigraphic Nomenclature (for example, Ford, 1967; Tobin, 1986; see fig. 16-1). Some stratigraphers developed informal lithostratigraphic nomenclatures using a combination of traditional units and new lithostratigraphic units (for example, Lee, 1974; Hay, 1981; see fig. 16-1). Also, the reexamination of traditional stratigraphy led to the recognition of the Grand Avenue Member and Wesselman Tongue of the Kope Formation, the North Bend Tongue of the Fairview Formation, and the Miamitown Shale (Ford, 1967).

CURRENT LITHOSTRATIGRAPHY

The Ohio Division of Geological Survey decided to test the mappability of the updated traditional stratigraphic units of Ford (1967) and Tobin (1986) and to examine the merits of reinstating the name Cincinnati Group for the Late Ordovician rocks of Ohio. Many of these units, including the Miamitown Shale of Ford (1967), were found to be mappable over all or part of southwestern Ohio.

However, as mapping proceeded, it became apparent that certain units were not mappable at all or mappable only at scales of 1:24,000 or larger. These units will be discussed in ascending stratigraphic order beginning with the primarily sub-Cincinnati Point Pleasant Tongue of the Clays Ferry Formation (fig. 16-2).

Field mapping and subsurface correlation of the Point Pleasant Tongue of the Clays Ferry Formation of Swadley and others (1975) indicate that this unit in southwestern Ohio is not a single, limestone-dominant tongue as recognized in northern Kentucky but consists of a number of limestone- and shale-dominant tongues. These tongues grade northward in the subsurface into the brown shales of the "Utica shale" filling the Sebree Trough (Mitchell and Bergström, 1991).

I feel the use of the name Point Pleasant Tongue of the Clays Ferry Formation does not describe the multiple limestone- and shale-dominant tongues present in southwestern Ohio. No formal name has been proposed for this

		Madison to Brookville, Indiana, region				Maysville, Ky., region		Cincinnati, Ohio, region				
		Brown & Lineback (1966)	Hatfield (1968)	Gray (1972)	Hay (1981)	Peck (1966)	Lee (1974)	Caster, Dalvé, & Pope (1955)	Ford (1967)	Tobin (1986)		
CINCINNATIAN SERIES	RICHMONDIAN STAGE	Whitewater Fm.	Whitewater Fm.	Whitewater Fm.	Elkhorn Fm. Whitewater Fm.	Drakes Fm.	Drakes Fm.	Elkhorn Fm. Red Shale	beds	Saluda Fm. shale facies		
	MAYSVILLIAN STAGE	Saluda Fm.	Saluda Fm.	Saluda Mbr. Fm.	Saluda Fm.	Bull Fork Fm.	Bull Fork Fm.	Whitewater Fm.		Upper Mbr.	unnamed	Whitewater
		EDENIAN STAGE	Dillsboro Fm.	Tanner's Creek Fm.	Dillsboro Fm.			Brookville Fm.		Grant Lake Fm.		Grant Lake Fm.
	Liberty Mbr.					Liberty Fm.	Blanchester Mbr.					
	Randolph Co. Mbr.					Waynesville Fm.	Clarksville Mbr.					
	Waynesville Mbr.					Waynesville Fm.	Ft. Ancient Mbr.					
	Excello Mbr.					Waynesville Fm.	Oregonia Mbr.					
	Station Hollow Mbr.					Grant Lake Fm.	Sunset Mbr.					
	Bellevue Ls.					Grant Lake Fm.	Mt. Auburn Mbr.					
	Miamitown Sh.					Grant Lake Fm.	Corryville Mbr.					
EDENIAN STAGE	Kope Fm.	Kope Fm.	Kope Fm.	Kope Fm.	Kope Fm.	Kope Fm.	McMillan Fm.	Bellevue Ls. Mbr.	Fairview Fm.	Bellevue Ls.		
							Fairview Fm.	Fairview Fm.		Fairmount Mbr.	Fairview Fm.	
EDENIAN STAGE	Kope Fm.	Kope Fm.	Kope Fm.	Kope Fm.	Kope Fm.	Kope Fm.	Latonia Fm.	Southgate Mbr.	Kope Fm.	Kope Fm.		
											McMicken Mbr.	Grand Ave. Mbr.
											Economy Mbr.	

FIGURE 16-1.—Development of the lithostratigraphic nomenclature of the Cincinnati Series in its type area between 1960 and 1990 (modified from Schumacher, 1984, fig. 2).

stratigraphic interval, although the use of the name Point Pleasant Formation is under consideration by the Ohio Division of Geological Survey.

The Bellevue Limestone, the Miamitown Shale, and the Mt. Auburn Formation were mappable, valid units at map scales of 1:24,000 or larger. However, these units would be mapped as solid lines at map scales of 1:62,500 or smaller. Schumacher, Swinford, and Shrake (1991) stated that, at smaller map scales, in areas of high topographic relief, this line would represent the contacts between the Fairview Formation, the Miamitown Shale, the Bellevue Limestone, and the Corryville Formation.

Schumacher, Swinford, and Shrake (1991) also discussed the lithologic changes that take place between the Cincinnati, Ohio, region and the Maysville, Kentucky, region in the Bellevue Limestone, the Corryville Formation, and the Mt. Auburn Formation. The shale-dominant Mt. Auburn Formation grades laterally into a limestone-dominant unit. Also, the entire Bellevue, Corryville, and Mt. Auburn stratigraphic interval becomes increasing limestone dominant toward Maysville.

Schumacher, Swinford, and Shrake (1991) proposed some changes in the existing lithostratigraphic nomenclature to accommodate mapping at 1:62,500 and smaller map scales and to address the lateral lithologic changes present in this interval. They reduced the Bellevue Limestone, the Corryville Formation, and the Mt. Auburn Formation to members of the newly adopted Grant Lake Formation.

This reduction in rank: (1) retained the traditional stratigraphic units defined at the rank used by Bassler (1906), (2) eliminated the problem of multiple stratigraphic units being represented by solid lines at smaller map scales, and (3) provided convenient lithostratigraphic units to trace the lateral changes in lithologies present in these units. The introduction of the Grant Lake Formation provided a lithostratigraphic unit thick enough to map at smaller map scales and addressed the lateral change from a shale-dominant section in the Cincinnati, Ohio, region to a limestone-dominant section in the Maysville, Kentucky, region. The Grant Lake Limestone of Peck (1966) was adopted for the limestone-dominant portion of this section. The Bellevue and Corryville Members were recognized as members in the Grant Lake Limestone, and the new Straight Creek Member was introduced to address the lateral change from the shale-dominant Mt. Auburn Member to a limestone-dominant member in the Maysville, Kentucky, region.

The Sunset and Oregonia Formations of Tobin (1986) were not mappable over much of southwestern Ohio. The lithologies characterizing the Sunset and the Oregonia Formations were recognizable; however, the contact between these units was rarely exposed. The lack of a mappable contact results from a sizeable increase in the amount of shale in the upper part of the Sunset Formation. The large amount of shale weathers rapidly and commonly buries the contact between the Sunset and the Oregonia. The Ohio Division of Geological Survey has informally mapped the Arnheim




FIGURE 16-2.—Schematic diagram illustrating the Upper Ordovician lithostratigraphic nomenclature in southwestern Ohio as recognized by the Ohio Division of Geological Survey. Formal lithostratigraphic units are indicated by the unit name in upper-case type; informal lithostratigraphic units, by lower-case type. The area under the bold line represents the Upper Ordovician stratigraphic units exposed in the area. The “valley” depicts Late Ordovician or post-Ordovician erosion of the Drakes, Whitewater, and Liberty Formations in Clermont and Hamilton Counties, Ohio. The stippled area is the transition zone between the Grant Lake Limestone and the Grant Lake Formation (from Schumacher, Swinford, and Shrake, 1991, fig. 1).


Formation of Foerste (1912) in place of the Sunset and Oregonia Formations. We plan to formalize our use of the Arnheim Formation when field mapping of southwestern Ohio is complete.¹

The Waynesville, the Liberty, and the Whitewater Formations have been recognized and mapped throughout all or part of Butler, Clermont, Clinton, Greene, Hamilton, Montgomery, Preble, and Warren Counties of southwestern Ohio. The Liberty and Whitewater Formations are not present in Adams, Brown, or Highland Counties of south-central Ohio (fig. 16-2). In south-central Ohio, we have not decided whether to map the “Arnheim” and the Waynesville Formations or use a redefined Bull Fork Formation of Peck (1966) for this area.

The uppermost units of the Cincinnati Series consist of the Elkhorn Shale of western Ohio and the Preachersville

Member of the Drakes Formation in south-central Ohio. The stratigraphic relationships between these two units is poorly understood because of extensive burial by glacial drift along much of the outcrop belt. Recent core drilling conducted by the Ohio Division of Geological Survey in west-central Ohio has provided 10 new stratigraphic sections of this interval. Preliminary study of these cores indicates the presence of a number of repetitive stratigraphic sequences that are easily recognized on gamma ray geophysical logs (fig. 16-3). Each sequence consists of a basal unit (lithofacies 3) of laminated calcareous or dolomitic shale beds and minor interbeds of fossiliferous limestone and dolomite. This unit is overlain by an interval (lithofacies 2) of interlaminated or interbedded limestone and/or dolomite and shale grading into an argillaceous, bioturbated, fine-grained dolomite. These sequences range in thickness from 1 meter (3.3 ft) to over 6 meters (19.7 ft) and are interpreted as repetitive shoaling cycles, possibly as the result of Late Ordovician glaciation. Additional lithologies recognized in this interval include: intervals of massive-bedded, vuggy dolomite (lithofacies 1), intervals of micritic to coarse-grained limestone or

¹Editor’s note: Mapping by the Division of Geological Survey in southwestern Ohio has been completed since this paper was written in 1992. The unpublished 1:24,000-scale bedrock-geology, bedrock-topography, and structure-contour maps are on file at the Division.



LITHOFACIES: 1. Massive-bedded, vuggy dolomite.
 2. Interbedded or interlaminated dolomite and shale and dolomite; argillaceous with moderate to heavy bioturbation.
 3. Shale with rare interbeds of dolomite and limestone.
 4. Limestone and dolomite; limestone micritic to coarse grained; dolomite microcrystalline, heavy bioturbation.
 5. Interbedded, fossiliferous, wavy-bedded limestone and shale.

FIGURE 16-3.—Preliminary interpretation of the lithostratigraphy, generalized lithofacies, and shoaling cycles present in Ohio Division of Geological Survey core 3066 drilled in Shelby County, Ohio.

microcrystalline, bioturbated dolomite (lithofacies 4), and zones of wavy-bedded limestone and shale (lithofacies 5). Ongoing core drilling will continue to provide additional information to assist in the interpretation of the lithostratigraphy and the depositional environment of this interesting stratigraphic interval.

The Division of Geological Survey is considering reinstating the name Cincinnati Group introduced by Meek and Worthen (1865) for all the strata between the top of the sub-Cincinnatian Lexington Limestone and the Ordovician-Silurian systemic boundary in southwestern Ohio. I feel the name Cincinnati Group would be a useful lithostratigraphic term because: (1) Upper Ordovician rocks in the subsurface of southwestern and south-central Ohio are presently unnamed or misnamed, (2) Upper Ordovician surface and subsurface lithostratigraphic units are not included in one common unit, and (3) the introduction of a lithostratigraphic unit at the group level would avoid possible confusion with the chronostratigraphic unit represented by the Cincinnati Series. However, the use of the term Cincinnati Group is similar to the term Cincinnati Series, which may create, not avoid confusion in some geologists' minds. Presently, this issue is under consideration, and I would welcome your suggestions.

BIOSTRATIGRAPHY

HISTORICAL DEVELOPMENT

The fossil content of the interbedded shales and limestones of the Cincinnati Series is one of the characteristic features of these rocks. This fact was reflected in the original descriptions of the traditional stratigraphic units. For example, Nickles (1902) named the "Mt. Auburn or *Platystrophia lynx* Beds" for the predominantly blue shale with irregularly bedded limestone containing an abundance of the brachiopod *Platystrophia lynx* in the lower 5 to 12 ft (1.5 to 3.7 meters) of this unit. (What was then called *Platystrophia lynx* now is referred to *Platystrophia ponderosa*.) Nickles concluded his description with a faunal list describing the fossils present in the Mt. Auburn beds. This example illustrates the emphasis placed on recognition of fossil constituents and index fossils in the original descriptions of the traditional units. However, systematic collecting to determine the vertical and lateral ranges of the majority of the index fossils of the Cincinnati Series has yet to be undertaken.

The first truly extensive systematic studies of the vertical and lateral distribution of a group of Cincinnati index fossils were the conodont biostratigraphic investigations conducted by Ohio State University professor Walter C. Sweet and a host of students (see, for example, Sweet and others, 1959; Pulse and Sweet, 1960; Carpenter and Ory, 1961; Kohut and Sweet, 1968). These studies, along with conodont biostratigraphic studies of the underlying Lexington Limestone (Bergström and Sweet, 1966), resulted in (1) the determination of the stratigraphic ranges of 35 multi-element conodont species; (2) the recognition of representatives of conodont faunas inhabiting two distinct faunal provinces, a warm-temperate North American Midcontinent Province, and a cold-temperate Anglo-Scandinavian-Appalachian Province; (3) the development of relative-abundance logs for common Midcontinent species for purposes of regional correlation; and (4) the determination of the vertical ranges of diagnostic species of conodonts from the Upper Ordovician stratotype in order to develop the biostratigraphic framework for the Ordovician System (Sweet and others, 1971).

In the 1970's and 1980's, conodont biostratigraphic investigations focussed on the correlation of other Upper Ordovician stratigraphic sequences to the stratotype. Sweet (1984), using the graphic correlation methods of Shaw (1964), correlated the distribution of conodonts from 61 stratigraphic sections throughout much of the North American Midcontinent Province in order to develop a chronostratigraphic framework for the Mohawkian and Cincinnati Series. A Composite Standard Section was synthesized and subdivided into 80 vertically continuous, 6-meter (19.7-ft)-thick units representing approximately equal time intervals. These subdivisions, termed Standard Time Units, are believed to represent periods of time 462,500 years in length. When compared to existing chronostratigraphic schemes, these subdivisions increase the resolution by 10 to as much as 27 times.

CURRENT BIOSTRATIGRAPHY

Recent biostratigraphic studies of the Cincinnati Series in its type area have focussed on graptolite biostratig-

raphy (Bergström and Mitchell, 1986, 1990a, 1990b; Mitchell and Bergström, 1991). These studies have described the diverse fauna of graptolites from the dark shales present in the subsurface Sebree Trough. This new information has resulted in better definition of the base of the Cincinnati Series in its type area and allows correlation of these sediments with other graptolite-bearing sequences throughout the world. Stig Bergström (The Ohio State University), Charles Mitchell (State University of New York at Buffalo), and I hope to publish a paper discussing the origin and evolution of the Sebree Trough through time and the tectonic relationships between the Sebree Trough and the Taconic Orogeny.

DEPOSITIONAL ENVIRONMENT

Regional studies of Anstey and Fowler (1969), Hay (1981), and Tobin (1982) have developed the framework necessary to understand the depositional environment of the Cincinnati Series. These authors have recognized the recurrence of distinctive lithofacies throughout the Cincinnati Series. These lithofacies were developed as the result of interaction between eustasy, basinal tectonics, and biological productivity. A typical succession ranged from an offshore, "deeper water," shale-dominant lithofacies through a lower shoreface, transitional, mixed limestone and shale lithofacies to an upper shoreface, "shallow-water," limestone- or dolomite-dominant lithofacies. Additional characteristics of each lithofacies are summarized in table 16-1.

The "deeper water" lithofacies was deposited in water depths estimated to have been between 20 and possibly 50 meters (66-164 ft) (Bucher, 1917; Anstey and Fowler, 1969; Anstey and others, 1987). These authors compared Cincinnati megaripples and bryozoan colony heights to modern analogs to make their paleobathymetric estimates.

Three shoaling cycles (shoaling-upward cycles of some authors) have been recognized in the Cincinnati Series in the vicinity of Cincinnati, Ohio (Anstey and Fowler, 1969; Hay, 1981; and Tobin, 1982). These authors generally agree in their placement of the beginning and end points of these cycles (fig. 16-4). The first shoaling cycle ranges from the Kope Formation to the top of the Bellevue Member of the Grant Lake Formation. The second, from the base of the Corryville Member of the Grant Lake Formation to the top of the "Arnheim formation," and the third, from the base of the Waynesville Formation to the Ordovician-Silurian systemic boundary.

The lithofacies succession of a given shoaling cycle is readily observed in the field and has been recognized in cores and in shale-percentage and geophysical logs. In the

Maysville, Kentucky, region, I have identified five shoaling cycles in the Cincinnati Series (Schumacher, 1992) (figs. 16-4 and 16-5). I have subdivided the Kope Formation-to-Bellevue Limestone shoaling cycle of previous workers into two cycles on the basis of the presence of a "shallow-water" lithofacies in the base of the Fairview Formation.

The third cycle ranges from the transitional lithofacies in the basal portion of the Corryville Member of the Grant Lake Limestone through the top of the "shallow-water" lithofacies of the Straight Creek Member of the Grant Lake Limestone. The fourth shoaling cycle is from the base to the top of the "Arnheim formation," and the fifth cycle ranges from the basal Waynesville Formation to the top of the Drakes Formation.


I contend that the five shoaling cycles recognized in the Maysville, Kentucky, region also are present in the Cincinnati, Ohio, region. Field mapping supplemented with shale-percentage logs for continuous cores indicates that the "shallow-water" lithofacies of the basal portion of the Fairview Formation in the Maysville, Kentucky, region can be correlated to the North Bend Tongue of the Fairview Formation of the Cincinnati, Ohio, region (fig. 16-5). The "shallow-water" lithofacies present in the Maysville, Kentucky, region grades laterally into a limestone-dominant transitional lithofacies, then to the shale-dominant transitional lithofacies of the North Bend Tongue. This lateral progression of lithofacies illustrates the progressive increase in water depth from Maysville to Cincinnati.

Ford (1967) documented the intertonguing relationship between the shale-dominant transitional lithofacies of the North Bend Tongue of the Fairview Formation and the shale-dominant, "deeper water" lithofacies of the Wesselman Tongue of the Kope Formation. The Wesselman Tongue of the Kope Formation overlying the North Bend Tongue of the Fairview Formation reflects a return to "deeper water" conditions. Thus, the top of the North Bend Tongue of the Fairview Formation is the top of the first shoaling cycle in the Cincinnati region. The lower part of this cycle originates in the middle of the Kope Formation. Laterally, the Wesselman Tongue is overlain by and grades into the shale-dominant transitional lithofacies characterizing the middle part of the Fairview Formation. This lithofacies records the beginning of the second shoaling cycle which ends at the top of the "shallow-water" lithofacies represented by the Bellevue Member of the Grant Lake Formation (fig. 16-5).

Within the second shoaling cycle in the Cincinnati, Ohio, region, the normal progression from the basal, shale-dominant, transitional lithofacies through shallower, limestone-dominant, transitional lithofacies to limestone-dominant "shallow-water" lithofacies is interrupted by the presence

TABLE 16-1.—Diagnostic features of the lithofacies observed in the Cincinnati Series of southwestern Ohio

Lithofacies	Dominant lithology	Subordinate lithology	Dominant lithology bedding		Average range of shale percentage	Fossil content of shales
			Thickness	Style		
"deeper water"	shale	limestone	thick (<.3048 m)	planar	60-100%	sparsely fossiliferous
transitional	interbedded limestone and shale or shale and limestone		medium (between .3048 m and .1016 m)	planar to irregular	40-60%	sparsely fossiliferous
"shallow water"	limestone	shale	thin (>.1016 m)	wavy, nodular, irregular	0-40%	fossiliferous



¹After Schumacher, Swinford, and Shrake, 1991

FIGURE 16-4.—Schematic diagram comparing the lithostratigraphy, the lithofacies, and the interpretation of shoaling cycles of the Cincinnati Series exposed in the Cincinnati, Ohio, and Maysville, Kentucky, regions.

of the Miamitown Shale.

The shale-dominant lithofacies of the Miamitown Shale intertongues with the upper part of the limestone-dominant transitional lithofacies of the Fairview Formation and the lower portion of the limestone-dominant “shallow-water” lithofacies of the Bellevue Member of the Grant Lake Formation in Butler, Hamilton, and Warren Counties and the western portion of Clermont County, Ohio. Ford (1967) and Hay (1981) interpreted the Miamitown Shale as representing a shale-dominant “deeper water” lithofacies. Tobin (1982) believed this unit represents a shale-dominant, “shallow-water” lithofacies. Dattilo (paper 7 in this volume) provides additional information on the interpretation of the

lithofacies of the Miamitown Shale.

My experience with the Miamitown Shale is limited to a few exposures mapped in Clermont and Warren Counties. I recognized two discrete tongues of Miamitown Shale and the meter-scale shoaling-upward cycles described by Dattilo (1991). The tongues of Miamitown rapidly pinch out or grade into the Fairview Formation or the Bellevue Member of the Grant Lake Formation. The limits of the Miamitown as a mappable unit have been delineated in parts of Clermont and Warren Counties by Schumacher (1990) and Shrake and Schumacher (1992). I do not know what the depositional setting of the Miamitown Shale is because of my lack of experience with this unit.




FIGURE 16-5.—Correlation of the upper part of the Kope Formation, the Fairview Formation, and the lower part of the Grant Lake Limestone/Formation between the Cincinnati, Ohio, and the Maysville, Kentucky, regions, illustrating the vertical and lateral changes in lithofacies. The basal two shoaling cycles of the Cincinnati Series are illustrated by the numbered arrows at the margins of the cross section. Note the lateral gradation of lithofacies in the lower and upper portions of the Fairview Formation.

The third shoaling cycle is from the base of the transitional lithofacies represented by the Corryville Member of the Grant Lake Formation to the top of the Mt. Auburn Member of the Grant Lake Formation. The transitional to “shallow-water” lithofacies of the Mt. Auburn Member is the lateral equivalent of the “shallow-water” lithofacies of the Straight Creek Member of the Grant Lake Limestone.

The fourth cycle ranges from the transitional lithofacies of the lower part of the “Arnheim formation” to the nodular top of the “Arnheim” throughout southwestern Ohio. The fifth shoaling cycle ranges from the base of the “deeper water” lithofacies of the Waynesville Formation to the top of the Cincinnati Series in southwestern Ohio.

The fifth shoaling cycle is likely an oversimplification of the cyclicity present in the uppermost part of the Cincinnati Series. The ongoing investigation of the repetitive stratigraphic sequences discovered in the subsurface of west-central Ohio will lead to the recognition of a number of additional shoaling cycles. Presently, I am unsure whether these cycles will be recognized as independent shoaling cycles or as subcycles within the fifth shoaling cycle.

The recognition of intervals characterized by shoaling is beneficial in reconstructing the local geologic setting of the Cincinnati Series throughout its deposition. Weir and

others (1984) and Tobin (1986) suggested the Cincinnati Series was deposited on a gently northward sloping, mixed siliciclastic-carbonate ramp throughout Cincinnati time. This depositional setting affected sedimentation throughout much of the sub-Cincinnati and through the first shoaling cycle in the Cincinnati Series. In the second and third shoaling cycles, the area of shoaling environments shifted from the Lexington, Kentucky, region to the Maysville, Kentucky, region.

Apparently, as the area of shoaling shifted, so did the direction of the slope of the associated siliciclastic-carbonate ramp. The “deeper water” portion of this ramp was oriented to the northwest, explaining the progressive change from predominantly “shallow-water” lithofacies present in the Maysville, Kentucky, region to the predominantly transitional lithofacies of the Cincinnati, Ohio, region.

The area of shoaling environments shifted from the Maysville, Kentucky, region to the Richmond, Indiana, region in the fifth shoaling cycle. The “shallow-water” lithofacies associated with the Whitewater Formation can be correlated along the outcrop belt of southeastern Indiana into the subsurface of the eastern third of Indiana and portions of west-central Ohio (Gray, 1972; Schumacher, Bergström, and Mitchell, 1991). A mixed siliciclastic-carbonate ramp

flanked either side of this north-south-trending shoal, and "deeper water" lithofacies occurred in central Ohio and central Indiana.

Anstey and Fowler (1969), Hay (1981), and Tobin (1982) suggested that shoaling cycles occurred as the result of changes in sea level caused by local or regional tectonics, basin sedimentation patterns, Late Ordovician glaciation, or a combination of these. The exact cause or causes of the shoaling cycles cannot be determined at this time. This uncertainty is the result of the poor understanding of the stratigraphic relationships between the Cincinnati, Ohio, stratigraphic section and coeval sections in the Appalachian Mountains and Midcontinent region. This poor understanding is unfortunate, because recent investigations have refined our understanding of the Taconic Orogeny and its effect on Midcontinent sedimentation (for example, Ettensohn, 1991). The Ohio Division of Geological Survey has correlated surficial Cincinnati lithostratigraphic units into the subsurface adjacent to the outcrop belt, but more study is needed to bridge the gap between Upper Ordovician outcrops in the Appalachian Mountains and the Midcontinent region.

Tobin (1982) and Jennette and Pryor (1993) have recognized additional sedimentary cycles from the Cincinnati Series of southwestern Ohio. These cycles, termed megacycles and storm cycles by Tobin (1982) and shallowing-upward bedding cycles and storm beds by Jennette and Pryor (1993), range in thickness from 1 to 3 meters (3.3-9.8 ft) and less than 1 meter (3.3 ft), respectively. Tobin suggested that megacycles were the result of fluctuating sediment supply because of deltaic shifting, changing tectonics, or climatic changes. Jennette and Pryor (1993) believed that shallowing-upward bedding cycles are the result of cyclic fluctuation of sea level caused by glacio-eustasy.

Megacycles, shallowing-upward cycles, and storm cycles have been recognized in the course of our field mapping. The Ohio Division of Geological Survey has not attempted detailed correlation of these cycles. However, a few observations can be stated concerning them. Megacycles and shallowing-upward cycles occur in many of the formations of the Cincinnati Series, but not everywhere in the abundance in which they occur in the upper portion of the Kope Formation and the lower part of the Fairview Formation. Storm cycles or beds occur in all the lithostratigraphic units of the Cincinnati Series, and many limestone and shale beds have been produced by or influenced by storm-generated sedimentary processes.

What do the various scales of sedimentary cycles tell us about the formation of the Cincinnati Series? For over a decade, I have been working to resolve this question. I have not finalized my answer, because detailed subsurface correlation to coeval rocks exposed in the Appalachian Mountains and the Midcontinent region has not been completed. My feeling is that, when these correlations are completed, some shoaling cycles will be traceable between the Cincinnati, Ohio, region and the Appalachian exposures. The recognition of these cycles will allow the interpretation of how the tectonic events associated with the Taconic Orogeny affected sedimentation in the Cincinnati Series.

With updated regional correlations, my guess is that the Cincinnati Series will be interpreted as mixed siliciclastic-carbonate sedimentation on at least three gently sloping, storm-dominated ramps. The orientation of these ramps varied through time because of the influence of tectonic pulses associated with the Taconic Orogeny.

REFERENCES CITED

- American Commission on Stratigraphic Nomenclature, 1961, Code of Stratigraphic Nomenclature: American Association of Petroleum Geologists Bulletin, v. 45, p. 645-665.
- Anstey, R. L., and Fowler, M. L., 1969, Lithostratigraphy and depositional environment of the Eden Shale (Ordovician) in the tri-state area of Indiana, Kentucky, and Ohio: *Journal of Geology*, v. 77, p. 668-682.
- Anstey, R. L., Rabbio, S. F., and Tuckey, M. E., 1987, Bryozoan bathymetric gradients within a Late Ordovician epeiric sea: *Paleoceanography*, v. 2, p. 165-176.
- Bassler, R. S., 1906, A study of the James types of Ordovician and Silurian Bryozoa: *Proceedings of the U.S. National Museum*, v. 30, no. 1442, p. 1-66.
- Bergström, S. M., and Mitchell, C. E., 1986, The graptolite correlation of the North American Upper Ordovician Standard: *Lethaia*, v. 19, p. 247-266.
- _____, 1990a, Trans-Pacific graptolite faunal relations: the biostratigraphic position of the base of the Cincinnati Series (Upper Ordovician) in the standard Australian graptolite zone succession: *Journal of Paleontology*, v. 64, p. 992-997.
- _____, 1990b, The Utica Shale of northern Ohio and its relationship to the Utica Shale of the northern Appalachian Basin and lithologically similar rocks in the central Great Lakes region: *Appalachian Basin Industrial Associates Program for Meeting*, v. 17, p. 2-39.
- Bergström, S. M., and Sweet, W. C., 1966, Conodonts from the Lexington Limestone (Middle Ordovician) of Kentucky and its lateral equivalents in Ohio and Indiana: *Bulletins of American Paleontology*, v. 50, no. 229, p. 271-441.
- Brown, G. D., Jr., and Lineback, J. A., 1966, Lithostratigraphy of the Cincinnati Series (Upper Ordovician) in southeastern Indiana: *American Association of Petroleum Geologists Bulletin*, v. 50, p. 1018-1023.
- Bucher, W. H., 1917, Large current-ripples as indicators of paleogeography: *Proceedings of the U.S. National Academy of Sciences*, v. 3, p. 285-291.
- Carpenter, J. W., and Ory, T. R., 1961, The American Upper Ordovician Standard. VI. The Covington sequence at Maysville, Kentucky: *Ohio Journal of Science*, v. 61, p. 372-378.
- Caster, K. E., Dalvé, E. A., and Pope, J. K., 1955, Elementary guide to the fossils and strata of the Ordovician in the vicinity of Cincinnati: *Cincinnati Museum of Natural History*, 47 p.
- Caster, K. E., and Kjellesvig-Waering, E. N., 1964, Upper Ordovician eurypterids of Ohio: *Palaeontographica Americana*, v. 4, no. 32, p. 301-358.
- Cummings, E. R., 1908, The stratigraphy and paleontology of the Cincinnati Series of Indiana: *Indiana Department of Geology and Natural Resources, Thirty-second Annual Report for 1907*, p. 605-1189.
- Dattilo, B. F., 1991, The Miami town shale (Upper Ordovician): a facies after all (abs.): *Geological Society of America Abstracts with Programs*, v. 23, no. 3, p. 9.
- Drake, Daniel, 1825, Geological account of the Valley of the Ohio: *Transactions of the American Philosophical Society*, new ser., v. 2, p. 124-139.
- Ettensohn, F. R., 1991, Flexure interpretation of relationships between Ordovician tectonism and stratigraphic sequences, central and southern Appalachians, U.S.A., in Barnes, C. R., and Williams, S. H., eds., *Advances in Ordovician geology: Geological Survey of Canada Paper 90-9*, p. 213-224.
- Foerste, A. F., 1903, The Richmond Group along the western side of the Cincinnati Anticline in Indiana and Kentucky: *American Geologist*, v. 31, p. 333-361.
- _____, 1905, The classification of the Ordovician rocks of Ohio and Indiana: *Science*, new ser., v. 22, no. 553, p. 149-152.
- _____, 1909, Preliminary notes on Cincinnati and Lexington fossils: *Bulletin of the Scientific Laboratories of Denison University*, v. 14, art. 17, p. 289-333.
- _____, 1912, The Arnheim Formation within the areas traversed by the Cincinnati Geanticline: *Ohio Naturalist*, v. 12,

- p. 429-456.
- Ford, J. P., 1967, Cincinnatian geology in southwest Hamilton County, Ohio: American Association of Petroleum Geologists Bulletin, v. 51, p. 918-936.
- Gray, H. H., 1972, Lithostratigraphy of the Maquoketa Group (Ordovician) in Indiana: Indiana Geological Survey Special Report 7, 31 p.
- Hatfield, C. B., 1968, Stratigraphy and paleoecology of the Saluda Formation (Cincinnatian) in Indiana, Ohio, and Kentucky: Geological Society of America Special Paper 95, 34 p.
- Hay, H. B., 1981, Lithofacies and formations of the Cincinnatian Series (Upper Ordovician), southeastern Indiana and southwestern Ohio: Ph.D. dissertation (unpub.), Miami University, 236 p.
- James, J. F., 1886, Cephalopoda of the Cincinnati Group: Journal of the Cincinnati Society of Natural History, v. 8, p. 235-253.
- Jennette, D. C., and Pryor, W. A., 1993, Cyclic alternation of proximal and distal storm facies: Kope and Fairview Formations (Upper Ordovician), Ohio and Kentucky: Journal of Sedimentary Petrology, v. 63, p. 183-203.
- Kohut, J. J., and Sweet, W. C., 1968, The American Upper Ordovician Standard. X. Upper Maysville and Richmond conodonts from the Cincinnati region of Ohio, Indiana, and Kentucky: Journal of Paleontology, v. 42, p. 1456-1477.
- Lee, G. B., 1974, Lithostratigraphy of the Cincinnatian Series (Upper Ordovician) from Maysville, Kentucky to Dayton, Ohio: M.S. thesis (unpub.), Miami University, 127 p.
- Meek, F. B., and Worthen, A. H., 1865, Descriptions of new species of Crinoidea, etc., from the Paleozoic rocks of Illinois and some of the adjoining states: Academy of Natural Sciences of Philadelphia Proceedings, v. 17, p. 143-155.
- Mitchell, C. E., and Bergström, S. M., 1991, New graptolite and lithostratigraphic evidence from the Cincinnati region, U.S.A., for the definition and correlation of the base of the Cincinnatian Series (Upper Ordovician), in Barnes, C. R., and Williams, S. H., eds., Advances in Ordovician geology: Geological Survey of Canada Paper 90-9, p. 59-77.
- Nickles, J. M., 1902, The geology of Cincinnati: Cincinnati Society of Natural History Journal, v. 20, p. 49-100; reprint edition, p. 1-53.
- Nickles, J. M., 1903, The Richmond Group in Ohio and Indiana and its subdivisions, with a note on the genus *Strophomena* and its type: American Geologist, v. 32, p. 202-218.
- Orton, Edward, 1873, Report on the Third Geological District; geology of the Cincinnati Group; Hamilton, Clermont, Warren and Butler Counties: Ohio Division of Geological Survey, v. 1, pt. 1, Geology, p. 365-480.
- Peck, J. H., 1966, Upper Ordovician formations in the Maysville area, Kentucky: U.S. Geological Survey Bulletin 1244-B, 30 p.
- Pulse, R. R., and Sweet, W. C., 1960, The American Upper Ordovician Standard. III. Conodonts from the Fairview and McMullan Formations of Ohio, Kentucky, and Indiana: Journal of Paleontology, v. 34, p. 237-264.
- Schumacher, G. A., 1984, Sedimentology, biostratigraphy, and paleoecology of Cincinnatian (Upper Ordovician) *Iocrinus subcrassus* crinoid (Echinodermata) assemblages: M.S. thesis (unpub.), University of Cincinnati, 210 p.
- _____, 1990, Bedrock geology of the Goshen, Ohio, quadrangle: Ohio Division of Geological Survey Open-File Map BG-B5B2.
- _____, 1992, Lithostratigraphy, cyclic sedimentation, and event stratigraphy of the Maysville, Kentucky, area, Stop 8 in Ettensohn, F. R., ed., Changing interpretations of Kentucky geology: layer cake, facies, flexure, and eustasy (field trip for 1992 Annual Meeting of Geological Society of America): Ohio Division of Geological Survey Miscellaneous Report 5, p. 165-172.
- Schumacher, G. A., Bergström, S. M., and Mitchell, C. E., 1991, The Sebree Trough project. 4: preliminary tracing of surficial Upper Ordovician lithostratigraphic units into the subsurface of western and northern Ohio (abs.): Geological Society of America Abstracts with Programs, v. 23, no. 3, p. 59.
- Schumacher, G. A., Swinford, E. M., and Shrake, D. L., 1991, Lithostratigraphy of the Grant Lake Limestone and Grant Lake Formation (Upper Ordovician) in southwestern Ohio: Ohio Journal of Science, v. 91, p. 56-68.
- Shaw, A. B., 1964, Time in stratigraphy: New York, McGraw-Hill, 365 p.
- Shrake, D. L., and Schumacher, G. A., 1992, Bedrock geologic map of the South Lebanon, Ohio, quadrangle: Ohio Division of Geological Survey Open-File Map BG-B5C2.
- Swadley, W. C., Luft, S. J., and Gibbons, A. B., 1975, The Point Pleasant Tongue of the Clays Ferry Formation, northern Kentucky: U.S. Geological Survey Bulletin 1405-A, p. 30-31.
- Sweet, W. C., 1984, Graphic correlation of upper Middle and Upper Ordovician rocks, North American Midcontinent Province, U.S.A., in Bruton, D. L., ed., Aspects of the Ordovician System: Palaeontological Contributions from the University of Oslo 295, p. 23-25.
- Sweet, W. C., Ethington, R. L., and Barnes, C. R., 1971, North American Middle and Upper Ordovician conodont faunas, in Sweet, W. C., and Bergström, S. M., eds., Symposium on Conodont Biostratigraphy: Geological Society of America Memoir 127, p. 163-193.
- Sweet, W. C., Turco, C. A., Warner, Earl, Jr., and Wilkie, L. C., 1959, The American Upper Ordovician Standard. I. Eden conodonts from the Cincinnati region, Ohio and Kentucky: Journal of Paleontology, v. 33, p. 1029-1068.
- Tobin, R. C., 1982, A model for cyclic deposition in the Cincinnatian Series of southwestern Ohio, northern Kentucky, and southeastern Indiana: Ph.D. dissertation (unpub.), University of Cincinnati, 483 p.
- _____, 1986, An assessment of the lithostratigraphic and interpretive value of the traditional "biostratigraphy" of the type Upper Ordovician of North America: American Journal of Science, v. 286, p. 673-701.
- Warn, J. M., and Strimple, H. L., 1977, The disparid inadunate superfamilies Homocrinacea and Cincinnaticrinacea (Echinodermata, Crinoidea), Ordovician-Silurian, North America: Bulletins of American Paleontology, v. 72, no. 296, p. 1-138.
- Weir, G. W., Peterson, W. L., Swadley, W. C., and Pojeta, John, Jr., 1984, Lithostratigraphy of Upper Ordovician strata exposed in Kentucky; with a section on biostratigraphy: U.S. Geological Survey Professional Paper 1151-E, 121 p.

Address of author: Gregory A. Schumacher, Ohio Department of Natural Resources, Division of Geological Survey, 4383 Fountain Square Drive, Columbus, OH 43224-1362.

17. PALEOGEOGRAPHY AND PALEOENVIRONMENTS, FAIRVIEW THROUGH WHITEWATER FORMATIONS (UPPER ORDOVICIAN, SOUTHEASTERN INDIANA AND SOUTHWESTERN OHIO)

by
Helen B. Hay

INTRODUCTION

The focus of this paper is the lithofacies of about 122-168 meters (400-550 ft) of section that includes the upper part of the Kope Formation through the Whitewater Formation to the Ordovician/Silurian contact in southeastern Indiana and southwestern Ohio. The goal is to determine what the stratigraphic and geographic distribution of the facies suggests about the paleogeography, paleoenvironments, and geologic history of the area. The rocks are thin gray limestones composed of whole and broken marine fossils interbedded with compacted, but not cemented, terrigenous clay and silt that are traditionally called shales, although they are more accurately classified as mudstones. Facies intervals differ from each other in the proportion of shale to limestone and in internal and bedding characteristics of both rock types.

Gray (1972) assigned the Upper Ordovician of Indiana to the Maquoketa Group and identified four provinces: a deep basin in the southwest, a western shelf in the northwest, an eastern shelf, and a southeastern shelf. Most of the limestones are in the eastern part of the eastern shelf and in the southeastern shelf, the eastern parts of which include the area of this study. Most of the Cincinnati of the central and western parts of the state is shale. Weir and Peck (1968) and Cressman (1973) studied the Middle and Upper Ordovician rocks of Kentucky, most of which are shallow-water limestones. So this paper deals with details of an area viewed as shelf from central Indiana and basin from central Kentucky.

In general, these sediments accumulated on a shallow marine shelf at depths that ranged from above normal wave base to, more commonly, near or below storm wave base. The large number of thin facies intervals suggests that the depositional environment was subject to frequent minor changes that would be expected in very shallow seas rather than in the more stable environments of deeper shelves. The regional facies patterns show wedges of >70 percent shale that thicken toward the northwest or northeast. These shale wedges intertongue with limestone-rich zones of <70 percent shale that thicken first toward the south then, later in the Cincinnati, along a more nearly north-south axis. Most of the contacts between units are gradational and, therefore, are believed to be time transgressive. The characteristics of the lithofacies, as well as their stratigraphic relationships based on isopach maps, leads to a reasonable interpretation of depositional environments, paleogeography, and transgressive/regressive cycles.

Localities included in the study are shown in figure 17-1 and are listed at the end of this paper. Hay (1981) described the geographically named localities, five of which are cores: New Point, Wayne County, Randolph County, Miamisburg, and Middletown. Descriptions of part of the Brookville composite section and of some other localities were published by Hay and others (1981), and a portion of the Madison composite section by Totten and Hay (1987). The numbered localities (1-22) are mostly from Kentucky GQ maps. Information from open-file core descriptions provided by the Ohio Division of Geological Survey (2627 in Warren County; 2537,




FIGURE 17-1.—Index map (modified from Hay, 1981). Localities marked "core" are cores of the Ohio Division of Geological Survey. For references for numbered localities (1-22) see Localities on p. 132-133.

2981, 2982, and 2984 in Butler County; and 3023 in Preble County) are incorporated, to the extent possible, in this discussion. Several localities are described in other papers in this volume: the Indiana Route 1 road cut at South Gate Hill (IN-FR-0005; paper 12 in this volume); the Madison, Indiana, road cuts (IN-JE-0001, IN-JE-0002, and IN-JE-0003; paper 6); the Brookville Dam spillway (IN-FR-0002; paper 8); Bon Well Hill (IN-FR-0001; paper 10); Garr Hill/Brookville North (IN-FR-0003; paper 11); and the U.S. Route 27 road cut near Richmond, Indiana (IN-WY-0001; paper 15).

LITHOFACIES AND STRATIGRAPHY

LITHOFACIES CLASSIFICATION

The lithofacies classification that underpins this discussion was developed through detailed study of the Brookville and Richmond composite sections (Hay, 1975), then modified and applied regionally (Hay, 1981). It is a field classification that has proven effective in describing both outcrops and cores. The key characteristics of the lithofacies are summarized in figure 17-2. There are four facies groups. Groups 1, 2, and 3 are distinguished on the basis of the shale percentage in an interval: >70 percent, 55-70 percent, and




FIGURE 17-2.—Principal diagnostic characteristics of lithofacies and illustration of lithofacies types. Modified from Hay (1981) and Hay and others (1981).

<55 percent, respectively. Within these three groups the letter designations are based on the nature of the shales and limestones. The letter “a” indicates a predominance of planar rather than nodular or rubbly limestones and shales that are predominantly free of calcareous stringers, lenses, or nodules. The “b” facies have calcareous stringers, lenses, or nodules in the majority of the shales. Group 3 (<55 percent shale) has two additional facies: 3c, containing nodular, rubbly limestones and limy shales, and 3d, containing planar limestones, >20 percent of which are laminated, burrowed, and composed of silt- and fine-sand-sized allochems. The latter facies appears to be unfossiliferous in hand specimen or outcrop. The facies of group 4 are much less common but are useful environmental indicators where they occur. The classification is open ended in that all four groups can be expanded, if the need arises.

FORMATIONS AND MEMBERS

Regional correlation of lithofacies (Hay, 1981, pls. 1 and 2) suggests a reasonable division of the Cincinnati Series

into formations and members (Hay 1981, pl. 3) that is somewhat different from that currently in use in Indiana and Ohio (fig. 17-3). The Indiana Geological Survey recognizes four formations: the Kope, Dillsboro, Saluda, and White-water Formations. The Ohio Division of Geological Survey recognizes the Kope and Fairview Formations, Miami-town Shale, Grant Lake Formation (divided into the Bellevue, Corryville, and Mt. Auburn Members by Schumacher, Swinford, and Shrake, 1991), the Arnheim formation (informal), the Waynesville, Liberty, and Whitewater Formations, and the Elkhorn shale (informal). In south-central Ohio, the Ohio Survey recognizes the Drakes Formation at the top of the Ordovician.

My suggestion, presented here informally, incorporates some of the units of Ohio and Indiana, but, in some respects, is different from either. It is illustrated schematically along a south-north cross section in figure 17-4. The Kope (facies 1a), Fairview (facies 2a and 3a), Miami-town (facies 1a), and Bellevue (facies 3b and 3c) can be traced throughout the area and should be retained in Ohio and recognized in Indiana.

The Bellevue lithology is similar to that of the Whitewa-

Series	Stage	Current Classification of Others				(not to scale)	Hay (1981) (informal)		
		(1)	Ohio (2)	Kentucky (3)	Indiana (4)		Formations	Members	
Cincinnati (Upper Ordovician)	Richmondian	Elkhorn	Elkhorn / Drakes	Drakes	Whitewater		Whitewater	Elkhorn sh.	
		Whitewater	(as in column to left)	Bull Fork	Dillsboro		Saluda		Saluda
		Liberty					Brookville	Liberty sh.	
		Waynesville						Waynesville sh.	
		Arnheim						undifferentiated	
	Maysvillian	McMillan				Mount Auburn	Mount Auburn		Grant Lake Limestone
			Corryville	Corryville	Station Hollow sh.				
		Fairview	Bellevue	Bellevue	Bellevue				
			Fairmount	Fairview	Fairview				
	Edenian	Latonia	Kope	Kope	Kope	Kope	Kope		

FIGURE 17-3.—Rock-stratigraphic classification (modified from Hay, 1981, and Hay and others, 1981). In the “current classification of others,” (1) = Caster and others (1955) and Davis (1992). (2) = Kope from Weiss and Sweet (1964); Fairview and Miami town from Ford (1967); Grant Lake Formation and members upward from Schumacher and others (1991). (3) = Peck (1966). (4) = Brown and Lineback (1966) and Gray (1972).




FIGURE 17-4.—Generalized south-north distribution of formations and members as identified in this paper. The Bellevue Formation of the diagram is the Bellevue Member of the Grant Lake Formation as used by Schumacher and others (1991). Modified from Hay (1981) and Hay and others (1981).

ter (facies 3b and/or 3c), and I suggest recognition of a new formation, the “Brookville formation,” for the interval of variable lithology between the Bellevue and the Whitewater or the Saluda, whichever is lower in a given section. The “Brookville formation” can be divided into facies intervals that could be considered members or tongues: the “Excello member,” the Waynesville Shale Member, the Liberty Member and the “Randolph County Shale member.” Within the

“Excello” there are two shale tongues (>70 percent shale) called the “Station Hollow shales.” In 1981 I believed these shale tongues merged in the vicinity of the Miamisburg core and therefore called them tongues of the same unit. In view of new information in nearby cores, I now think these are two separate shale wedges and should not be made a single member of the “Brookville formation,” although they should be identified in the description of cores and outcrops because of what they can show about sedimentation patterns in the area. The “Station Hollow shales” are at least in part equivalent to the Corryville Member of the Grant Lake Formation in Ohio. The “Excello” includes the Corryville and Mt. Auburn Members of the Grant Lake Formation and the Arnheim formation of Ohio nomenclature. The “Excello” consists of a variety of facies, mostly of groups 2 (55-70 percent shale) but including some group 3 (<55 percent shale), and mostly in “b” or “c” divisions (limy shales and/or rubbly limestones). Overall, the greatest volume of the “Excello” is facies 2b.

The Waynesville-Liberty interval is characterized by decreasing shale percentage upward. On the basis of the descriptions of their cores, the Ohio Division of Geological Survey seems to put the contact approximately where the shale percentage drops below 55 percent, which correlates with my boundary between facies 2a and 3a. I prefer to

place the Waynesville-Liberty contact lower in the section so that the Liberty includes facies of Group 2, mainly 2a, as well as 3a, and limits the Waynesville to the predominantly 1a facies. This placement seems reasonable in view of the fact that the Liberty, as here defined, includes the same facies sequence as the lithologically similar Fairview Formation, facies 2a overlain by 3a. This placement makes the Waynesville Shale Member comparable in facies to the Miamitown and "Station Hollow shales" wedges that underlie and overlie the Bellevue, respectively. Unfortunately, at the present time, I do not know where this 70 percent-shale boundary occurs in the Ohio cores, so I cannot relate the Waynesville-Liberty boundary in those cores to its placement based on the 70 percent-shale criterion. Within the Waynesville Shale Member, as defined here, there is a thin but widespread interval of facies 2a or 3a. It would also be useful to know where this interval falls within the Ohio cores, because it constitutes a useful datum for regional correlation.

In the Randolph County and Wayne County cores, and possibly in the Miamisburg core, a tongue of facies 1a (>70 percent shale) within the Liberty has been called the "Randolph County Shale member." Inspection of the Ohio Survey cores, particularly those in the north, should make it possible to identify this shale tongue. It probably is within the upper part of their Waynesville Formation.

In summary, the "Brookville formation" consists of shale wedges—the "Station Hollow shales" in the "Excello member," the Waynesville Shale Member, and the "Randolph County shale member"—that lie between the intervals of lower clastic ratio, the "Excello" and Liberty Members. Most of the "Excello" and the Liberty consist of facies in groups 2 and 3. The "Excello member" generally has limy shales and some rubbly limestones of facies 2b and, less commonly, 3b or 3c. The Liberty Member is predominantly facies 2a in the lower part and 3a in the upper part, a decrease in shale upward. The "Excello member" differs from the Bellevue and Whitewater Formations in that most of it is within the 55-70 percent shale range, whereas the Bellevue and the Whitewater have much less shale, placing them in facies of group 3 (3b and/or 3c). This facies division of the "Brookville formation" is useful because it identifies shale tongues and facies sequences that are significant in understanding the geologic history of the area. Finally, the Elkhorn Shale member, a tongue of facies 1a within the Whitewater Formation, is informally proposed as a member of the Whitewater Formation. In Preble County (Ohio) core 3023, the Elkhorn Shale was identified as an informal unit above the Whitewater, with undifferentiated Upper Ordovician strata above it. The nearby Richmond outcrops (see paper 15 in this volume) and the Wayne County core contain Whitewater rubbly lithology (cemented with dolomite) as well as some other facies above the Elkhorn Shale member, so it seems reasonable to extend the Whitewater Formation to the top of the Ordovician, at least in this area.

DEPOSITIONAL ENVIRONMENTS OF LITHOFACIES

Interpretation of depositional environments depends on the characteristics of individual facies and on their stratigraphic relations as revealed in transition and probability matrix analysis (Hay, 1981). The matrix analysis gave the probability that each facies would be overlain by each other facies and compared that result with the probabilities if

the facies sequence were random. This was done for one-step transitions and for reversible two-step transitions, for example, 1a overlain by 2a overlain by 1a. The geological significance of facies successions that are in conformable vertical contact is that they represent adjacent depositional environments (Walther's Principle; Blatt and others, 1972, p. 187-188). The environmental conditions responsible for some facies are relatively easy to interpret on the basis of individual facies characteristics. For others, the evidence is less clear, and interpretation must rely heavily on the association of these facies with those that are less ambiguous. This discussion of facies associations and environments is followed in the next section by isopach maps of formations and members. The combined information makes it possible to interpret paleogeography and history of these Cincinnati rocks.

INTERPRETATIONS OF FACIES OF GROUPS 3 AND 4

Figure 17-5 shows schematically the suggested lateral relations of the lithofacies based on facies associations and characteristics. The facies for which environmental interpretations are least ambiguous are those of group 4, all of




FIGURE 17-5.—Possible depositional environments and lateral relations of facies. From Hay (1981) and Hay and others (1981). Fan-shaped symbol in diagram C represents *Tetradium* colonies. Approximate lateral scale for diagrams A, B, and C is tens of miles; for diagrams D and E, a few hundred or a few thousand feet. S.L. = sea level; Saluda dol = Saluda Formation; 1a, 1b, 2a, etc. are lithofacies types diagnosed and illustrated in figure 17-2.

which probably indicate very shallow water. Facies 4d is the most significant of these because it is widespread and allows interpretation of the strongly associated facies in group 3, particularly 3c, the rubbly rock of the Whitewater Formation. Facies 4d, exposed at Garr Hill (IN-FR-0003; see paper 11 in this volume) near the top of the Brookville composite section about 4 miles (6.4 km) north of Brookville, Indiana, is a thoroughly burrowed, micritic limestone with very little shale. It contains algae in which the structure is preserved and a fauna of ostracodes and small snails but lacks the normal marine fauna of bryozoans, brachiopods, and echinoderms characteristic of most of the Cincinnati. It is underlain by massive *Tetradium* colonies that are a part of the Saluda Formation described by Hatfield (1968). Farther south and west in Indiana, facies 4d is overlain by Saluda dolomite containing mud cracks. Hatfield (1968) thought that the *Tetradium* masses ringed and acted as a wave baffle for a shallow and intermittently exposed lagoon in which the dolomite formed. The stratigraphic position and fossil characteristics of facies 4d suggest that it formed in quiet, very shallow, probably somewhat hypersaline water just lagoonward of the baffle. The Garr Hill outcrop is near the northern margin of the Saluda, and no dolomite occurs there (Hay and others, 1981). Facies 4d is overlain by, and in some cases interbedded with, facies 3c, the rubbly Whitewater; therefore, facies 3c must have formed in a laterally adjacent environment (fig. 17-5C) that was only slightly deeper, but, as the diverse fauna indicates, the water was of normal marine salinity.

Although formed in shallow water, facies 3c probably was deposited under relatively low energy conditions compared to 3b and 3a, with which it is stratigraphically associated. Both its field characteristics and limestone thin sections support this interpretation. Of the limestones studied, 60 percent of the facies 3c thin sections are micritic compared with 26 percent for facies 3a, and 71 percent of the facies 3c limestones are rudites compared with 50 percent for facies 3a. With the exception of the horn corals, which may be biochemically eroded, the fossils of facies 3c are not significantly abraded, brachiopods are commonly articulated, and molluscan steinkerns are abundant. These factors are consistent with a relatively quiet environment, as is the poor separation of the fine clastics and carbonates into discrete beds. Such conditions could occur on the interior of a shallow platform as illustrated in figure 17-5A. Marginal to this platform, under higher energy conditions, facies 3b and 3a could have formed as illustrated. This distribution is analogous to the facies distribution of the Bahama Bank (Imbrie and Purdy, 1962), where higher energy environments are marginal to the bank.

The major facies of the Bellevue is facies 3b, which has thin planar or wavy carbonate beds separated by thin fossiliferous shales and rubbly limestones. The Bellevue environment, although similar to that of the Whitewater facies 3c, was shallow but more strongly agitated, at least when the sediments of the more laterally continuous, less rubbly limestone beds were deposited.

Facies 3a, which has planar limestones and blocky or fissile, relatively unfossiliferous shales, is characteristic of the upper part of the Fairview and Liberty. This facies probably was deposited below normal wave base but periodically was subjected to agitation by storm waves and strong currents that separated the coarser biogenic debris from the fine clastics, thereby producing the limestone beds, and at least

the lower part of the overlying shale, as current velocity decreased.

The concept of storms as a mechanism for producing the interbedded limestones and shales of the Cincinnati is supported by the fact that the lower bedding surfaces of the limestone beds commonly are in sharp contact with the underlying shales, but the tops of the beds are commonly silty or gradational with the overlying shales. The siltstone beds that commonly are interbedded with shales (for example, in the Fairview and the lowest part of the Waynesville) also are likely to be the result of storms that sorted the silts rather than biogenic debris from the muddy sediments. Surely these thin siltstones are best explained as the result of localized mechanical separation of the coarser grains of the muds and not by periodic influx of silt from a distant source area. There is no evidence that individual siltstone layers thicken in any particular direction or can be traced to any source area. Therefore, the siltstones and at least many of the limestones have a similar origin. In addition, the broad lateral extent of the limestone beds compared to their thinness strongly suggests that storms played an important role in their final characteristics and burial by spreading into thin sheets any localized biogenic buildups.

Meyer and others (1981) described another possible way in which storms may have influenced the sediments. Terrigenous clastics suspended by storms may have been carried to nearby localities and deposited on top of the shelly substrate and living organisms without producing any significant reworking of the carbonate debris. The characteristics of the limestone beds in this case would depend on whether the biogenic material had been reworked prior to the event that buried it.

INTERPRETATION OF FACIES GROUPS AND THE SHALE PROBLEM

A major problem or ambiguity in facies interpretation arises when one considers the cause of changes in clastic ratios that distinguishes groups 1, 2, and 3. A change in clastic ratio can be caused by a change in the sedimentation rate of terrigenous clastics, by a change in the rate of production of carbonates, or by both. The sedimentation rate of clastics can be a function either of rate of supply from the source area or of hydrodynamic conditions in the basin. The rate of biogenic-limestone production can be influenced by water depth, substrate texture (softness), turbidity, or variation in chemical conditions such as salinity or Eh, all of which may affect the density and diversity of carbonate-producing organisms.

The question of interpretation is complicated further by uncertainty about the conditions under which muds are deposited. The traditional view for shales, such as those in the Cincinnati of Ohio and Indiana that were deposited tens to hundreds of miles from land, is that sedimentation of the muds took place under relatively low energy conditions. Pryor's (1975) work on modern muds suggests that flocculation and biogenic pelletization may cause the clays and silts to behave more like sand particles. If this idea applies to the Ordovician shales, then there may have been very little difference in the energy levels responsible for the shales and the sand-sized allochems that are common constituents of the limestones.

Another traditional view about the shale-rich intervals such as the Miamitown Shale, the Waynesville Shale

Member, and the Elkhorn Shale member is that they were deposited at greater depth than the enclosing limestone-rich intervals. This view probably is true for some of the Cincinnati shales but not for all. Each of these shale tongues needs to be interpreted individually. A facies sequence of 1a, 2a, and 3a upward such as the Kope through the Fairview, or the 2a to 3a transition upward in the Liberty, probably indicates shoaling-upward conditions. The fact that the Kope is quite uniform and thick over a large geographic area suggests a stable, relatively deep water environment. The limestone content increases near the top of the Kope where it grades into the Fairview, probably as a consequence of progressive shoaling. Figure 17-5B illustrates this interpretation of lateral relations of these facies as well as an alternative environment of very shallow water for facies 1a.

When facies 1a or 1b is in direct contact with those of group 3, it may not be appropriate to attribute the facies change to an abrupt change in water depth. The Elkhorn Shale member may be an example of this situation in which an influx of clastics, or a difference in water chemistry, rather than a change in water depth, is a more reasonable explanation for the facies change.

I believe that the contact between the Waynesville Shale Member and the "Excello member" is a disconformity; if so, the Waynesville Shale Member, at least in the lower part, probably was deposited in very shallow water. In most localities the contact between the "Excello" and the Waynesville is sharp and involves an abrupt lithologic change. The upper part of the "Excello" is nodular or rubbly limestones and limy shales of facies 1b or 2b. In many localities these facies are overlain by a few feet of facies 3a with brown, sandy-textured, friable sediment between fairly well sorted limestone beds. These brown beds owe their texture to sand-sized, phosphatized steinkerns of minute brachiopods, mollusks, and other small fossils. This band of facies 3a is commonly oxidized, even in cores; this oxidation suggests it may be the top of the regressive phase of the "Excello" and lies just below the disconformity. On the other hand, this 3a band may be the basal unit of a transgressive Waynesville cycle. The basal beds of the Waynesville are facies 1a shales with interbeds of siltstone, rather than limestone, and very few fossils. The Waynesville Shale Member becomes more fossiliferous upward. To the north, in the Randolph County, Wayne County, and Preble County (3023) cores, the contact is somewhat less sharp, but throughout the region this contact is the most consistently abrupt lithologic change in the Cincinnati. Cumings and Galloway (1913) noted an abrupt faunal change at this contact, where about 30 species of bryozoans either disappear or appear; this faunal change, they said, is greater than at any other stratigraphic horizon in the Cincinnati. The proper assignment of the 3a zone that I placed at the top of the "Excello" probably could be determined by a study of the bryozoans within it. So the interpretation of a disconformity is supported both by faunal and lithologic evidence, although the precise position of the disconformity is not certain.

If the barren shales and siltstones at the base of the Waynesville represent the basal beds of a transgression, then the sediments must have been deposited in very shallow water unless the transgression was extremely rapid. Unlike the Kope, this occurrence of facies 1a would not be a deep-water deposit. The scarcity of fossils may be due to abnormal

water chemistry or a high rate of mud sedimentation that created turbid or otherwise unstable conditions, possibly an unusually soft, soupy substrate. As the transgression proceeded, water depth increased, as did the biological diversity. The gradual increase in fossil diversity and abundance argues against a rapid transgression and deposition of the Waynesville sediments in deep water. Linguloid brachiopods are common in the lower part of the Waynesville, and their presence also suggests a shallow-water environment (Hay and others, 1981).

The fauna of the upper part of the Waynesville Shale Member (as limited here to facies 1a) has characteristics in common with the upper part of the Kope (Hay and others, 1981) and may have formed at similar depth, although this is not the only possible explanation for the similarity of the fossil assemblages. They both may have been controlled by turbidity or substrate texture rather than depth. The fine clastics of the Waynesville shales may have been derived from weathering and erosion of a low landmass exposed during the hiatus as well as from source areas to the east and southeast, the likely source of most of the Cincinnati terrigenous clastics. The disconformity may not exist in the northern cores where the contact is somewhat less abrupt than elsewhere. This contact is well exposed near the base of both the new Indiana Route 1 road cut at South Gate Hill (IN-FR-0005) and Bon Well Hill (IN-FR-0001) (see papers 10 and 12 in this volume).

In summary, the facies with >70 percent shale, those of group 1, may have formed in a variety of environments. The difference between facies 1a and 1b, which has more limy shales and/or nodular rubbly limestones, may be in the degree of effectiveness of storms in segregating the biogenic debris from the finer clastics; if so, facies 1a would have resulted when sorting was effectively accomplished. Facies 2b, for example, in the upper part of the "Excello," is quite similar to facies 3c of the Whitewater except that it contains more shale. It may be that the supply of clastics was greater when the "Excello" sediments accumulated.

REGIONAL ISOPACH AND STRUCTURE-CONTOUR MAPS

Isopach and structure-contour maps (figs. 17-6 through 17-8) show a tentative regional picture that is very likely to be revised as more information from the Ohio Division of Geological Survey is incorporated. To the extent possible based on available descriptions, these maps include information from Ohio Survey cores 2537 (D. A. Stith and E. M. Swinford, written and oral commun., 1983, 1991), 2627 (Shrake and others, 1990), 2981, 2982, 2984, and 3023. There are problems in attempting to incorporate information from these new cores that relate to illustration of isopachs for intertonguing facies and a lack of information on placement of boundaries for units within the "Brookville formation."

The structure-contour map for the base of the Fairview Formation (fig. 17-6B) shows patterns similar to those of the base and top of the "Brookville formation" (Hay, 1981) that reflect post-Ordovician deformation.

ISOPACH MAPS OF THE FAIRVIEW, MIAMITOWN, AND BELLEVUE

Figures 17-6C, 17-6D, and 17-7A are isopach maps for the Fairview Formation, the Miamitown Shale, and the




FIGURE 17-6.—Index map (A), structure and isopach maps of the Fairview Formation (B, C), and isopach map of the Miamitown Shale (D). Modified from Hay (1981). The value (85?) for core 2984 on C is plotted but not contoured.




FIGURE 17-7.—Isopach maps of the Bellevue Member of the Grant Lake Limestone/Formation (A), the "Brookville formation" (B), the "Excello member" plus "Station Hollow shales" wedges (C), and the "Station Hollow shales" wedges (D). Modified from Hay (1981). For Ohio cores, thicknesses of the "Excello member" are obtained from the sum of the Grant Lake and the Arnheim Formations, minus the Bellevue Member.




FIGURE 17-8.—Isopach maps of the Waynesville Shale Member (A), the Liberty Member plus the "Randolph County Shale member" (B), the Saluda Formation (C), and the Whitewater Formation (D). Modified from Hay (1981).

Bellevue Member of the Grant Lake Formation, respectively. Intertonguing of these units has been described in cores 2981 and 3023. For core 2981, the tongues of Fairview and Bellevue lithology within the Miamitown Shale were added to the Fairview Formation and the Bellevue Member, respectively, in making the isopach maps (figs. 17-6C and 17-7A), and the remainder was included in the Miamitown Shale (fig. 17-6D). For core 3023, the lowest 27.0 ft (8.2 meters) of Fairview lithology are put in the Fairview Formation, and the uppermost 5 ft (1.5 meters) of Fairview lithology are included in the Bellevue, which immediately overlies it and has a lower shale percentage than the Miamitown. Even including the two tongues of Fairview lithology within the Miamitown, the interval, overall, probably is still above the 70 percent shale boundary characteristic of the Miamitown elsewhere. Another problem in the Fairview isopach map (fig. 17-6C) is the thickness of 85 ft (25.9 meters) of Fairview in core 2984 compared to only 48 ft (14.6 meters) about 7 miles (11 km) to the northwest in the Middletown core (Hay, 1981) and about 70 ft (21 meters) approximately 10 miles (16 km) south in northern Hamilton County (Osborne, 1970, 1974). This apparent change in thickness may be a consequence of where the boundaries of the Fairview are placed, rather than a true stratigraphic difference. The value for core 2984 is shown on the Fairview isopach map but is not contoured.

The Fairview Formation thins toward the north and pinches out between the Wayne County and Randolph County cores (fig. 17-6C) where the Kope and the Miamitown merge. Ford (1967) commented on the steeper northward slope indicated by the structure contours on the top of the Fairview compared to the base; he thought this slope indicated a northward-sloping sea floor in the vicinity of Cincinnati. This pattern continues northward from Cincinnati; the base of the Fairview from Bellevue Hill in Cincinnati (OH-HA-0003, no. 21 on fig. 17-6A) to the Wayne County core decreases in elevation by 210 ft (64.0 meters) (fig. 17-6B), but the top of the Fairview between those localities drops 299 ft (91.1 meters). This difference is consistent with Ford's interpretation of a north-sloping ramp. The Fairview also thins somewhat toward the southeast and southwest and reaches a maximum thickness in north-central Kentucky. The base of the Fairview is probably younger toward the north.

The articulate brachiopod *Strophomena planoconvexa* occurs in the middle of the Fairview (60 ft/18.3 meters above the base) at Bedford, Kentucky, in about the middle of the formation in Cincinnati (J. K. Pope, oral commun., 1980), and probably in the middle at Aurora, Indiana, but at the base at Ripley, Ohio. The occurrence of this fossil seems to be limited to a thin stratigraphic interval and, thus, may represent a timeline; if so, then its lower position at Ripley compared with the localities farther west means that the base of the Fairview is younger at Ripley.

The Miamitown Shale (fig. 17-6D) thickens toward the northwest in the direction of the Maquoketa Shale basin that occupied much of Indiana during this part of the Cincinnati (Gray, 1972). The Miamitown isopachs are not perfectly complementary to those of the Fairview, although the two units clearly are related. The Miamitown appears to be time transgressive from the northwest toward the southeast, so that it was being deposited in the north at the same time as the upper part of the Fairview farther south. At least the lower part of the Miamitown probably represents a return to the slightly deeper water conditions

of the Kope, indicating a relatively greater subsidence of the shelf toward the northwest.

The Bellevue isopach map (fig. 17-7A) show an entirely different pattern, thick toward the southeast and southwest and thinner but of relatively uniform thickness elsewhere, implying relatively uniform depositional conditions for the Bellevue on a shallow platform. The complementary thicknesses of the Fairview and the Bellevue in the southwest and southeast may imply that the Fairview and Bellevue are partly time equivalent. So the environments of the Fairview, Miamitown, and Bellevue may have existed simultaneously in different parts of the basin. Meyer and others (1981) reached this same conclusion.

The facies sequences and isopachs together suggest this history: shoaling in the upper part of the Kope as the limestone content increases, and continuing through the Fairview, beginning in the south or southwest and progressing northward and eastward. Then followed subsidence and a return to a deeper water environment in the northwest that resulted in the lower part of the Miamitown Shale. The Bellevue lithology (facies 3b and 3c) indicates very shallow conditions. The upper part of the Miamitown may record a rather rapid shoaling that ended the northwestward and northward sloping of the sea floor while the Bellevue sediments accumulated on a shallow platform of fairly uniform depth. In the south, if the upper part of the Fairview is time equivalent to the lower part of the Bellevue in the southeast and southwest, and if the Fairview represents deeper water than the Bellevue, then there appear to have been shoals (Bellevue) in the southwest and southeast with a deeper channel (Fairview) between them. Furthermore, both the Fairview and the Bellevue could be partly time correlative with the Miamitown farther north.

ISOPACH MAPS OF THE "BROOKVILLE FORMATION"

Isopach maps of the "Brookville formation" and its members, the interval between the top of the Bellevue and the base of the Whitewater or the Saluda Formation, are shown in figures 17-7B through 17-8B. Unlike the Fairview Formation that thins northward, the "Brookville formation" is more nearly uniform in thickness but thickens toward the north and northeast. In the south-central part of the area the original thickness cannot be determined due to post-Ordovician erosion. The north-south trend of the isopachs of the "Brookville formation" in the southwest is complementary to the thickening of the Saluda Formation in that direction (fig. 17-8C).

In Indiana, if the thickness of the shaly intervals of the "Brookville formation" is subtracted from the total thickness, the trend of the isopachs changes from east-west to nearly north-south (just slightly northeast-southwest), and the unit thins westward. For example, the "Brookville formation," without the shales, thins by 50 ft (15.2 meters) between Brookville and New Point. So the northward thickening of the "Brookville formation" as a whole in Indiana (fig. 17-7B) is due to greater thickness of the shales, not to an increase in thickness of the limestone-rich intervals. Not enough information is available for Ohio to be sure what happens there.

Preble County (Ohio) core 3023 has a described thickness of only 256 ft (78.0 meters) between the top of the Bellevue and the base of the Whitewater (the "Brookville formation") compared with 301 ft (91.7 meters) just 10 miles (16 km) northwest in the Wayne County core. The

Whitewater Formation (including everything above the "Brookville formation") for core 3023 is 61 ft (18.6 meters) thicker than in outcrops near Richmond only 2 or 3 miles (3-5 km) away. So, in core 3023, the "Brookville formation" is thinner than expected, and the Whitewater is thicker than expected. These discrepancies are caused either by a difference in the placement of the "Brookville"/Whitewater boundary or there is a significant facies change over a very short distance. Thickness for this core is not contoured on the Brookville isopach map.

The "Excello member" and the "Station Hollow shales" wedges within it include the interval from the top of the Bellevue to the top of the Arnheim of Ohio nomenclature. The "Excello" isopachs (fig. 17-7C) indicate thickening toward the east and are consistent with the trend shown by Schumacher and others (1991, fig. 4C) for the Corryville Member of the Grant Lake Formation and the Grant Lake Formation as a whole in Butler and Warren Counties.

Isopachs of the "Station Hollow shales" also are plotted separately in figure 17-7D. At the present time these shale wedges have not been identified in the Ohio cores, although approximations of thicknesses are interpreted for two of the cores, 2537 (D. A. Stith and E. M. Swinford, oral and written commun., 1983, 1991) and 2627 (Shrake and others, 1990). The lower "Station Hollow shale" wedge has the same general trend as does the Miamitown, thickening toward the northwest, and is likely the result of subsidence in the northwest and a return to deeper water conditions. On the other hand, the thickening of the upper tongue toward the northeast may be the result of an increased supply of clastic sediments rather than a deepening of the sea, in view of the fact that the entire "Excello member" also thickens in that direction. This shale seems to produce a net addition of sediment to the section; it is not accompanied by thinning of the limestone-rich portions of the "Excello."

The depositional strike of isopachs for the Waynesville Shale Member in Indiana is generally northeast-southwest; the unit thickens toward the west and northwest (fig. 17-8A). There are not enough data points for confidence in the trends in Ohio at the present time, although it will be possible to get this information from the Ohio cores to clarify both this and the Liberty map (fig. 17-8B). The two points that are mapped in Ohio suggest that the Waynesville may thicken toward the northeast as well as the northwest. This wrap-around pattern, if real, suggests that the Cincinnati Arch, or a nonstructural shoal, was present and may be related to the probable disconformity at the top of the "Excello." If the Waynesville is transgressive across a disconformity, its base should be older and its thickness greater in the direction from which the transgression came, from the northwest and northeast. It would thin toward the axis of the arch, where sedimentation began later. This pattern for the Waynesville Shale Member also may suggest that the arch was present.

Isopachs for the Liberty Member plus the "Randolph County shale member" have a roughly north-south axis; the units thin toward the west and east and thicken toward the north (fig. 17-8B). The northward thickening is modified if the thickness of the "Randolph County shale member," shown in parentheses on this map, is subtracted, but the overall pattern does not change significantly. There are not enough data points for the "Randolph County shale member" to tell whether it thickens toward the northwest, north, or northeast; new data from Ohio, when incorporated, should clarify this trend. The Liberty Member thins in the south-

west owing to early onset of the conditions that produced the Saluda Formation and terminated the environment in which the Liberty formed.

ISOPACH MAPS OF THE WHITEWATER AND SALUDA FORMATIONS

The Saluda isopachs (fig. 17-8C) indicate that the Saluda Formation does not extend as far north as Richmond, although there are Saluda facies in western Butler County, Ohio. The description of Ohio core 2981 makes no mention of Saluda, but the top of the core is within the Whitewater Formation and may be below any Saluda occurrence there. Perhaps it is more likely that facies 4d, by which the Saluda is identified in that area, was not recognized in the core. This reasoning seems likely in view of the fact that 50 ft (15.2 meters) of Whitewater are described in the core, but nearby, at Camden, there are only 21 ft (6.4 meters) of Whitewater below the Saluda, and only 18 ft (5.5 meters) below it along East Fourmile Creek.

The Whitewater Formation (fig. 17-8D) is preserved only around the margins of the outcrop area and in cores in areas where it has been protected from erosion by a cover of Silurian rock. Even where protected, it is possible, or likely, that erosion between the times of deposition of the Ordovician and Silurian rocks in this area removed some of the original latest Ordovician rock. The thinning of the Whitewater in the southwest is complementary to the thickening of the Saluda there. In the north, part of the thickness of the Whitewater Formation is the Elkhorn Shale member, which is 29 ft (8.8 meters) thick in Randolph County, 30 ft (9.1 meters) thick in Wayne County, and 28 ft (8.5 meters) thick in Preble County core 3023. On the basis of only these three points, the Elkhorn Shale member may thicken toward the northeast. Farther southeast in Ohio, at Manchester, for example, the unit at the top of the Ordovician is the shale-rich Drakes Formation, which is underlain by the Bull Fork Formation. In core 2627 in Warren County, Ohio, 74 ft (25.6 meters) of Whitewater is overlain by more than 42 ft (12.9 meters) of dolomitic, silty shale assigned to the Drakes Formation. The stratigraphic relationship between the Drakes Formation in the southeast and the Elkhorn Shale Member to the north is not clear.

SUMMARY AND OVERALL INTERPRETATION OF THE ISOPACH MAPS

Inspection of these isopach maps as a series suggests that the Cincinnati Arch, or at least a north-south-trending shoal, may have developed during the Cincinnati Epoch. Of the limestone-rich units, only the Fairview Formation and the Saluda Formation thin to the north. The Bellevue, except in the southwest and southeast corners where it is quite thick, maintains a relatively uniform thickness and may mark the change from a north-sloping sea floor to a more uniform shoal. The "Excello" is thicker toward the northeast, and the Liberty is thicker northward and thins both toward the northwest and the northeast. The Whitewater thins southward, possibly owing to pre-Silurian as well as post-Silurian erosion; in the southwest the thinning of the Whitewater is complementary to the thickening of the Saluda.

The shale-rich intervals thicken toward the northwest, the northeast, or both. The Miamitown and the lower "Station Hollow shale" wedge thicken northwestward, but the

upper "Station Hollow shale" wedge thickens toward the northeast. The Waynesville Shale Member possibly thickens toward both the northwest and the northeast. The "Randolph County shale member" may thicken toward the north, the northwest, or the northeast. The Elkhorn Shale Member may thicken toward the northeast, although this trend is based on only a few data points.

The intervals with lobate contours include both shale-rich and limestone-rich units: the "Station Hollow shales," the Waynesville, the Liberty/"Randolph County," and the Saluda. All of these units have a roughly north-south-trending axis. It seems possible that this axis is caused by a north-south-trending shoal that replaced a northward slope of the sea floor that existed during deposition of the Fairview sediments. As discussed earlier, the Bellevue platform may indicate the beginning of this shoal, which became more pronounced in the lower half of the "Brookville formation," where shales thin toward the axis. This interpretation is consistent with the likely disconformity at the top of the "Excello" and its relation to the lobate isopachs of the overlying Waynesville Shale Member. This tentative interpretation is subject to confirmation or revision when the data from the Ohio Division of Geological Survey are incorporated.

SEDIMENTARY CYCLES

Figure 17-9 presents an interpretation of sedimentary cycles within the Cincinnati based on the facies sequences and isopach maps; the three localities depicted are Bedford, Kentucky (in the southwest), Manchester, Ohio (in the southeast), and the Wayne County, Indiana, core (in the north). Although the details are somewhat different, the interpretation of all three localities is that there are

three major regressive phases separated by two transgressive phases and some minor oscillations within them. The datum is the middle of the Grant Lake Formation or the Bellevue Formation. The positions of possible index fossils *Holtehdahlina*, *Strophomena concordensis*, and *Strophomena planoconvexa*, and the peak zone of *Thaerodonta* are marked.

Others have recognized cycles in the Cincinnati. For example, Weir and Peck (1968) concluded that the facies of the Grant Lake (Bellevue) indicated shallower conditions than those of the underlying Fairview, and that at least two regressive phases were evident. Fox (1962) thought the Whitewater and Saluda sediments were deposited in a more shallow sea than the underlying Tanners Creek Formation (upper part of the "Excello member" and the Waynesville Shale and Liberty Members of the "Brookville formation"). If there is a disconformity between the "Excello" and the Waynesville Shale Member, then there must have been a third transgression at the time of deposition of the Waynesville sediments; maximum depths probably are represented by facies 2a of the Liberty (fig. 17-9, Wayne County). The South Gate Hill locality paper in this volume (Hay, Kirchner, and Cuffey, paper 12) describes these cycles from the "Excello" to the Saluda Formation. The graph of Wayne County also illustrates a transgression to explain the Miamitown Shale. Hay (1981) discussed this figure in more detail.

In summary, there appear to be three regressive phases, the first and third culminating in the similar shallow-water facies of the Grant Lake/Bellevue and the Whitewater and Saluda Formations, and the second at the top of the "Excello member" of the "Brookville formation." The transgressive peak in the second cycle is within the "Excello member," within the "Station Hollow shales," or both. The peak in the third cycle may not have been synchronous throughout




FIGURE 17-9.—Transgressive and regressive cycles for the Bedford, Kentucky, Wayne County, Indiana, and Manchester, Ohio, sections. Fossils: H = *Holtehdahlina*, T = *Thaerodonta*, Sc = *Strophomena concordensis*, Sp = *S. planoconvexa*. From Hay (1981).

the region, although it is everywhere below the Whitewater and Saluda. Minor oscillations are indicated by the facies sequences, the most notable of which is the transgression that resulted in the Miamitown Shale and interrupted the major regression of the first cycle in the north.

This discussion would not be complete without considering the possibility that what appear to be transgressions and regressions in the sense of changing water depths due either to eustatic fluctuation or to differential rates of subsidence in fact may have been caused by pulses of clastic influx or progradation of clastics from the east resulting in shale wedges, separated by times of lower clastic supply in which the more limestone-rich strata were deposited. In fact, the presence or absence of carbonate sediments on modern tropical shelves seems to depend more on the supply or lack of supply of clastics rather than on small variations in water depth.

Although there may have been some such pulses of clastic influx, the nature of the depositional environments of the least ambiguous facies, those of the Bellevue, the Whitewater, and the Saluda Formations, argues in favor of cyclic changes in water depth. Compared with the Fairview Formation and the Liberty Member of the "Brookville formation," these formations were deposited in very shallow seas, so the interpretation of changing water depths seems valid. Facies 2a and 3a of the Fairview and "Liberty" have characteristics other than shale percentage that distinguish them from the Saluda, Whitewater, and Bellevue and lead to the interpretation of somewhat deeper water for these sediments.

Another possibility is that some of the shoaling could have been caused by an influx of clastics that partially filled the basin and thereby caused the depth to decrease. In view of the fact that the Miamitown Shale directly underlies or intertongues with the shallow-water Bellevue, at least the upper part of the Miamitown may represent such an influx. But the Miamitown thickens toward the northwest and the only known source of terrigenous clastics is to the east or southeast, so the Miamitown is not thicker in the direction of any known source. Sediment by-passing is a possibility. Similarly, the Elkhorn Shale member may have been produced by increased influx of clastics based on its association with the shallow-water facies of the Whitewater Formation.

A possible approach to the problem of rate of supply of muds, which bears on interpretation of changes in clastic ratios, will be available when there are time markers in which we can have confidence. The total stratigraphic thickness between timelines could be determined for different localities. If subtraction of the shale-rich intervals (facies 1a/1b) decreases the difference in total thickness for different localities, one might reasonably conclude that the shale zone was caused by a net increase in siliciclastic sedimentation rather than inhibition of carbonate-producing organisms. The thicker stratigraphic intervals between timelines would occur in those localities that also had more shale. In view of the abundance and quality of preservation of macroscopic fossils, it seems likely that modern biostratigraphic work, as well as review of the older paleontological literature, should yield good index fossils.

The cause of these transgressions and regressions may be epeirogenic, either limited to this geographic region or related to tectonism in the Appalachians, or it may be glacio-eustatic (Berry and Boucot, 1973), or both. If it is glacio-eustatic, then there should be widespread evidence

of the cycles in shallow marine sediments of other parts of the stable craton.

RELATION TO APPALACHIAN TECTONISM

It is widely accepted that the source of the clastic sediments of the Midwest was Late Ordovician erosion of tectonic highlands in the Appalachian belt (Dott and Batten, 1988). The result was formation of a clastic wedge which was deposited in a linear trough or foreland basin that was parallel to the orogenic belt and that prograded westward. Bretsky (1970) illustrated a westward migration of increasingly coarser clastics through Maysvillian and Richmondian time. This migration should have produced an increase in terrigenous clastics upward through the Cincinnati in southeastern Indiana and southwestern Ohio. In fact, that is the case near Manchester, Ohio, the easternmost locality of this study. There (fig. 17-9), the Grant Lake Limestone is succeeded by the Bull Fork Formation, which increases in shale upward beginning at the time of, or just after, deposition of the Waynesville Shale Member farther northwest. The Drakes Formation at the top of the Ordovician at Manchester, Ohio, contains very little limestone.

But just the opposite lithologic change occurs to the west, where the upper part of the "Brookville formation" and the overlying Whitewater and Saluda Formations generally contain less shale upward. This is not what we should expect unless a shoal, possibly the Cincinnati Arch, already existed marginal to the eastern basin. Such a shoal would have helped trap sediments derived from the east in the linear trough of the basin, which thus would have prevented transport onto the shoal on which carbonates could have been forming. The result would be the observed increase in limestone relative to shale upward. The north-south-trending arch or a nonstructural shoal could be an isostatic bulge at the distal margin of the migrating foreland basin (Quinlan and Beaumont, 1984). Therefore, the change to a roughly north-south axial orientation of isopachs noted in some of the units above the Bellevue may be related to tectonism in the Appalachians and development of the clastic wedge of the foreland basin. This speculative model should be tested eventually by examining deep cores or geophysical logs from localities in Ohio east of the area of this study.

LOCALITIES

REFERENCES FOR NUMBERED LOCALITIES IN FIGURE 17-1:

1. Swadley, 1972a, Lawrenceburg, Aurora, Hooven, GQ-989
2. Gibbons, 1973, Newport, Withamsville, GQ-1072
3. Gibbons, 1971, Alexandria, GQ-926
4. Gibbons and others, 1975, New Richmond, GQ-1228
5. Luft, 1970, De Mossville, GQ-862
6. Luft, 1972a, Butler, GQ-982
7. Gibbons, 1975, Worthville, GQ-1265
8. Swadley and Gibbons, 1976, Campbellsburg, GQ-1354
9. Swadley, 1976, Carrollton, GQ-1281
10. Swadley, 1973a, Vevay South, Vevay North, GQ-1123
11. Swadley, 1973b, Sanders, GQ-1095
12. Swadley, 1974, Glencoe, GQ-1154
13. Swadley, 1972b, Elliston, GQ-994
14. Luft, 1973, Williamstown, GQ-1104

15. Luft, 1972b, Falmouth, GQ-1037
16. Luft, 1975, Berlin, GQ-1256
17. Outerbridge, 1971a, Brooksville, GQ-905
18. Outerbridge, 1971b, Germantown, GQ-971
19. Gibbons and Weiss, 1972, Maysville West, GQ-1005
20. Weiss and others, 1972, Maysville East, GQ-1006
21. Ford, 1967
22. Osborne, 1968

NAMED LOCALITIES:

For details of localities followed by numbers in parentheses, see Appendix A at the end of this volume.

Aurora, Indiana (IN-D-0001).

Bedford, Kentucky. Series of road cuts along U.S. Route 42 about 2½ miles (4 km) east of Bedford and along a side road about 1 mile (1.6 km) east of the U.S. Route 42 outcrops (turn northwest on side road). NE rectangle, Bedford, Kentucky, 7.5-minute quadrangle, Trimble County, Kentucky.

Bon Well Hill (IN-FR-0001).

Brookville Dam spillway (IN-FR-0002). The described section extends from the top of the dam down to river level.

Garr Hill (Brookville North) (IN-FR-0003). Described section is on north side of the valley.

Madison, Indiana (IN-JE-0001, IN-JE-0002, IN-JE-0003). The base of the south cut is in the Bellevue Formation and the top of the north cut is Silurian.

Manchester, Ohio (in part OH-AD-0003). Road cuts along Ohio Route 136 from Manchester north to junction with Ohio Route 41.

Miamisburg, Ohio, core. From Mound Laboratory, SE corner, near Miamisburg, Ohio, 7.5-minute quadrangle; core stored at Miami University, Oxford, Ohio.

Middletown, Ohio, core. From SW¼ sec. 19, Trenton, Ohio, 7.5-minute quadrangle; core stored at Miami University, Oxford, Ohio.

New Point, Indiana, core (Indiana Geological Survey drill hole 124). Sec. 8, T. 10 N., R. 11 E., New Point, Indiana, 7.5-minute quadrangle; uppermost Ordovician is exposed in quarry at New Point.

Randolph County, Indiana, core. NW¼NW¼ sec. 16, T. 19 N., R. 13 E., Carlos, Indiana, 7.5-minute quadrangle; core stored at Ball State University, Muncie, Indiana.

Richmond/U.S. Route 27 (IN-WY-0001). See paper 15 in this volume.

Ripley, Ohio (see OH-BR-0008). Road cut about 3 miles (9.8 km) northeast of Ripley on U.S. Routes 62 and 68, SC rectangle, Russellville, Ohio-Kentucky, 7.5-minute quadrangle; the Fairview Formation is exposed just to east on Chicken Hollow Road (see OH-BR-0001) in road cut and along creek about ¼ mile (0.4 km) from U.S. Routes 62/68.

South Gate Hill (Indiana Route 1) (IN-FR-0005). See paper 12 in this volume.

Wayne County, Indiana, core (Indiana Geological Survey drill hole 57). Sec. 12, T. 14 N., R. 1 W., Whitewater, Indiana-Ohio, 7.5-minute quadrangle.

OHIO DIVISION OF GEOLOGICAL SURVEY CORES:

Core 2537. NW¼SE¼NW¼SE¼ sec. 5, T. 3 N., R. 3 E., West Elkton, Ohio, 7.5-minute quadrangle, Wayne Township, Butler County, Ohio.

Core 2627. NE¼ sec. 14, T. 3 N., R. 5 W., Waynesville, Ohio, 7.5-minute quadrangle, Wayne Township, Warren County, Ohio; in American Aggregates quarry.

Core 2981. Reily, Ohio, 7.5-minute quadrangle, Reily Township, Butler County, Ohio; Ohio Coordinates (South Zone): x = 1,346,650, y = 534,900.

Core 2982. College Corner, Ohio, 7.5-minute quadrangle, Oxford Township, Butler County, Ohio; Ohio Coordinates (South Zone): x = 1,363,700, y = 565,200.

Core 2984. Mason, Ohio, 7.5-minute quadrangle, Liberty Township, Butler County, Ohio; Ohio Coordinates (South Zone): x = 1,477,800, y = 504,275.

Core 3023. NE¼NW¼SE¼ sec. 17, T. 8 N., R. 1 E., New Paris, Ohio-Indiana, 7.5-minute quadrangle, Jackson Township, Preble County, Ohio.

REFERENCES CITED

- Berry, W. B. N., and Boucot, A. J., 1973, Glacio-eustatic control of Late Ordovician-Early Silurian platform sedimentation and faunal changes: *Geological Society of America Bulletin*, v. 84, p. 275-284.
- Blatt, Harvey, Middleton, Gerard, and Murray, Raymond, 1972, *Origin of sedimentary rocks: Englewood Cliffs, New Jersey, Prentice-Hall*, 634 p.
- Bretsky, P. W., Jr., 1970, Upper Ordovician ecology of the central Appalachians: *Peabody Museum of Natural History, Yale University, Bulletin* 34, 150 p.
- Brown, G. D., Jr., and Lineback, J. A., 1966, Lithostratigraphy of the Cincinnati Series (Upper Ordovician) in southeastern Indiana: *American Association of Petroleum Geologists Bulletin*, v. 50, p. 1018-1023.
- Caster, K. E., Dalvé, E. A., and Pope, J. K., 1955, Elementary guide to the fossils and strata of the Ordovician in the vicinity of Cincinnati, Ohio: *Cincinnati Museum of Natural History*, 47 p.
- Cressman, E. R., 1973, Lithostratigraphy and depositional environments of the Lexington Limestone (Ordovician) of central Kentucky: *U.S. Geological Survey Professional Paper* 768, 61 p.
- Cummings, E. R., and Galloway, J. J., 1913, The stratigraphy and paleontology of the Tanner's Creek section of the Cincinnati Series in Indiana: *Indiana Department of Geology and Natural Resources, Thirty-seventh Annual Report*, for 1912, p. 353-479.
- Davis, R. A., ed., 1992, Cincinnati fossils: an elementary guide to the Ordovician rocks and fossils of the Cincinnati, Ohio, region: *Cincinnati Museum of Natural History*, 61 p.
- Dott, R. H., and Batten, R. L., 1988, *Evolution of the Earth* (4th ed.): New York, McGraw-Hill, 643 p.
- Ford, J. P., 1967, Cincinnati geology in southwest Hamilton County, Ohio: *American Association of Petroleum Geologists Bulletin*, v. 51, p. 918-936.
- Fox, W. T., 1962, Stratigraphy and paleoecology of the Richmond Group in southeastern Indiana: *Geological Society of America Bulletin*, v. 73, p. 621-641.
- Gibbons, A. B., 1971, Geologic map of the Alexandria quadrangle, Campbell and Kenton Counties, Kentucky: *U.S. Geological Survey Geologic Quadrangle Map* GQ-926, scale 1:24,000.
- _____, 1973, Geologic map of parts of the Newport and Withamsville quadrangles, Campbell and Kenton Counties, Kentucky: *U.S. Geological Survey Geologic Quadrangle Map* GQ-1072, scale 1:24,000.
- _____, 1975, Geologic map of the Worthville quadrangle, north-central Kentucky: *U.S. Geological Survey Geologic Quadrangle Map* GQ-1265, scale 1:24,000.
- Gibbons, A. B., Kohut, J. J., and Weiss, M. P., 1975, Geologic map of the New Richmond quadrangle, Kentucky-Ohio: *U.S. Geological Survey Geologic Quadrangle Map* GQ-1228, scale 1:24,000.
- Gibbons, A. B., and Weiss, M. P., 1972, Geologic map of the Maysville

- West quadrangle, Kentucky–Ohio: U.S. Geological Survey Geologic Quadrangle Map GQ-1005, scale 1:24,000.
- Gray, H. H., 1972, Lithostratigraphy of the Maquoketa Group (Ordovician) in Indiana: Indiana Geological Survey Special Report 7, 31 p.
- Hatfield, C. B., 1968, Stratigraphy and paleoecology of the Saluda Formation (Cincinnatian) in Indiana, Ohio, and Kentucky: Geological Society of America Special Paper 95, 34 p.
- Hay, H. B., 1975, Lithofacies classification for the Cincinnatian Series (Upper Ordovician), southeastern Indiana: M.S. thesis (unpub.), Miami University, 147 p.
- _____, 1981, Lithofacies and formations of the Cincinnatian Series (Upper Ordovician), southeastern Indiana and southwestern Ohio: Ph.D. dissertation (unpub.), Miami University, 236 p.
- Hay, H. B., Pope, J. K., and Frey, R. C., 1981, Lithostratigraphy, cyclic sedimentation, and paleoecology of the Cincinnatian Series in southwestern Ohio and southeastern Indiana, *in* Roberts, T. G., ed., GSA Cincinnati '81 Field Trip Guidebooks: American Geological Institute, v. 1, p. 73-86.
- Imbrie, John, and Purdy, E. G., 1962, Classification of modern Bahamian carbonate sediments, in Ham, W. E., ed., Classification of carbonate rocks: American Association of Petroleum Geologists Memoir 1, p. 253-272.
- Kirchner, Brian, 1991, Description, correlation, and interpretation of Cincinnatian Series lithofacies in the Indiana State Route 1 roadcut near Brookville, Indiana: Independent study in geology, Earlham College, 42 p.
- Luft, S. J., 1970, Geologic map of the De Mossville quadrangle, north-central Kentucky: U.S. Geological Survey Geologic Quadrangle Map GQ-862, scale 1:24,000.
- _____, 1972a, Geologic map of the Butler quadrangle, Pendleton and Campbell Counties, Kentucky: U.S. Geological Survey Geologic Quadrangle Map GQ-982, scale 1:24,000.
- _____, 1972b, Geologic map of the Falmouth quadrangle, Pendleton County, Kentucky: U.S. Geological Survey Geologic Quadrangle Map GQ-1037, scale 1:24,000.
- _____, 1973, Geologic map of the Williamstown quadrangle, Grant and Pendleton Counties, Kentucky: U.S. Geological Survey Geologic Quadrangle Map GQ-1104, scale 1:24,000.
- _____, 1975, Geologic map of the Berlin quadrangle, Bracken and Pendleton Counties, Kentucky: U.S. Geological Survey Geologic Quadrangle Map GQ-1256, scale 1:24,000.
- Meyer, D. L., Tobin, R. C., Pryor, W. A., Harrison, W. B., Osgood, R. G., Hinterlong, G. D., Krumpolz, B. J., and Mahan, T. K., 1981, Stratigraphy, sedimentology, and paleoecology of the Cincinnatian Series (Upper Ordovician) in the vicinity of Cincinnati, Ohio, *in* Roberts, T. G., ed., GSA Cincinnati '81 Field Trip Guidebooks: American Geological Institute, v. 1, p. 31-72.
- Osborne, R. H., 1968, The American Upper Ordovician Standard. IX. Bedrock geology of eastern Hamilton County, Ohio: American Association of Petroleum Geologists Bulletin, v. 52, p. 2137-2152.
- _____, 1970, Bedrock geology of the Madeira quadrangle, Hamilton and Clermont Counties, Ohio: Ohio Division of Geological Survey Report of Investigations 77, 1 sheet, scale 1:24,000.
- _____, 1974, Bedrock geology of the Cincinnati East quadrangle, Hamilton County, Ohio: Ohio Division of Geological Survey Report of Investigations 94, 1 sheet, scale 1:24,000.
- Outerbridge, W. F., 1971a, Geologic map of the Brooksville quadrangle, Bracken County, Kentucky: U.S. Geological Survey Geologic Quadrangle Map GQ-905, scale 1:24,000.
- _____, 1971b, Geologic map of the Germantown quadrangle, Mason and Bracken Counties, Kentucky: U.S. Geological Survey Geologic Quadrangle Map GQ-971, scale 1:24,000.
- Peck, J. H., 1966, Upper Ordovician formations in the Maysville area, Kentucky: U.S. Geological Survey Bulletin 1244-B, 30 p.
- Pryor, W. A., 1975, Biogenic sedimentation and alteration of argillaceous sediments in shallow marine environments: Geological Society of America Bulletin, v. 86, p. 1244-1254.
- Quinlan, G. M., and Beaumont, Christopher, 1984, Appalachian thrusting, lithospheric flexure, and the Paleozoic stratigraphy of the eastern interior of North America: Canadian Journal of Earth Sciences, v. 21, p. 973-996.
- Schumacher, G. A., Swinford, E. M., and Shrake, D. L., 1991, Lithostratigraphy of the Grant Lake Limestone and Grant Lake Formation (Upper Ordovician) in southwestern Ohio: Ohio Journal of Science, v. 91, p. 56-68.
- Shrake, D. L., Wolfe, P. J., Richard, B. H., Swinford, E. M., Wickstrom, L. H., Potter, P. E., and Sitler, G. W., 1990, Lithologic and geophysical description of a continuously cored hole in Warren County, Ohio, including description of the Middle Run Formation (Precambrian?) and a seismic profile across the core site: Ohio Division of Geological Survey Information Circular 56, 11 p.
- Swadley, W C, 1972a, Geologic map of parts of the Lawrenceburg, Aurora, and Hooven quadrangles Boone County, Kentucky: U.S. Geological Survey Geologic Quadrangle Map GQ-989, scale 1:24,000.
- _____, 1972b, Geologic map of the Elliston quadrangle, Grant County, Kentucky: U.S. Geological Survey Geologic Quadrangle Map GQ-994, scale 1:24,000.
- _____, 1973a, Geologic map of parts of the Vevay South and Vevay North quadrangles, north-central Kentucky: U.S. Geological Survey Geologic Quadrangle Map GQ-1123, scale 1:24,000.
- _____, 1973b, Geologic map of the Sanders quadrangle, north-central Kentucky: U.S. Geological Survey Geologic Quadrangle Map GQ-1095, scale 1:24,000.
- _____, 1974, Geologic map of the Glencoe quadrangle, north-central Kentucky: U.S. Geological Survey Geologic Quadrangle Map GQ-1154, scale 1:24,000.
- _____, 1976, Geologic map of part of the Carrollton quadrangle, Carroll and Trimble Counties, Kentucky: U.S. Geological Survey Geologic Quadrangle Map GQ-1281, scale 1:24,000.
- Swadley, W C, and Gibbons, A. B., 1976, Geologic map of the Campbellsburg quadrangle, north-central Kentucky: U.S. Geological Survey Geologic Quadrangle Map GQ-1354, scale 1:24,000.
- Totten, S. M., and Hay, H. B., 1987, Madison, Indiana: geomorphology, and Paleozoic and Quaternary geology, *in* Biggs, D. L., ed., North-Central Section of the Geological Society of America: Geological Society of America Centennial Field Guide, v. 3, p. 365-370.
- Weir, G. W., and Peck, J. H., 1968, Lithofacies of Upper Ordovician rocks exposed between Maysville and Stanford, Kentucky: U.S. Geological Survey Professional Paper 600-D, p. 162-168.
- Weiss, M. P., Schilling, F. A., Pierce, K. L., and Ali, S. A., 1972, Geologic map of the Maysville East quadrangle, Ohio–Kentucky: U.S. Geological Survey Geologic Quadrangle Map GQ-1006, scale 1:24,000.
- Weiss, M. P., and Sweet, W. C., 1964, Kope Formation (Upper Ordovician): Ohio and Kentucky: Science, v. 145, no. 3638, p. 1296-1302.

Address of author: Helen B. Hay, deceased, formerly Geology Department, Earlham College, Richmond, IN 47374.

18. SEQUENCE STRATIGRAPHY OF THE CININNATIAN SERIES (UPPER ORDOVICIAN, CININNATI, OHIO, REGION)

by
Steven M. Holland

INTRODUCTION

Cyclicality in the Cincinnatian Series has been recognized since the earliest studies of these rocks. Orton (1878) noted several large-scale shifts in shale content within the Cincinnatian Series; these cycles were described later in more detail by many workers (most prominently by Anstey and Fowler, 1969; Hay, 1981; and Tobin 1982). These later workers argued that the Cincinnatian Series in southwestern Ohio, southeastern Indiana, and northernmost Kentucky consists of three shoaling-upward cycles characterized by systematic changes in shale percentage, bedding style, and carbonate lithology. Subsequent work in this same region, in central Kentucky, and in the Valley and Ridge Province from Tennessee to Maryland indicates that the Cincinnatian Series is composed of five sequences, each of which is dominantly shoaling upward in Ohio and Indiana (Holland, 1990, 1993). The principal features of these sequences and their implications for lithostratigraphic work are discussed in this paper.

SEQUENCE STRATIGRAPHY. A BRIEF SUMMARY

Sequence stratigraphy divides the stratigraphic record into sequences—relatively conformable successions of genetically related strata bounded by unconformities and their correlative conformities (Mitchum, 1977). Because each unconformity separates overlying strata from underlying strata that are everywhere older, the unconformities are chronostratigraphically significant surfaces. This does not imply that the magnitude of the hiatus is the same everywhere; the ages of rocks overlying or underlying the unconformity can differ considerably. This feature of unconformities allows explicit chronostratigraphic hypotheses to be framed with sequence stratigraphy, one aspect that distinguishes sequence stratigraphy from other analyses of cyclicality. Because sequences are composed of several suites of genetically related facies (these suites are called systems tracts), explicit predictions on how facies composition of a sequence changes laterally are made, thereby avoiding debates over “ideal” cycles that have plagued the cyclothem concept.

Sequence stratigraphy is undergoing rapid changes in concepts and terminology; what follows here is only a brief outline of the most important terms and concepts used in this paper (table 18-1). For a more thorough explanation, the reader is referred to Wilgus and others (1988). Sequences are characteristically tens to hundreds of meters thick and of a few million years in duration. However, these thicknesses and durations are not part of the sequence definition.

Sequence stratigraphy is not strictly a form of lithostratigraphy, nor of chronostratigraphy, nor of allostratigraphy. However, sequence stratigraphy does contain attributes of each of these forms of stratigraphy. A sequence can be thought of as a rock unit having thickness and as a time unit having a duration. Sequence stratigraphy also can be the basis of lithostratigraphic, chronostratigraphic, or allostratigraphic subdivisions. For example, the presence of unconformities can be used to define formational boundaries

(Article 23 (d) of the North American Stratigraphic Code, North American Commission on Stratigraphic Nomenclature, 1983) or to define allostratigraphic units (Article 58 (c) of the Code). If unconformities represent widespread synchronous breaks in the stratigraphic record, they are physical features that can be used to define chronostratigraphic units (Article 67 of the Code).

SEQUENCE ARCHITECTURE

The sequence boundary is a surface of subaerial erosion or exposure indicating a significant hiatus (fig. 18-1). A sequence boundary passes seaward into a correlative conformable surface bearing no significant hiatus. A sequence boundary can display channeling, erosional relief, paleokarst, or other features of subaerial exposure, but also can be a nearly planar surface lacking obvious indications of subaerial exposure or erosion. The subaqueous extension of the sequence boundary—its correlative conformity—may be recognizable only by stratal termination patterns on seismic profiles.

Lying above the sequence boundary are sediments belonging to the lowstand systems tract. The lowstand systems tract is a three-dimensional unit representing all contemporaneous depositional systems present during the lowstand. In any given outcrop, however, only a few of the lowstand depositional environments will be present. Lowstand systems tracts commonly display an overall shallowing-upward trend and are composed of individual shallowing-upward units, called parasequences. A parasequence is characterized by a flooding surface at its base, a shallowing-upward succession, and is generally 1 to 10 meters (3-33 ft) thick.

The lowstand systems tract is capped by the transgressive surface, the first significant marine-flooding surface within a sequence. A marine-flooding surface is characterized by marine deposits resting in sharp contact on nonmarine deposits or by deeper water marine deposits resting in sharp contact on shallower water marine deposits. The transgressive surface is a disconformity and commonly is demonstrably diachronous, becoming younger landward.

In shallow cratonic settings, the sequence boundary and the transgressive surface are combined into the same stratal surface, and the lowstand systems tract is absent. Where the transgressive surface and the sequence boundary are combined into a single surface, all rocks above this surface are younger than all rocks below this surface; therefore, the combined sequence boundary/transgressive surface still can be used as a chronostratigraphic marker.

Lying above the transgressive surface is a generally thin interval of transgressive deposits belonging to the transgressive systems tract. The transgressive systems tract is a three-dimensional unit representing all contemporaneous depositional systems present during the transgression. In any given outcrop, however, only one or a few of the transgressive depositional environments will be present. Transgressive systems tracts commonly display an overall deepening-upward trend, but may be composed of several parasequences.

The transgressive systems tract is capped by the maxi-

TABLE 18-1.—Terms of sequence stratigraphy¹

Condensed section	a facies consisting of thin marine beds deposited at very slow rates; commonly consists of hemipelagic or pelagic sediments
Conformity	bedding surface separating younger from older strata, along which there is no evidence of erosion or nondeposition, and containing no significant hiatus
Depositional system	a three-dimensional assemblage of lithofacies
Highstand systems tract	uppermost systems tract within a sequence; overlies maximum flooding surface and underlies sequence boundary; displays aggradational to progradational parasequence stacking
Lowstand systems tract	lowermost systems tract within a sequence; overlies sequence boundary and underlies transgressive surface; displays progradational to aggradational parasequence stacking
Marine-flooding surface	surface that separates younger from older strata, across which there is evidence of an abrupt increase in water depth; commonly accompanied by minor submarine erosion; may contain a minor hiatus
Maximum flooding surface	marine-flooding surface that marks change from retrogradational to aggradational parasequence stacking; called downlap surface by Van Wagoner and others (1988)
Parasequence	relatively conformable succession of genetically related beds bounded by marine-flooding surfaces and their correlative surfaces
Sequence	fundamental unit of sequence stratigraphy; relative conformable succession of genetically related strata bounded by unconformities (sequence boundaries) and their correlative conformities
Sequence boundary	surface separating younger from older strata, along which there is evidence of subaerial erosional truncation or subaerial exposure; may indicate a significant hiatus
Systems tract	a subdivision of a sequence, defined by types of bounding surfaces, position within sequence, and stacking patterns of parasequences. Three systems tracts are recognized: lowstand, transgressive, and highstand
Transgressive surface	first significant marine-flooding surface across the shelf within a sequence
Transgressive systems tract	middle systems tract, overlies transgressive surface and underlies maximum flooding surface; displays retrogradational parasequence stacking
Unconformity	surface separating younger from older strata, along which there is evidence of subaerial erosional truncation or subaerial exposure; contains a significant hiatus. Van Wagoner and others (1988) essentially equate the terms unconformity and sequence boundary; however, to avoid confusion with the extensive presequence literature, unconformity is used here in the traditional sense, and sequence boundaries represent one particular type of unconformity
Sequence stratigraphy	study of rock relationships within a chronostratigraphic framework of repetitive, genetically related strata bounded by surfaces of erosion or nondeposition, or their correlative conformities

¹Adapted from Van Wagoner and others (1988).

maximum flooding surface (also called the downlap surface in seismic stratigraphy). In outcrop, the maximum flooding surface is characterized by a sharp increase in water depth, in many cases signaled by a sharp increase in shale content. Thus, transgressions commonly involve two unconformities: a lower surface (the sequence boundary/transgressive surface) that indicates initial flooding and an upper surface (the maximum flooding surface) that indicates rapid deepening.

The maximum flooding surface is overlain by the highstand systems tract. The highstand systems tract is a three-dimensional unit representing all contemporaneous depositional systems present during the highstand phase of a sequence. As for the other systems tracts, only a few of the highstand depositional environments may be present within any given local section. Highstand systems tracts commonly display an overall shallowing-upward trend and are composed of numerous shallowing-upward subunits (parasequences). The terms lowstand and highstand refer only to specific phases of a sequence and do not refer to any specific altitude of sea level; for example, every sequence contains a highstand phase whether sea level is high above or far below present sea level.

A condensed section is an interval of rock representing slow net deposition and commonly is present in the transgressive systems tract and lower highstand systems tract. Although net deposition may be very slow, condensed sections can have a complicated history of deposition, non-

deposition, and erosion. Consequently a condensed section can vary widely in appearance in outcrop. Condensed sections can contain pelagic or hemipelagic deposits, abundant phosphate or glauconite, extensive burrowing, hardgrounds, anomalously abundant microfossils, or taphonomically degraded macrofossils.

SEQUENCE GENESIS

Sequences, disconformable surfaces, and systems tracts are interpreted to have formed in response to specific changes in relative sea level (fig. 18-1). Relative sea level is the sum of eustatic sea level (or sea level measured relative to some fixed reference frame) and tectonic subsidence. Note that relative sea level and water depth are not equivalent terms; for example, relative sea level could increase while water depth actually decreased—if sedimentation rate outstripped the rate of relative sea-level rise.

During times of maximum rate of relative sea-level fall, landward areas are drained and exposed to weathering, and streams incise to form a sequence boundary. The fall of relative sea level shifts deposition basinward. The lowstand systems tract forms and continues to develop as relative sea level stops falling and begins to rise slowly. Because of the low relief of cratonic areas, lowstand deposition may be confined to shelf margins and slopes, leaving the craton exposed. As the rate of sea-level rise increases, landward areas are flooded, and a transgressive surface forms, ini-




FIGURE 18-1.—Cross sections of a stratigraphic sequence, plotted in thickness (upper drawing) and geologic time (lower drawing). See table 18-1 for definitions of terms. From Haq and others (1987).

tiating the transgressive systems tract. Rising sea level traps terrigenous sediment in estuaries and lagoons as on the Atlantic Coast today. In addition, the rate of sea-level rise commonly exceeds the deposition rate for all nonreefal

carbonates, particularly where water is deeper than a few meters. Thus, in both siliciclastic and carbonate settings, rapid sea-level rise concentrates deposition in shallow nearshore and estuarine areas, but starves all other areas

of sediment. A condensed section characteristically begins to form during this time and into early highstand time. Transgressive systems tracts generally deepen upward and typically are thin because depositional rates are generally slow during a transgression. A maximum flooding surface forms and signals the beginning of highstand deposition, when the rate of sea-level rise reaches a maximum. As sea-level rise slows and eventually begins to fall, sediments are no longer hindered from dispersal to the shelf, and the areas in which condensed sections were forming retreat. Because highstand sedimentation rates are higher and because relative sea level is either slowly rising or falling, highstand systems tracts generally shallow upward.

FACIES AND DEPOSITIONAL ENVIRONMENTS

Although at first glance the Cincinnati Series may seem a monotonous pile of limestones and shales, it contains a number of discrete facies that can be defined on purely lithologic criteria, if one wishes to equate facies with formal lithostratigraphic nomenclature (discussed on p. 146). The lithologic definition of a facies can be supplemented with paleontologic criteria such as characteristic fossil assemblages, ichnology, and taphonomy to aid in field identification, particularly where an outcrop is small or poor. Of the numerous facies descriptions of the Cincinnati Series (for example, Hay, 1981; Tobin, 1982; Weir and others, 1984; Holland, 1990), most of the facies recognized in the various studies are largely synonymous (table 18-2). The descriptions of the principal Cincinnati facies are listed in tables 18-3 and 18-4. More detailed descriptions can be found in the references listed above.

HIGHSTAND SYSTEMS TRACT FACIES

The highstand facies consistently are stacked into conformable, shallowing-upward hemicycles, such that a deeper water facies grades upward into a shallower water facies. Where a deeper water facies overlies a shallower water facies, the contact is invariably sharp, and the two highstand facies commonly are separated by a very thin transgressive facies. The consistent stacking of highstand facies into a repeated vertical shallowing-upward succession suggests highstand deposition along a series of shore-parallel belts on a gently sloping, mixed-carbonate-clastic ramp. The northward pinchout of supratidal and intertidal facies and the southward pinchout of deeper subtidal facies indicate a northward-dipping ramp. Megaripples, groove casts, gutter casts, and aligned fossils also indicate northward-directed storm flow downramp (Hofmann, 1966; Meyer and others, 1981; Jennette, 1986; Duke, 1990).

Highstand facies form the bulk of the Cincinnati Series, and the subtidal highstand facies that dominate the Cincinnati Series in its type area are quite similar to one another at first glance. Closer examination reveals numerous and in some cases subtle differences between facies. Several depth trends are apparent within the highstand facies (table 18-3). The deepest subtidal facies exposed at the surface (the shale-dominated facies) contains abundant shale and dominantly fine grained carbonate lithologies. Fossil morphologies tend to be thin and fragile (for example, the brachiopods *Resserella* and *Sowerbyella*), and multi-element skeletons of arthropods and echinoderms commonly are articulated. (Note: some workers prefer the name *Onniella* instead of the name *Resserella* for the forms

found in the type-Cincinnati.) Shell breakage, abrasion, encrustation, and boring are generally low. Bioturbation is limited to discrete traces of the *Cruziana* ichnofacies. Deep-water storm beds are dominated by distal types, characterized by relatively thin, basal shell lags; thick, laminated to hummocky, cross-stratified siltstones; and thick capping shales (Aigner, 1985). Collectively, these features suggest offshore deposition in generally quiet water disturbed only by major storms.

The mixed packstone-shale facies is intermediate between the shale-dominated facies and the wavy/nodular limestone facies in nearly all aspects, including shale content, carbonate grain size, fossil morphologies, taphonomy, ichnofacies assemblage, and storm-bed structure. Peak species diversity and morphological variability are reached in this facies. Given its intermediate character and stratigraphic position, the mixed packstone-shale facies was deposited within the transition zone, where the water was agitated frequently by storms.

The shallowest subtidal facies (wavy/nodular limestone) is characterized by less shale. Carbonate lithologies are commonly coarse grained. Fossil morphologies are characteristically thick and robust (for example, the articulate brachiopods *Platystrophia* and *Hebertella*, and thick ramose to massive trepostome bryozoans), and multi-element skeletons are more rarely articulated than in deeper water facies. Shell breakage, abrasion, encrustation, and boring are variable, but commonly are more intense than in deeper water facies. Bioturbation is pervasive, and the few discrete traces are elements of the *Skolithos* ichnofacies. Shallow-water storm beds are dominated by proximal types characterized by an upward succession of thick basal shell lags, thin or absent laminated siltstones, and thin fossiliferous capping shales (Aigner, 1985). These features, taken together with the stratigraphic position of this facies, suggest deposition in the shoreface zone. However, the presence of an abundant, sessile, soft-bottom epifauna and thoroughly bioturbated sediments indicates that this environment was not subject to large breaking waves, but instead was a fairly low energy shoreface.

The intertidal and supratidal portions of the shallowing-upward highstand deposits are not well exposed in most of Ohio and Indiana. Intertidal facies of the Cincinnati Series are characterized by a pervasively bioturbated fabric and generally lack abundant and diverse body fossils; ramose trepostome bryozoans are the most abundant form. The supratidal facies contains abundant desiccation cracks and parallel lamination and uncommon vertical burrows, but lacks bioturbated texture and body fossils.

TRANSGRESSIVE SYSTEMS TRACT FACIES

Transgressive facies in the Cincinnati Series are unusual in several respects. They are generally thin, always less than 5 meters (16.4 ft) thick, and generally less than 2 meters (6.6 ft) thick. They are geographically restricted, commonly traceable for only tens of kilometers rather than hundreds. Although highstand facies are generally similar to one another, transgressive facies are unlike each other or any highstand facies in their combination of grain size, sedimentary structures, bedding, faunal composition, and taphonomy. Although contact relationships between successive highstand facies are highly predictable, contact relationships between transgressive facies and overlying and underlying highstand facies are less predictable.

TABLE 18-2.—*Synonymy of Cincinnati Series facies and lithostratigraphic nomenclature*

Facies of this paper and Holland (1990)	Facies of Tobin (1982), Tobin (1986)	Facies of Hay (1981), Hay and others (1981)	Facies of Weir and others (1984)	Previous lithostratigraphic nomenclature
Laminated Carbonate Mudstone (LCM)	Facies D	did not name	M. Dolomite and calcitic dolomite	Parts of Saluda, Preachersville, Rowland, Terrill, and Tate
Bioturbated Carbonate Mudstone (BCM)	Facies D	did not name	K. Dolomitic mudstone L. Dolomitic mudstone and dolomite	Parts of Saluda, Preachersville, Rowland, Terrill, and Tate
Variegated Mudstone (IVM)	Facies D	tentatively assigned to 1a	not studied	Parts of Preachersville and Elkhorn
Wavy/Nodular Limestone (WNL)	Facies C	3a, 3b, 3c, 3d	E. Nodular-bedded fossiliferous limestone and shale F. Nodular-bedded limestone and calcarenite G. Nodular-bedded mudstone	All of Whitewater, Oregonia, Reba, Mt. Auburn, Straight Creek, Stingy Creek, and Bellevue; parts of Calloway Creek, Grant Lake, and Bardstown
Mixed Packstone-Shale (PS)	Facies B	2a, 2b	A. Thin-bedded fossiliferous limestone B. Even-bedded fossiliferous limestone and shale C. Even-bedded fossiliferous limestone and shale	All of Liberty, "Sunset" (of Tobin, 1986, not Weir and others, 1984), Corryville, Fairview, and Clays Ferry; parts of Calloway Creek, Grant Lake, and Bardstown
Shale-Dominated (SD)	Facies A	1a, 1b	D. Shale and fossiliferous limestone	All of Waynesville and Kope
Wavy-Bedded Wackestone (WBW)	Facies D	4d	I. Micrograined limestone N. Dolomitic limestone and mudstone	All of Hitz Bed, Sunset (of Weir and others, 1984, not Tobin 1986), and Gilbert; parts of Saluda and Marble Hill Bed
Cross-Bedded Calcarenite (CBC)	not encountered in study	4a, 4b	F. Nodular-bedded limestone and calcarenite (in part) H. Calcarenite	Never separately named, generally incorporated in WNL units
Bioclastic Packstone (BP)	not distinguished as separate facies	3a	not distinguished as separate facies	Never separately named, generally incorporated in WNL units
Gastropod quina (GC)	not encountered in study	4c	not separately named as facies	Parts of Marble Co-Hill Bed

Transgressive facies always lie between overlying highstand facies that formed in shallower water than underlying highstand facies; in other words, they always occur in a transgressive position.

Because of the thinness of the transgressive systems tract, geographic relationships of the transgressive facies are much better known than the stacking relationships. The

transgressive systems tract is always thin, but is thicker in landward areas characterized by the wavy-bedded wackestone facies (table 18-3). The wavy-bedded wackestone facies (WBW) is characterized by bioturbated wackestones with pinch-and-swell bedding. Faunally, WBW is dominated by a stromatoporoid-tabulate coral-molluscan assemblage, but lacks abundant typical open-marine fauna such as bra-

TABLE 18-3.—*Highstand facies*

Lithofacies	Lithologies	Bedding ¹	Sedimentary structures	Body-fossil assemblage	Taphonomy	Trace fossils
Supratidal Laminated Carbonate Mudstone (LCM)	calcareous to dolomitic mudstone	planar lamination, even and regular	ubiquitous planar lamination, wave-ripple lamination, abundant desiccation cracks	absent	—	short <i>Skolithos</i> ?
Intertidal Bioturbated Carbonate Mudstone (BCM)	calcareous to dolomitic mudstone	very thin to medium bedded, unevenly but regularly bedded	rare desiccation cracks, wave-ripple lamination	scarce; thick ramose bryozoans, locally a rugosan-stromatoporoid assemblage	calcite forms preserved or locally dolomitized; moderate to severe breakage	discrete burrows to pervasive bioturbation; <i>Chondrites</i> , <i>Palaeophycus</i>
Intertidal Variegated Mudstone (IVM)	mudstone (95%), siltstone or dolomitized wackestone/packstone (5%)	indistinct in mudstones; thin to very thin siltstones, wackestones, and packstones	ripple-lamination	elements of the <i>Hebertella</i> assemblage: <i>Hebertella</i> , ramose trepostome bryozoans, cyclostome bryozoans	recrystallized and dolomitized calcitic forms; moderate breakage	pervasive bioturbation; <i>Palaeophycus</i> in siltstones
Shoreface Wavy/Nodular Limestone (WNL)	fossiliferous packstone (70%), fossiliferous mudstone (30%); minor wackestone, grainstone, and calcisiltite	very thin bedded, evenly but irregularly bedded	proximal storm beds, wave-ripple lamination, gutter casts	<i>Hebertella</i> assemblage: large thick-shelled brachiopods (<i>Hebertella</i> , <i>Platystrophia</i> , <i>Rafinesquina</i>); common thick ramose and massive bryozoans; also pelecypods, gastropods, cephalopods, crinoids, and trilobites	calcite shells well preserved in both limestones and shales; aragonitic forms preserved as molds; breakage and abrasion variable, but generally moderate to severe; widespread boring and encrustation	pervasive bioturbation; <i>Skolithos</i> assemblage: <i>Skolithos</i> , uncommon cone-shaped hypichnia
Transition Zone Mixed Packstone-Shale (PS)	fossiliferous packstone (45%), mudstone (45%), wackestone, grainstone, and calcisiltite (10% total)	medium to very thin, evenly bedded, meter-scale cycles	storm beds, planar laminations, wave-ripple lamination, small-scale hummocky cross-lamination	mixed brachiopod-bryozoan assemblage: wide range of brachiopod and bryozoan taxa and morphologies; also pelecypods, gastropods, trilobites, and echinoderms	calcite shells preserved, aragonitic forms preserved as molds; negligible to moderate breakage and abrasion (rarely severe); common articulated multi-element skeletons	mixed <i>Cruziana-Skolithos</i> ichnofacies; <i>Chondrites</i> , <i>Diplocraterion</i> , <i>Trichophycus</i> , <i>Palaeophycus</i>
Offshore Shale-Dominated (SD)	mudstone (70%), fossiliferous wackestone and packstone (20%), crinoidal grainstone (5%), calcisiltite (5%)	thin to thick, uneven to cyclical meter-scale cycles	storm beds, small-scale hummocky and trough cross-lamination, megaripples, gutter casts, groove casts, runzelmarken	<i>Onniella</i> assemblage: small, thin, flat brachiopods (<i>Onniella</i> , <i>Sowerbyella</i> , small <i>Rafinesquina</i>); also ramose bryozoans, trilobites (including blind forms such as <i>Cryptolithus</i>), crinoids, graptolites	calcite shells preserved, aragonitic forms preserved as molds; variable shell breakage and low shell abrasion; common to abundant articulated multi-element skeletons	<i>Cruziana</i> ichnofacies; <i>Chondrites</i> , <i>Diplocraterion</i> , <i>Trichophycus</i> , <i>Rasophycus</i> , <i>Diplichnites</i>

¹Bedding terminology of Ingram (1954).

TABLE 18-4.—*Transgressive facies*

Lithofacies	Lithologies	Bedding ¹	Sedimentary structures	Body-fossil assemblage	Taphonomy	Trace fossils
Wavy-Bedded Wackestone (WBW)	wackestone to micstone (80%), mudstone (20%)	medium wavy-bedded carbonates, regular but uneven; very thin mudstone beds	normal grading; planar lamination	stromatoporoid-tabulate coral-molluscan assemblage, including ramose bryozoans and ostracodes; grades upward into <i>Hebertella</i> assemblage	both calcitic and aragonitic bioclasts preserved intact or recrystallized; breakage and abrasion low	<i>Skolithos</i> assemblage to burrow-mottled <i>Skolithos</i>
Cross-Bedded Calcarenite (CBC)	calcarenite (90%), fossiliferous mudstone (10%)	medium- to thick-bedded carbonates, irregular to broadly lenticular beds, sigmoidal shaped; very thin mudstone beds	low-angle tabular cross-bedding, trough cross-bedding, erosional base to facies	thick-shelled elements of <i>Hebertella</i> assemblage	comminuted bioclasts; highly abraded, broken and bored bioclasts; some brachiopod shells phosphatized	absent
Bioclastic Packstone (BP)	bioclastic wackestone and packstone (80%), fossiliferous mudstone (20%)	medium bedded, irregular and uneven, lenticular in places	megaripples, cross-lamination, amalgamation	variable; generally contains deepening-upward sequence from <i>Hebertella</i> assemblage to mixed brachiopod-bryozoan assemblage to <i>Onniella</i> assemblage	highly abraded, broken and bored, but not as intensely as in cross-bedded calcarenite facies; phosphatic steinkerns of gastropods, pelecypods, and bryozoans	absent
Gastropod Coquina (GC)	gastropod grainstone (50%), crinoidal calcarenite (50%)	medium to thick bedded, tabular, even, regular	cross-lamination	almost entirely the gastropod <i>Loxoplocus</i> and crinoid ossicles; fragments of bryozoans and <i>Hebertella</i>	gastropods largely unbroken and unabraded, shell material dolomitized and lined internally and externally with calcite; other elements highly abraded and broken	<i>Skolithos</i> locally

¹Bedding terminology of Ingram (1954).

chiopods, echinoderms, and trilobites. These features and the paleogeographic setting of WBW in the most upramp areas are consistent with quiet shallow-water deposition, such as lagoons.

Wavy-bedded wackestones are bounded downramp by the cross-bedded calcarenite facies (CBC). This facies contains abundant sand-wave-scale cross-bedding and is dominated by echinodermal grains, highly abraded brachiopods, and massive bryozoans. CBC is limited to narrow (approximately 10 km/6 miles) shore-parallel belts, suggesting deposition as a series of bars or shoals, which may have protected the lagoons from wave energy.

The cross-bedded calcarenite facies is bounded downramp by very widespread but thin deposits of bioclastic packstones (BP). This facies is characterized by a consistently taphonomically degraded fauna that is intensely broken and abraded. BP is commonly enriched in resistant echinodermal grains and in phosphatic steinkerns of molluscan protoconchs and bryozoan zooecia. The bioclastic packstone lithology is not unusual in the Cincinnati Series and is present in nearly all facies, but the bioclastic packstone facies is distinctive in that it commonly occurs just above the sequence boundary in downramp sections and appears in outcrop as an anomalous band of tabular limestones.

The gastropod coquina is an unusual facies in the Cincinnati Series; it occurs at the base of the C4 sequence (see below) over a few square kilometers near Bedford, Kentucky. It consists of a series of decimeter-scale fining-upward units that grade from gastropod coquina into crinoidal grainstones. Faunally, it is dominated by abundant, nearly unbroken high-spined gastropods of the genus *Loxoplocus* and by echinodermal grains. Brachiopods and bryozoans occur sparingly and are highly abraded. On the basis of its geometry as a single, narrow, shore-parallel body and three shore-perpendicular tongues, the gastropod coquina is interpreted as a bar and tidal-channel system (Swadley, 1979).

Florida Bay is a possible modern analog to the Cincinnati Series transgressive systems tract. Florida Bay is an expanse of subtidal to supratidal seagrass beds that currently are forming mottled wackestones (Parkinson and Meeder, 1989). These seagrass beds are bounded seaward by a series of islands composed of coarse shell debris; these islands are bounded seaward by extensive poorly sorted shelly sand bottoms (Aigner, 1985).

Of the transgressive facies, only the bioclastic packstone facies is widely distributed in Ohio and Indiana. The wavy-bedded wackestone facies, cross-bedded calcarenite facies, and gastropod coquina facies are best developed in upramp areas of central Kentucky.

DESCRIPTIONS OF CINCINNATIAN SEQUENCES

The character of the Cincinnati sequences changes laterally, especially into upramp areas of central Kentucky (fig. 18-2). However, in southeastern Indiana, southwestern Ohio, and northernmost Kentucky, the character of each of the Cincinnati sequences is remarkably constant (figs. 18-3 and 18-4). The Cincinnati sequences are numbered sequentially from oldest (C1) to youngest (C5). Mohawkian sequences are currently under study and will be similarly numbered, with an M prefix; preliminary results suggest the presence of three Mohawkian sequences in the Cincinnati Arch, Nashville Dome, Valley and Ridge, and Upper Mississippi Valley areas.




FIGURE 18-2.—Map showing locations of sections in figures 18-3 and 18-4. A, Aiken High School (OH-HA-0012); AC, Ashlock Cemetery; B, Bedford; BN, Brookville North (= Garr Hill, IN-FR-0003); BRR, Blue Rock Road (OH-HA-0005); BWH, Bon Well Hill (IN-FR-0001); By, Bybee; CC, Caesar Creek spillway and dam (OH-WA-0001, OH-WA-0002); CF, Clays Ferry; CL, Cowan Lake (OH-CN-0001); FI, Flemingsburg; Fr, Frederickstown; G, Georgetown (OH-BR-0006); HGN, Halls Gap North; LCL, Lincoln County Line; M1, Madison #1 (IN-JE-0003); M3, Madison #3 (IN-JE-0001); O, Orphanage Road (KY-KE-0003); PP, Point Pleasant (OH-CT-0020); R, Richmond (= Richmond-U.S. Route 27, IN-WY-0001); RR, Riedlin Road (KY-KE-0001); SGH, South Gate Hill (IN-FR-0005); WU, West Union (OH-AD-0004); WBW, Wright Brothers Memorial. See section on Localities (p. 149-150) for descriptions.

C1 SEQUENCE

The C1 sequence is the longest and thickest of the Cincinnati sequences. Spanning all of the Edenian Stage and most of the Maysvillian Stage, it represents some 6 million years. The C1 sequence is approximately 105 meters (344 ft) thick at Cincinnati and thins southward into central Kentucky.

In the type-Cincinnati area, the C1 sequence is composed in vertical sequence of the Kope, Fairview, and Bellevue Formations. The C1 sequence boundary is placed at the contact between the Kope Formation and the Point Pleasant Formation and is well exposed in stream cuts near Point Pleasant, Ohio (OH-CT-0020), and in road cuts along U.S. Route 52 near the Beckjord Coal Plant (OH-CT-0019). Here, the shale-dominated facies belonging to the Kope Formation overlies mixed packstone-shale facies of the Point Pleasant Formation. A 1-meter-thick (3.3-ft-thick) interval dominated by cross-bedded and megaripped packstones at the top of the Point Pleasant may constitute the bioclastic packstone facies of the transgressive systems tract. The water depths near the base of the Kope Formation are probably the greatest achieved in the type-Cincinnati, as the fauna is characterized by abundant graptolites and the deep-water trilobite *Triarthrus*. The Kope Formation grades upward into the mixed packstone-shale facies of the Fairview For-

mation, which in turn grades upward into the wavy/nodular limestone facies of the Bellevue Formation. This shoaling-upward succession is well exposed in long road cuts along Interstates 275 and 471 in northern Kentucky. Overall, the C1 sequence at Cincinnati is dominated by the shoaling-upward succession from offshore to shoreface environments.

In central Kentucky, the C1 sequence boundary is placed at the sharp contact between the mixed packstone-shale facies of the Clays Ferry Formation and the cross-bedded calcarenites and wavy/nodular limestones of the Lexington Limestone. The C1 sequence boundary is well exposed along Kentucky Route 2328 at Clays Ferry, Kentucky. As at Cincinnati, a thin interval of megarippled and taphonomically degraded packstones at the contact may represent the bioclastic packstone facies of the transgressive systems tract. Alternatively, the transgressive systems tract may be absent here, in which case the sequence boundary, transgressive surface, and maximum flooding surface are combined into a single stratal surface. The top of the Clays Ferry Formation is marked by the Garrard Siltstone, a thick interval of convolute-bedded siltstones. The Garrard Siltstone may correlate lithostratigraphically and chronostratigraphically to thinner convolute-bedded siltstones with well-developed ball-and-pillow structures exposed from Maysville, Kentucky, to Batavia, Ohio (for example, Maysville/KY-MS-0002 and Chicken Hollow Road/OH-BR-0001 localities). These thinner siltstone units lie within the Fairview Formation and occur in approximately the same stratigraphic position as the Garrard Siltstone. Above the Garrard Siltstone is the mixed packstone-shale facies of the lower Calloway Creek Formation, which grades upward into the wavy/nodular limestones of the upper Calloway Creek Formation, which grade upward into the bioturbated and laminated calcareous mudstones of the Tate Formation. The C1 sequence in central Kentucky is dominated by a shallowing-upward succession from transition zone to supratidal environments, a shallower suite of environments than at Cincinnati.

C2 SEQUENCE

Spanning the upper part of the Maysvillian Stage, the C2 sequence represents only 1 million years. At Cincinnati, the C2 sequence is approximately 12 meters (39 ft) thick. The C2 sequence thins to the south; in central Kentucky, it is approximately 10 meters (33 ft) thick.

In the type-Cincinnati area, the C2 sequence boundary is placed at the contact between the mixed packstone-shale facies of the Corryville Formation and the wavy/nodular limestone facies of the Bellevue Formation. The character of the sequence boundary changes across the Cincinnati area. At Madison, Indiana (IN-JE-0001, IN-JE-0002, and IN-JE-0003), the sequence boundary is poorly exposed, but seems to lack any transgressive deposits such as bioclastic packstones. At Muddy Creek in Cincinnati (OH-HA-0020), the sequence boundary is marked by a single thick bed of bioclastic packstone at the sharp facies contact between the Corryville and Bellevue. Near Georgetown, Ohio (OH-BR-0006), the sequence boundary is not as sharp, but is placed at the sharp change from nodular bedding to tabular bedding. Several hardgrounds are present at this bedding change. The Corryville Formation grades upward into the wavy/nodular limestones of the Mt. Auburn Formation. The overall succession of the C2 sequence at Cincinnati passes from transition zone to shoreface environments.

In central Kentucky, the C2 sequence boundary is placed

at the sharp contact between the wavy-bedded wackestones of the Gilbert Formation and the laminated carbonate mudstones of the Tate Formation. The sequence boundary here is unremarkable except for its sharpness; it is not burrowed, enriched in glauconite or phosphate, nor does the surface bear more than a few centimeters of relief or obvious evidence of karstification. The wavy-bedded wackestones grade rapidly over a few tens of centimeters into wavy/nodular limestones of the Stingy Creek Formation; this zone of rapid gradation is equivalent to the maximum flooding surface. The Stingy Creek Formation grades upward into the bioturbated and laminated carbonate mudstones of the Terrill Formation. The overall succession of the C2 sequence in central Kentucky passes from lagoonal transgressive deposits upward into shoreface and supratidal environments. The C2 sequence in central Kentucky contains a thicker transgressive systems tract than at Cincinnati and a shallower suite of highstand facies.

C3 SEQUENCE

The C3 sequence represents slightly more than 1 million years and is limited to the basal Richmondian Stage. The C3 sequence is approximately 20 meters (66 ft) thick at Cincinnati and thins southward to 15 meters (49 ft) in central Kentucky.

In the type-Cincinnati area, the C3 sequence boundary is placed at the contact between the mixed packstone-shale facies of the "Sunset Formation" and the wavy/nodular limestone of the Mt. Auburn Formation (see section on Comments on Cincinnati lithostratigraphic nomenclature for discussion of Sunset usage). The C3 sequence boundary is well exposed at Blue Rock Road (OH-HA-0005) and is marked by a single thick bed of medium-grained bioclastic packstone at the sharp facies contact. The C3 sequence boundary is poorly exposed at Madison, Indiana (IN-JE-0001, IN-JE-0002, and IN-JE-0003), where transgressive deposits appear to be absent. The mixed-packstone shale facies grades upward into the wavy/nodular limestone facies of the Oregon Formation. Note that the C3 sequence in the type-Cincinnati corresponds to the entire Arnheim Formation, originally composed of the Sunset and Oregon Members. The C3 sequence in the type-Cincinnati is dominated by a shoaling-upward succession from transition zone to shoreface environments.

In central Kentucky, the C3 sequence boundary is marked by the sharp contact between the wavy-bedded wackestones of the Sunset Formation and the laminated carbonate mudstones of the Terrill Formation. The sequence boundary is well exposed at the Lincoln County Line locality, where it is sharp but otherwise unremarkable. The C3 sequence boundary also is well exposed at Frederickstown. Somewhat to the north at Bedford and Flemingsburg, the C3 sequence boundary is placed at the base of cross-bedded calcarenites, which are sharply overlain by wavy/nodular limestones of the Reba Formation. The wavy-bedded wackestones grade over several tens of centimeters into the wavy/nodular limestones of the Reba Formation, which grade upward into the bioturbated and laminated carbonate mudstones of the Rowland Formation. The C3 sequence in Kentucky is very similar in architecture to the C2 sequence in Kentucky; it consists of transgressive lagoonal deposits overlain by a shallowing-upward highstand succession passing from shoreface into supratidal environments. C3 transgressive deposits are thicker in central Kentucky than near Cincinnati.




FIGURE 18-3.—Measured sections of the northern Cincinnati Arch. See figure 18-2 for locations and figure 18-4 for explanation of facies patterns. Sections generally are arrayed from south to north, from left to right. Sequence boundaries are indicated by horizontal lines connecting sections. Because of the thinness of the transgressive units, correlations of the maximum flooding surface are not drawn; at this scale, they commonly would be plotted on the same line as the sequence boundaries. Where the transgressive units are thick (for example, Caesar Creek dam and South Gate Hill), the maximum flooding surface lies at the upper surface of the transgressive facies. Letters C1-C5 along left side indicate sequence number; M = an unnumbered Mohawkian sequence, and S = Silurian rocks.

nati, and a shallower suite of highstand environments is developed in Kentucky.

C4 SEQUENCE

The C4 sequence represents just over 2 million years and spans the majority of the Richmondian Stage. The C4 sequence is approximately 60 meters (197 ft) thick near the type area of the Richmondian Stage at Richmond, Indiana, and thins southward to central Kentucky, where it is 11 meters (36 ft) thick.

In the type-Cincinnati area, the C4 sequence boundary is placed at the base of the 1.5-meter (5-ft) interval of bioclastic packstones that are traditionally regarded as the uppermost part of the Oregonia Formation. These bioclastic packstones are well exposed at Bon Well Hill (IN-FR-0001) and Caesar Creek spillway (OH-WA-0001) and dam (OH-WA-0002) and consist of taphonomically degraded brachiopods and bryozoans in a disarticulated-echinoderm-rich packstone. The packstone beds are rich in phosphatic steinkerns of mollusks and bryozoan zooecia, are cross bedded, and display an overall faunal deepening-upward sequence passing from a shallower, *Rafinesquina*-dominated assemblage into a deeper water, *Resserella*-dominated assemblage. (Note: some workers prefer the name *Onniella*

instead of the name *Resserella* for the forms found in the type-Cincinnati). The upper surface of this packstone interval is a sharp contact with the overlying shale-dominated facies of the Waynesville Formation; this contact represents the maximum flooding surface. The shale-dominated facies of the Waynesville grades upward into the mixed packstone-shale facies of the Liberty Formation, which grades upward into the wavy/nodular limestone facies of the Whitewater Formation. On the western side of the Cincinnati Arch, the Whitewater (known in older usage as the Lower Whitewater) grades upward into the bioturbated and laminated carbonate mudstone facies of the Saluda Formation. The bioturbated carbonate mudstones contain prominent heads of the tabulate coral *Tetradium*. On the eastern side of the Cincinnati Arch, the C4 interval is poorly exposed and many of the lithofacies are somewhat shalier, but show the same shoaling-upward faunal succession passing from an offshore *Resserella*-dominated assemblage to a transition-zone mixed brachiopod-bryozoan assemblage to a shoreface *Hebertella*-dominated assemblage. Although its assignment to the C4 sequence is tentative, the intertidal variegated mudstone facies is present on the east side of the Cincinnati Arch at West Union (OH-AD-0004). Despite the differences between facies on the east and west sides of the Cincinnati Arch, the overall C4 succession passes from a thin zone of transgres-




FIGURE 18-4.—Measured sections of the southern Cincinnati Arch. See figure 18-3 for explanation. D above Halls Gap North section indicates Devonian rocks (New Albany Shale).

sive bioclastic packstones into highstand facies that shallow upward from offshore to intertidal environments.

In central Kentucky, the C4 sequence and its sequence boundary are variable, suggesting somewhat greater paleotopography. At Bybee, the sequence boundary is placed at the sharp contact between bioturbated carbonate mudstones of the Preachersville Formation and laminated carbonate mudstones of the Rowland Formation. The sequence passes upward into the variegated mudstone facies and back into the bioturbated carbonate mudstone facies, both of the Preachersville Formation, but otherwise shows little environmental change. At Frederickstown, the sequence boundary is more obvious and is placed at the sharp contact between laminated carbonate mudstones of the Rowland Formation and 1 meter (3.3 ft) of bioclastic packstones of the Bardstown Member of the Drakes Formation. These bioclastic packstones are sharply overlain by wavy/nodular limestones of the Bardstown Member; this contact represents the maximum flooding surface. The wavy/nodular limestones grade upward into bioturbated and laminated

carbonate mudstones of the Saluda Formation. At Bedford, the C4 sequence boundary is marked by the base of the only deposit of gastropod coquina in the Cincinnati; this gastropod coquina is named the Marble Hill Bed, and it sharply overlies bioturbated carbonate mudstones of the Rowland Formation. The gastropod coquina is sharply overlain by wavy-bedded wackestones, which are sharply overlain by mixed packstone-shale facies of the Liberty Formation; the sharp contact at the base of the Liberty is the maximum flooding surface. The Liberty grades upward into wavy/nodular limestones of the Whitewater Formation, which grades upward into bioturbated and laminated carbonate mudstones of the Saluda Formation. At Flemingsburg, the C4 sequence boundary is obscure and poorly exposed; it is placed at a horizon marking an increase in shale and the first occurrence of a number of distinctive C4 taxa, including the rugose coral *Grewingkia* and *Streptelasma* and the articulate brachiopods *Thaerodonta*, *Glyptorthis*, and *Hisco-beccus*. (Note, some workers assign *Hisco-beccus* to the genus *Lepidocyclus*.) Overall, the C4 sequence in central Kentucky

shows only localized transgressive deposits and variable amounts of shoaling during the highstand, but, in general, a shallower suite of facies than at Cincinnati.

C5 SEQUENCE

The C5 sequence is the thinnest and briefest of the Cincinnati sequences. Limited to the uppermost Richmondian, it represents fewer than a half a million years, although some strata may have been removed by latest Ordovician and earliest Silurian erosion. The C5 sequence is 11 meters (36 ft) at its thickest measured section at the Wright Brothers Memorial and pinches out to the south in northern Kentucky.

The C5 sequence boundary is placed at the base of wavy-bedded wackestones of the "Saluda Formation" and the Hitz Bed, where they overlie bioturbated and laminated carbonate mudstones of the true Saluda Formation. The C5 sequence boundary is well exposed at Madison (IN-JE-0001, IN-JE-0002, and IN-JE-0003) and at Garr Hill (= Brookville North, IN-FR-0003). The wavy-bedded wackestones are sharply overlain by wavy/nodular limestones of the Upper Whitewater Formation (as known in older literature); this contact is the maximum flooding surface. In northernmost outcrop areas, the wavy/nodular limestones grade upward into the shale-rich facies of the Elkhorn Formation, which is overlain by the Silurian Brassfield Formation.

The C5 sequence boundary also is well exposed at the Wright Brothers Memorial, where it is placed at the base of a prominent band of bioclastic packstones traditionally placed at the top of the Whitewater Formation; these bioclastic packstones sharply overlie wavy/nodular limestones of the Whitewater Formation. The sharp upper contact of the bioclastic packstones with variegated mudstones of the Preachersville Formation is interpreted as the maximum flooding surface. The Preachersville Formation is overlain here by the Belfast Bed of the Brassfield Formation.

Throughout most of Kentucky, the C5 sequence is absent, and the Silurian Brassfield Formation overlies C4 sequence rocks. The thinning of the C5 sequence into Kentucky is probably partly the result of depositional thinning to the south and partly the result of pre-Silurian erosion. The basal contact of the Brassfield Formation characteristically displays several tens of centimeters of relief at the scale of an outcrop. In south-central Kentucky, C4 and some C3 rocks were removed by pre-Devonian erosion such that the Devonian Boyle Limestone, New Albany Shale, and Chattanooga Shale lie atop Ordovician rocks as old as the C3 sequence.

COMMENTS ON CINCINNATIAN LITHOSTRATIGRAPHIC NOMENCLATURE

TRADITIONAL STRATIGRAPHIC NOMENCLATURE

The history of Cincinnati stratigraphic nomenclature is lengthy and has been summarized elsewhere (Cumings, 1922; Gutstadt, 1958; Weiss and Norman, 1960; Tobin, 1986). A brief summary is necessary as background to the current nomenclatural confusion. The earliest workers proposed only a few large and relatively undifferentiated units for the entire Cincinnati Series (Orton, 1873). Subsequent workers in the late 1800's and early 1900's developed a finely subdivided nomenclature consisting of many formations and members defined on both lithologic and paleontologic criteria. This classification was summarized by Bucher and others

(1939) and later published by Caster and others (1955).

The consistent expression of sequences in the type area of Cincinnati nomenclature was largely responsible for the "layer-cake" approach of early Cincinnati stratigraphers. However, problems arose when these units were traced out of the type area. One common nomenclatural problem arose several times as a worker would attempt to trace a formation across an unconformity—in some cases a sequence boundary, in others, a maximum flooding surface.

As facies at the bases of sequences pinch out into upramp areas of central Kentucky, these same formational names were commonly applied to dissimilar facies that occur at the top of the underlying sequence. For example, the Waynesville Formation consists of the shale-dominated facies and contains an abundant and diverse characteristic fauna. This deepest water facies of the C4 sequence pinches out into upramp areas of central Kentucky; in other words, Ordovician divers swimming southward into central Kentucky would have encountered progressively shallower water environments as they swam into upramp areas. Nonetheless, early workers confused the lithostratigraphic and chronostratigraphic nature of formations and attempted to recognize the Waynesville Formation in central Kentucky. They applied the name Waynesville Formation to a series of "unfossiliferous shales" in the same approximate stratigraphic position as the Waynesville farther north (that is, beneath the Liberty Formation). These unfossiliferous shales correspond to the bioturbated and laminated calcareous mudstone facies of the Rowland Formation. This uppermost facies of the C3 sequence pinches out northward into downramp areas; again, at the end of C3 time, a shallower set of environments would have been developed in downramp areas to the north. Early workers thus attempted to extend a formation from highstand facies above a sequence boundary/maximum flooding surface to highstand facies beneath those surfaces.

Similar problems arose with what are recognized here as transgressive facies, which have an especially narrow geographic distribution. The Saluda Formation in its type area is composed of the bioturbated and laminated carbonate mudstone facies. Near its top, it is overlain by the Hitz Bed, a thin unit of wavy-bedded wackestone with a characteristic molluscan fauna. The Saluda Formation is the uppermost facies of the C4 sequence and predictably thins to the north. The Hitz Bed is the lowermost facies of the C5 sequence and thickens predictably to the north. In this case, workers attempted to extend a formation from the highstand facies below a sequence boundary to the transgressive facies lying above the sequence boundary. However, the name Saluda has been applied to the wavy-bedded wackestone facies to the north where the type-Saluda lithology thins to a pinchout. Hence, in this paper, rocks that have been traditionally referred to the Saluda Formation but that are not type-Saluda lithology (that is, they are wavy-bedded wackestone facies) are referred to in this paper as "Saluda." The name Hitz, having priority, should be elevated to member or formation rank and be applied to these wavy-bedded wackestones of the C5 sequence.

A similar problem occurred with the Sunset Formation. In its type area near Sunset, Kentucky, the Sunset Formation is composed of wavy-bedded wackestones that belong to the transgressive systems tract of the C3 sequence. However, Foerste (1912) also applied the name Sunset Formation to the mixed packstone-shale facies that occurs northward in the same approximate stratigraphic position, below the

wavy/nodular limestones of the Oregonia Formation. In this case, early workers attempted to trace a formation from transgressive facies below a maximum flooding surface to highstand facies above the surface. This mixed packstone-shale facies is not the same lithology as that of the Sunset in its type area; thus, this facies is referred to in this paper as “Sunset” to distinguish it from the type Sunset. A new formational name should be proposed for the “Sunset.”

RECENT STRATIGRAPHIC NOMENCLATURE

Over time, these traditional stratigraphic units were recognized increasingly on paleontologic criteria, despite the original definitions of these units. This trend, the advent of the Stratigraphic Code, and problems arising from tracing formations across sequence boundaries fueled a re-analysis of Cincinnati stratigraphy and the proposal of units based solely on lithology (for example, Brown and Lineback, 1966; Peck, 1966; Hatfield, 1968; Lee, 1974). Some of these units were equivalent to traditional units. For example, the Kope Formation was a redefined Eden Group. The Fairview Formation was redefined with the same name, and the Bellevue Member was elevated to formational rank. Many of the newly proposed units, such as the Dillsboro and Bull Fork Formations, were much closer in scale and lithologic variability to the original units of Orton. Many of the traditional units, however, could be redefined in purely lithologic terms (Hay, 1981; Tobin, 1982; Tobin, 1986).

Despite the Stratigraphic Code, the presence of sequences has created nomenclatural problems in the Cincinnati Series in three ways. First, because any given facies may be overlain or underlain by another conformable facies, by a sequence boundary or by a maximum flooding surface, contact relationships of a unit are difficult to define. This is especially true at the formation scale in the geologic mapping of Kentucky. The Tate Formation, for example, rests “on the Calloway Creek Limestone; elsewhere it rests on the lower member of the Grant Lake Limestone. In southern east-central and southern central Kentucky, the Tate is overlain by the Gilbert Member of the Ashlock Formation; elsewhere it is overlain by the Grant Lake” (Weir and others, 1984, p. E43). Similarly, the Grant Lake Limestone can be defined in up to four ways. The base of the Grant Lake Limestone can be placed on the Calloway Creek Limestone, at the top of the Fairview Formation, or at the top of the Gilbert Member. The top of the Grant Lake Limestone can be placed at the base of the Rowland Formation, at the base of the “Sunset Formation,” or at the base of the Corryville Formation. Contacts defined in multiple ways may therefore be isochronous where the contact coincides with a sequence boundary or maximum flooding surface or diachronous where it coincides with a facies contact.

Second, because these sequences were developed on a ramp, the local succession of facies changes geographically, and separate systems of nomenclature have arisen in different areas, characteristically defined by state lines (fig. 18-5). Ohio, Indiana, and Kentucky currently have different systems of lithostratigraphic nomenclature, Indiana particularly so (Shaver and others, 1970; Weir and others, 1984; Schumacher and others, 1991). The different systems confound stratigraphic relationships by imposing state-line boundaries across a completely gradational system of facies. Even within Kentucky, different facies successions in northern and central Kentucky have led to “vertical

cutoffs” between formations that represent the same facies, such as the Clays Ferry and Fairview Formations (Weir and others, 1984). A more productive way to revise the nomenclature would be to treat the entire outcrop area as a single, genetically related body of strata, rather than to impose arbitrary political lines on a natural system. It should be noted that, wherever possible, the Ohio Division of Geological Survey has tried to extend units recognized in Kentucky into Ohio.

Third, the intertonguing of units and the thinning of units near their pinchouts promoted the lumping of facies into large homogeneous formations such as the Dillsboro and Bull Fork (fig. 18-6). These units straddle sequence boundaries and contain many dissimilar facies, making both environmental and chronostratigraphic interpretations




FIGURE 18-5.—Current stratigraphic terminology, particularly in Ohio and Indiana, for Upper Ordovician rocks in the Cincinnati Arch region. Most of these units refer to a single facies as recognized by Hay (1981), Tobin (1982), and Holland (1990). However, many of these facies are known by multiple stratigraphic names, the boundaries of which are placed at arbitrary vertical cutoffs. For example, The mixed packstone-shale facies of the C1 sequence is known as the Fairview Formation in Ohio and northernmost Kentucky, but as the Clays Ferry Formation in central Kentucky. Names in italics represent a name that should be abandoned on the basis of priority. “Sunset” and “Saluda” refer to stratigraphic units with lithologies unlike the original definitions of Sunset and Saluda. Although not shown on this figure, the Straight Creek Member of the Grant Lake Formation would occupy the same position as the Mt. Auburn Formation, and is regarded here as an equivalent, somewhat more calcareous variation of the Mt. Auburn Formation, similar to the distinction between Grant Lake Limestone and Grant Lake Formation (see Schumacher and others, 1991). UW = Upper Whitewater. Ma = million years.

difficult. In addition, the inconsistent boundary definitions of units such as the Grant Lake Limestone suggest many uses of the same name.

As stated previously, sequence stratigraphy is not a form of lithostratigraphy; its goal is the genetic interpretation of strata, not the recognition of mappable units. However, an understanding of sequence architecture can suggest guidelines for the naming of lithostratigraphic units. On one hand, formational contacts that straddle sequence boundaries or maximum flooding surfaces are characteristically sharp; however, because such a contact is a disconformity, one formation may overlie any of several other formations. In this case, contact definitions should emphasize the sharpness of the contact and the names of all the stratigraphic units that a formation might contact at the disconformity. On the other hand, formational contacts within systems tracts are characteristically gradational; however, the upper formation at a contact will always overlie the same lower formation. In this case, contact definitions should state the gradational yet predictable relationship between two formations.

These concepts suggest several revisions of Cincinnati stratigraphic nomenclature (fig. 18-7):

(1) Each facies within each sequence should be named separately and given formational rank for consistency. This approach has already been done for many of the traditional units (Tobin, 1986; Schumacher and others, 1991). For the sake of consistency, the scale and variability of individual

facies should be kept uniform throughout the section; in other words, if a facies can be recognized in one portion of the section, it should be similarly recognized as a distinct unit wherever it occurs in the section. As a counterexample, the shale-dominated facies of the Kope Formation is recognized as a formation by the Indiana Geological Survey, but the shale-dominated facies of the Waynesville Formation is not recognized as a distinct unit and is included within the lithologically variable Dillsboro Formation. This lack of consistency confuses geologic interpretations because it implies that the two formations represent equivalent units with which to interpret geologic history.

(2) Some units erected during the Kentucky Mapping Project should be abandoned on the basis of priority. The same facies within the same sequence was previously named (fig. 18-5). For example, the Clays Ferry Formation and the Fairview Formation refer to the same lithofacies within the same sequence, and the line between the two is drawn at an arbitrary vertical cutoff across which there is no lithologic contrast. Of these two, the Fairview Formation has priority. Similarly, both the Reba Formation and the Oregonia Formation refer to the wavy/nodular limestone facies of




FIGURE 18-6.—Upper Ordovician stratigraphic units that lump dissimilar facies and straddle disconformities, plotted on background of figures 18-5 and 18-7. Because these units offer little interpretive value, and commonly have ill-defined or variably defined contacts, it is recommended here that their usage be abandoned. Ma = million years.




FIGURE 18-7.—Proposed stratigraphic nomenclature for the Upper Ordovician rocks of the Cincinnati Arch region. Note that each facies within each sequence (that is, a parvafacies) is given a single name; for example, contrast the current usage of Fairview and Clays Ferry Formations for a single facies with the proposed name of Fairview Formation for this facies. In this and other cases, the older name is given priority. Other similar pairs include: Bellevue and Calloway Creek, Mt. Auburn and Gilbert, Oregonia and Reba, Lower Whitewater and Bardstown, Saluda and Preachersville, and Hitz and Saluda. Also note that the “Sunset” and Upper Whitewater on figure 18-5 are labeled here as “tbn,” indicating that new stratigraphic names will be proposed for these units.

the C3 sequence, and their contact is an arbitrary vertical cutoff. Although the Oregonia Formation was defined both lithologically and paleontologically for much of its history, the name is long standing and has been redefined in purely lithologic terms (Tobin, 1986). The Oregonia Formation should be retained in place of the Reba Formation on the basis of priority.

(3) Some previously named units offer little interpretive value because they lump dissimilar facies and straddle sequence boundaries (fig. 18-6). With the exception of a redefined Whitewater Formation (which should be restricted to what has been traditionally called the Lower Whitewater), these units should be abandoned. For example, the Dillsboro Formation straddles three sequence boundaries and includes at least three distinct facies. Although the Dillsboro may be a recognizable mappable unit, that should not be the only goal. Lithostratigraphic units should have some interpretive value as well.

These guidelines would create a much-simplified view of Cincinnati stratigraphy, with a return to many of the traditional names that offer real interpretive power. These guidelines also leave several units unnamed at the present, particularly within the C5 sequence (fig. 18-7). I am planning to propose names for these units following some additional field checking to establish the geographic and stratigraphic limits of these units.

It could be argued that the Stratigraphic Code insists that interpretations should not affect nomenclature and that these guidelines are therefore invalid, but this argument can be countered several ways. The Code's insistence on the lack of interpretation in the naming of a unit is a logical impossibility. The formulation of a stratigraphic code requires an implicit concept of the structure of the stratigraphic record, despite the fact that these concepts were never stated; thus, the Code itself implies a stratigraphic interpretation. Second, the Code states that lithostratigraphic units should not span known disconformities. The identification of a particular surface as a disconformity is a stratigraphic interpretation, not an observation; again, the Code is internally inconsistent because it argues that interpretation should not influence nomenclature. Finally, although sequence stratigraphy can lead to a set of lithostratigraphic guidelines, these guidelines do not necessarily imply a sequence structure to the stratigraphic record.

SUMMARY

The Cincinnati Series consists of five stratigraphic sequences. In the type area of the Cincinnati Series in northernmost Kentucky, southwestern Ohio, and southeastern Indiana, the sequences are dominantly shoaling upward and contain very thin transgressive intervals at their base. In upramp areas of central Kentucky, these same sequences are thinner and contain proportionally greater thicknesses of transgressive deposits. Sequence analysis of the Cincinnati Series offers several guidelines for clarifying the current confusing lithostratigraphic nomenclature.

LOCALITIES

For more information about the individual localities that bear locality codes, please, see Appendix A in this volume.

Aiken High School (OH-HA-0012). Exposures on north

side of driveway to Aiken High School on Belmont Avenue in Cincinnati, Ohio.

Ashlock Cemetery. Road cuts on U.S. Route 27, 0.8 mile (1.3 km) south of the Garrard-Lincoln County line. Lancaster, Kentucky, 7.5-minute quadrangle. Lincoln County, Kentucky. 37°34'25"N latitude, 84°36'49"W longitude.

Beckjord (OH-CT-0019). Road cuts on north side of east-bound U.S. Route 52, just west of Beckjord Coal Plant, 0.6 mile (1 km) east of Ohio Route 749.

Bedford. Road cuts on U.S. Route 42, 2.5 miles (4 km) east of intersection with U.S. Route 421 in Bedford, Kentucky. Bedford, Kentucky, 7.5-minute quadrangle. Trimble County, Kentucky. 38°36'08"N latitude, 85°16'25"W longitude.

Blue Rock Road (OH-HA-0005). Road cuts on Blue Rock Road, 0.1 mile (0.2 km) north of intersection with Cross County Highway and 0.3 mile (0.5 km) southeast of intersection with I-275.

Bon Well Hill (IN-FR-0001). Road cut at intersection of Indiana State Route 101 and Brookville Dam Road, 1 mile (1.6 km) north of Brookville, Indiana. Whitcomb (formerly Mount Carmel), Indiana, 7.5-minute quadrangle.

Brookville North. see Garr Hill.

Bybee. Road cuts on Kentucky Route 52, just east of Bybee, Kentucky. Panola, Kentucky, 7.5-minute quadrangle. Madison County, Kentucky. 37°43'13"N latitude, 84°06'43"W longitude.

Caesar Creek dam (= Caesar Creek gorge) (OH-WA-0002). Hillside outcrops on north side of tailwater area at Caesar Creek dam.

Caesar Creek spillway (OH-WA-0001). Exposures in emergency spillway of Caesar Creek Lake on Clarksville Road.

Chicken Hollow-2 (OH-BR-0001). Road cuts on south side of Chicken Hollow Road, 0.1 mile (0.2 km) east of intersection with U.S. Routes 62 and 68.

Clays Ferry. Road cuts on south side of Kentucky Route 2328 underneath south end of I-75 bridge over Kentucky River. Ford, Kentucky, 7.5-minute quadrangle. Madison County, Kentucky. 37°46'28"N latitude, 84°16'28"W longitude.

Cowan Lake (OH-CN-0001). Exposures in spillway of Cowan Lake and on north bank of Cowan Lake adjacent to spillway.

Flemingsburg. Road cuts along Kentucky Route 32, 1.8 miles (1.1 km) south of Kentucky Route 11. Flemingsburg, Kentucky, 7.5-minute quadrangle. Fleming County, Kentucky. Approximately 38°24'N latitude, 83°42'W longitude.

Frederickstown. Road cut on south side of U.S. Route 150, 0.5 mile (0.8 km) west of Nelson/Washington County line. Maud, Kentucky, 7.5-minute quadrangle. Nelson County, Kentucky. 37°46'09"N latitude, 85°21'25"W longitude.

Garr Hill (= Brookville North) (IN-FR-0003). Road cuts on opposite sides of valley along Indiana Route 101, 5.0 miles (8 km) north of Brookville, Indiana.

Georgetown (OH-BR-0006). Road cuts along Ohio Route 125 on both sides of valley of Whiteoak Creek, 0.8 mile (1.3 km) west of intersection with U.S. Route 68 in Georgetown, Ohio.

Halls Gap North. Road cuts on U.S. Route 27, 3.3 miles (5.3 km) north of intersection with Kentucky Route 643 in Halls Gap, Kentucky. Halls Gap, Kentucky, 7.5-minute quadrangle. Lincoln County, Kentucky. 37°29'52"N latitude, 84°39'00"W longitude.

Lincoln County Line. Road cuts on U.S. Route 27, 0.5 mile (0.8 km) south of Garrard-Lincoln County line and 3.5 miles

(5.6 km) south of Lancaster, Kentucky. Lancaster, Kentucky, 7.5-minute quadrangle. Lincoln County, Kentucky. 37°34'40"N latitude, 84°36'36"W longitude.

Madison #1 (= Madison South road cut) (IN-JE-0003). Road cut on east side of U.S. Route 421, 3.0 miles (4.7 km) south of intersection with Indiana Route 62 north of Madison, Indiana.

Madison #3 (= Madison North road cut) (IN-JE-0001). Road cut on east side of U.S. Route 421, 0.2-0.7 mile (0.3-1.1 km) south of intersection with Indiana Route 62 north of Madison, Indiana.

Maysville (= Maysville-Kentucky Route 11) (KY-MS-0002). Long road cut along Kentucky Route 11, 1.7 miles (2.7 km) north of intersection with Kentucky Routes 546 and 1448, just south of Maysville, Kentucky.

Muddy Creek (OH-HA-0020). Stream exposures on Muddy Creek at Muddy Creek Road bridge, 0.3 mile (0.5 km) west of Westbourne Road.

Orphanage Road (KY-KE-0003). Road cut on north side of Kentucky Route 371 (Orphanage Road), 0.4 mile (0.7 km) west of intersection with Kentucky Route 17.

Point Pleasant. Stream exposures on north side of U.S. Route 52, 0.3 mile (0.5 km) west of western city limits of Point Pleasant, Ohio. Laurel, Ohio-Kentucky, 7.5-minute quadrangle. Clermont County, Ohio. 38°54'02"N latitude, 84°14'13"W longitude.

Richmond (= Richmond-U.S. Route 27) (IN-WY-0001). Road cuts on U.S. Route 27, 0.8 mile (1.3 km) north of Farlow Road and 1.0 mile (1.6 km) south of Richmond, Indiana.

Riedlin Road/Mason Road (KY-KE-0001). Road cuts on Riedlin (Mason) Road at intersection with Kentucky Route 16, 0.4 mile (0.6 km) north of I-275.

South Gate Hill (IN-FR-0005). Road cuts on Indiana Route 1, 1.9 miles (3.0 km) south of its intersection with U.S. Route 52 at Cedar Grove, Indiana.

West Union #3 (= West Union) (OH-AD-0004).

Wright Brothers Memorial. Railroad cut on north border of Wright Brothers Memorial Park. Fairborn, Ohio, 7.5-minute quadrangle. Greene County, Ohio. 39°47'46"N latitude, 84°05'13"W longitude.

REFERENCES CITED

- Aigner, Thomas, 1985, Storm depositional systems: Berlin, Springer-Verlag, 174 p.
- Anstey, R. L., and Fowler, M. L., 1969, Lithostratigraphy and depositional environment of the Eden Shale (Ordovician) in the tri-state area of Indiana, Kentucky, and Ohio: *Journal of Geology*, v. 77, p. 668-682.
- Brown, G. D., Jr., and Lineback, J. A., 1966, Lithostratigraphy of the Cincinnati Series (Upper Ordovician) in southeastern Indiana: *American Association of Petroleum Geologists Bulletin*, v. 50, p. 1018-1023.
- Bucher, W. H., Caster, K. E., and Jones, Stewart, 1939, Elementary description of Cincinnati fossils and strata and plates of commoner fossils in the vicinity of Cincinnati, Ohio: University of Cincinnati, 13 p.
- Caster, K. E., Dalvé, E. A., and Pope, J. K., 1955, Elementary guide to the fossils and strata of the Ordovician in the vicinity of Cincinnati, Ohio: Cincinnati Museum of Natural History, 47 p.
- Cummings, E. R., 1922, Nomenclature and description of the geological formations of Indiana, in Logan, W. N., and others, *Handbook of Indiana geology*: Indiana Department of Conservation Publication 21, p. 403-570.
- Duke, W. L., 1990, Geostrophic circulation or shallow marine turbidity currents? The dilemma of paleoflow patterns in storm-influenced prograding shoreline systems: *Journal of Sedimentary Petrology*, v. 60, p. 870-883.
- Foerste, A. F., 1912, The Arnheim Formation within the areas traversed by the Cincinnati Geanticline: *Ohio Naturalist*, v. 12, p. 429-456.
- Gutstadt, A. M., 1958, Upper Ordovician stratigraphy in eastern interior region: *American Association of Petroleum Geologists Bulletin*, v. 42, p. 513-547.
- Haq, B. U., Hardenbol, J., and Vail, P. R., 1987, Chronology of fluctuating sea levels since the Triassic: *Science*, v. 235, p. 1156-1167.
- Hatfield, C. B., 1968, Stratigraphy and paleoecology of the Saluda Formation (Cincinnati) in Indiana, Ohio, and Kentucky: *Geological Society of America Special Paper* 95, 34 p.
- Hay, H. B., 1981, Lithofacies and formations of the Cincinnati Series (Upper Ordovician), southeastern Indiana and southwestern Ohio: Ph.D. dissertation (unpub.), Miami University, 236 p.
- Hofmann, H. J., 1966, Ordovician paleocurrents near Cincinnati, Ohio: *Journal of Geology*, v. 74, p. 868-890.
- Holland, S. M., 1990, Distinguishing eustasy and tectonics in foreland basin sequences: the Upper Ordovician of the Cincinnati Arch and Appalachian Basin: Ph.D. dissertation (unpub.), University of Chicago, 390 p.
- _____, 1993, Sequence stratigraphy of a carbonate-clastic ramp: the Cincinnati Series (Upper Ordovician) in its type area: *Geological Society of America Bulletin*, v. 105, p. 306-322.
- Ingram, R. L., 1954, Terminology for the thickness of stratification and parting units in sedimentary rocks: *Geological Society of America Bulletin*, v. 65, p. 937-938.
- Jennette, D. C., 1986, Storm-dominated cyclic ramp deposits of the Kope-Fairview transition (Upper Ordovician), southwestern Ohio and northern Kentucky: M.S. thesis (unpub.), University of Cincinnati, 210 p.
- Lee, G. B., 1974, Lithostratigraphy of the Cincinnati Series (Upper Ordovician) from Maysville, Kentucky to Dayton, Ohio: M.S. thesis (unpub.), Miami University, 127 p.
- Meyer, D. L., Tobin, R. C., Pryor, W. A., Harrison, W. B., Osgood, R. G., Hinterlong, G. D., Krumpolz, B. J., and Mahan, T. K., 1981, Stratigraphy, sedimentology, and paleoecology of the Cincinnati Series (Upper Ordovician) in the vicinity of Cincinnati, Ohio, in Roberts, T. G., ed., *GSA Cincinnati '81 Field Trip Guidebooks*: American Geological Institute, v. 1, p. 31-72.
- Mitchum, R. M., 1977, Seismic stratigraphy and global changes of sea level, part 1: glossary of terms used in seismic stratigraphy: *American Association of Petroleum Geologists Memoir*, v. 26, p. 205-212.
- North American Commission on Stratigraphic Nomenclature, 1983, *North American Stratigraphic Code*: American Association of Petroleum Geologists Bulletin, v. 67, p. 841-875.
- Orton, Edward, 1873, Report on the Third Geological District; geology of the Cincinnati Group; Hamilton, Clermont, Warren and Butler Counties: Ohio Division of Geological Survey, v. 1, pt. 1, Geology, p. 365-480.
- _____, 1878, Report on the geology of Warren County; Butler County; Preble County; Madison County: Ohio Division of Geological Survey, v. 3, p. 381-428.
- Parkinson, R. W., and Meeder, J. F., 1989, Mudbank destruction and the formation of a transgressive sand sheet, westernmost Florida Bay, Florida (abs.): *Geological Society of America Abstracts with Programs*, v. 21, no. 6, p. A290.
- Peck, J. H., 1966, Upper Ordovician formations in the Maysville area, Kentucky: *U.S. Geological Survey Bulletin* 1244-B, 30 p.
- Schumacher, G. A., Swinford, E. M., and Shrake, D. L., 1991, Lithostratigraphy of the Grant Lake Limestone and Grant Lake Formation (Upper Ordovician) in southwestern Ohio: *Ohio Journal of Science*, v. 91, p. 56-68.
- Shaver, R. H., Burger, A. M., Gates, G. R., Gray, H. H., Hutchison, H. C., Keller, S. J., Patton, J. B., Rexroad, C. B., Smith, N. M., Wayne, W. J., and Wier, C. E., 1970, *Compendium of rock-unit stratigraphy in Indiana*: Indiana Geological Survey Bulletin 43, 229 p.
- Swadley, W. C., 1979, The Marble Hill Bed. an offshore bar-tidal

- channel complex in the Upper Ordovician Drakes Formation of Kentucky: U.S. Geological Survey Professional Paper 1126-D, 8 p.
- Tobin, R. C., 1982, A model for cyclic deposition in the Cincinnati Series of southwestern Ohio, northern Kentucky, and southeastern Indiana: Ph.D. dissertation (unpub.), University of Cincinnati, 483 p.
- Tobin, R. C., 1986, An assessment of the lithostratigraphic and interpretive value of the traditional "biostratigraphy" of the type Upper Ordovician of North America: *American Journal of Science*, v. 286, p. 673-701.
- Van Wagoner, J. C., Posamentier, H. W., Mitchum, R. M., Vail, P. R., Sarg, J. F., Loutit, T. S., and Hardenbol, J., 1988, An overview of the fundamentals of sequence stratigraphy and key definitions: *Society of Economic Paleontologists and Mineralogists Special Publication 42*, p. 39-45.
- Weir, G. W., Peterson, W. L., Swadley, W. C., and Pojeta, John, Jr., 1984, Lithostratigraphy of Upper Ordovician strata exposed in Kentucky; with a section on biostratigraphy: U.S. Geological Survey Professional Paper 1151-E, 121 p.
- Weiss, M. P., and Norman, C. E., 1960, The American Upper Ordovician Standard. II. Development of stratigraphic classification of Ordovician rocks in the Cincinnati region: *Ohio Division of Geological Survey Information Circular 26*, 14 p.
- Wilgus, C. K., Hastings, B. S., Posamentier, H. W., Van Wagoner, J. C., Ross, C. A., and Kendall, C. G. S. C., ed., 1988, Sea level changes: an integrated approach: *Society of Economic Paleontologists and Mineralogists*, 381 p.

Address of author: Steven M. Holland, Geology Department, University of Georgia, Athens, GA 30602.

APPENDIX A.—TYPE-CINCINNATIAN LOCALITIES

by

Richard Arnold Davis, Sharon C. St. Louis Diekmeyer, Lawrence I. Goldman,
Benjamin F. Dattilo, Steven M. Holland, and Roger J. Cuffey

EXPLANATION

The following list is in two parts. The first part is arranged by state and by county within each state. Within each county the localities are arranged by a code number in the format OH-HA-0002, beginning with the U.S. Postal Service code for the state. The next two letters (or, in a few cases, one letter) are the county code; it is taken from the standard county designations used nationwide in the archaeological community. The four digits refer to the specific locality within the county.

The second part is an alphabetical list by the name of the locality. In cases in which a given locality has more than one designation, the known alternatives all are listed.

Measurements are given in the format "1.0 mile (1.6 km)." The first value is in the units in which the measurement was made or published; the number in parentheses is the value converted to the other system of measurement. In this case, the actual measurement was 1.0 mile, which converts to 1.6 km. If the measurement had been stated as 1.6 km (1.0 mile), the actual measurement was 1.6 km, which converts to 1.0 mile. In the case of map distances in miles, values are

given to the nearest 1/10 mile, because that is the precision of standard automobile odometers.

Some of the original surveying in this particular part of the country was done at an early enough date that the United States Land Survey System has not been applied consistently. Thus, many townships are not the 36-square-mile rectangles that they are elsewhere. Moreover, many do not bear the standard township and range designation, but, instead, have been named. What's worse, in some places there are no real townships at all. Add to that the fact that, even in the areas in which sections have been surveyed, they may not be numbered in the usual boustrophedon (alternating left to right, right to left) manner. In short, folks who are used to the regularity of locality designations west of Ohio are liable to be frustrated by the lack of such regularity in this part of the country.

The OGS numbers in the Ohio portion of this appendix refer to descriptions of measured sections on open file at the Ohio Division of Geological Survey, Columbus, Ohio.

This appendix does not have a bibliography of its own. Full references to works cited herein are listed in Appendix B of this volume, the Bibliography on the type-Cincinnati.

LOCALITIES BY STATE AND COUNTY

INDIANA

Indiana, Dearborn County

IN-D-0001 AURORA

Road cut on U.S. Route 50 about 1 mile (1.6 km) SW of Aurora; SE $\frac{1}{4}$ NE $\frac{1}{4}$ NW $\frac{1}{4}$ sec. 6, T. 4 N., R. 1 W., Aurora, Indiana, 7.5-minute quadrangle; Dearborn County, Indiana

Units.—Bellevue, Fairview, Kope

References.—Anstey and Perry (1973), Hay (1977), Diekmeyer (paper 3 in this volume), Hay (paper 17 in this volume)

IN-D-0002 ST. LEON

Road cuts on north and south sides of I-74, $\frac{1}{4}$ to 1 mile east of St. Leon-Lawrenceburg exit (intersection with Indiana Route 1); sec. 13, T. 7 N., R. 2 W., Cedar Grove, Indiana, 7.5-minute quadrangle; Dearborn County, Indiana; elevation at base: 585 ft (178.3 meters)

Units.—Liberty, Waynesville

References.—Reinhart (1977), Frey (1987b, loc. 6)

IN-D-0003 TANNERS CREEK

East end of first NYC railroad cut west of Guilford; base at ball of inside rail; center S $\frac{1}{2}$ sec. 19, T. 6 N., R. 1 W., Guilford, Indiana, 7.5-minute quadrangle; Miller Twp., Dearborn County, Indiana; 39°09'55"N, 84°54'50"W; elevation at base: 545 ft (166.1 meters)

Units.—Fairview (41 ft/12.5 meters), Kope (22 ft/6.7 meters)

References.—Cumings and Galloway (1913), Weiss, Edwards, Norman, and Sharp (1965), Diekmeyer (paper 3 in this volume)

IN-D-0004 WEST HARRISON

Road cut on Pin Hook Road adjacent to I-74, 3 miles (4.8 km) west of West Harrison; NE $\frac{1}{4}$ NE $\frac{1}{4}$ NW $\frac{1}{4}$ sec. 16, T. 7 N., R. 1 W., Harrison, Ohio-Indiana, 7.5-minute quadrangle; Harrison Twp., Dearborn County, Indiana; 39°15'40"N, 84°51'50"W

Units.—Kope

References.—Anstey and Perry (1973), Diekmeyer (paper 3 in this volume)

IN-D-0005

Road cut on north side of I-74, 4 km (2.5 miles) west of Harrison; NE $\frac{1}{4}$ NW $\frac{1}{4}$ sec. 18, T. 2 N., R. 1 E., Harrison, Ohio-Indiana, 7.5-minute quadrangle; Dearborn County, Indiana

Units.—Mt. Hope Member of Fairview Formation

References.—Giuseffi (1982, loc. 3), Frey (1987a, loc. B)

Indiana, Franklin County

IN-FR-0001 BON WELL HILL

Road cut on Indiana Route 101, 1.3-1.7 miles (2.1-2.7 km) northeast of junction with U.S. Route 52 inside north edge of Brookville; S $\frac{1}{2}$ SW $\frac{1}{4}$ SE $\frac{1}{4}$ sec. 16, T. 9 N., R. 2 W., Whitcomb, Indiana, 7.5-minute quadrangle (formerly Mount Carmel, Indiana, quadrangle); Franklin County, Indiana; 39°26'15"N, 84°59'25"W; 4367024 m N, 672966 m E, UTM zone 16; elevation at base: 747 ft (227.7 meters); elevation at top: 837 ft (255.1 meters)

Units.—"Brookville Fm." (Liberty, Waynesville, and "Excello" Mbrs.); in terms of the "traditional nomenclature": Liberty, Waynesville, Arnheim

References.—Hay (1977), Hay, Pope, and Frey (1981); Brandt Velbel (1985, loc. A), Frey (1987a, loc. D; 1987b, loc. 4), Hay and Cuffey (paper 10 in this volume), Hay (paper 17 in this volume), Holland (paper 18 in this volume)

IN-FR-0002 BROOKVILLE DAM SPILLWAY

Spillway below Brookville Dam, north of Brookville; SW $\frac{1}{4}$ SE $\frac{1}{4}$ SE $\frac{1}{4}$ sec. 17 and NW $\frac{1}{4}$ NE $\frac{1}{4}$ NE $\frac{1}{4}$ sec. 20, T. 9 N., R. 2 W., Brookville, Indiana, 7.5-minute quadrangle; Franklin County, Indiana; 39°26'13"N, 85°00'19"W; 4366972 m N, 671679 m E, UTM zone 16; elevation of base of spillway: 623 ft (189.9 meters) at river level in valley bottom out in front of dam

Units.—Waynesville, "Excello mbr." of "Brookville fm.," "Sta-

tion Hollow mbr.” of “Brookville fm.” Bellevue, Miamitown; in terms of the “traditional nomenclature”: Waynesville, Arnheim, Mt. Auburn, Corryville, Bellevue, Miamitown; Fairview and higher strata exposed in highwall opposite measured section

Note.—Across spillway from measured section is a highwall exposing 60 meters (200 ft) of Ordovician rock that correlates with strata exposed at Garr Hill (IN-FR-0003) and upper part of Bon Well Hill (IN-FR-0001). Height and steepness of this highwall make access very difficult; in any case, access is not normally permitted

References.—Hay (1977), Hay (1981), Hay, Pope, and Frey (1981), Tobin (1982), Diekmeyer (paper 3 in this volume), Hay (paper 17 in this volume), Hay and Cuffey (paper 8 in this volume)

IN-FR-0003 GARR HILL

Alternate designation.—Brookville North

Road cut on both sides of Indiana Route 101, 6.0-6.4 miles (9.6-10.2 km) northeast from junction of U.S. Route 52 and Indiana Route 101 within north edge of Brookville; E¹/₂SE¹/₄ NE¹/₄ and E¹/₂NE¹/₄SE¹/₄ sec. 35, T. 10 N., R. 2 W., Whitcomb, Indiana, 7.5-minute quadrangle; Franklin County, Indiana; 39°29'11"-39°29'29"N, 84°56'58"W; 4372556-4373108 m N, 676410 m E, UTM zone 16; elevation: bottom, 900 ft (274.3 meters); top, 990 ft (301.8 meters)

Units.—Basal beds of upper part of upper Whitewater, Saluda, lower Whitewater, Liberty, Liberty Mbr. of “Brookville fm.”

References.—Hay (1977, p. 1-18 to 1-22), Hay, Pope, and Frey (1981, stop 4, p. 80, 84-86), Bliss (1984), Frey (1987a, locs. G and J), Hay and Cuffey (paper 11 in this volume), Hay (paper 17 in this volume), Holland (paper 18 in this volume)

IN-FR-0004 FAIRFIELD CAUSEWAY

Alternate designations.—Causeway Road, Fairfield Road

Cuts along Fairfield Road, which runs west from Indiana Route 101, passes near village of New Fairfield, and traverses causeway between north and south parts of Brookville Reservoir; causeway is in sec. 28, T. 10 N., R. 2 W., New Fairfield, Indiana, 7.5-minute quadrangle; Fairfield Twp., Franklin County, Indiana; road cut at west end of causeway is in sec. 28, but Everton, Indiana, 7.5-minute quadrangle; center of causeway is at 39°30'23"N, 84°59'43"W

Units.—Liberty, Waynesville

IN-FR-0005 SOUTH GATE HILL

Alternate designation.—Indiana Route 1; Cedar Grove

Road cuts on both sides of Indiana Route 1, 1.0-1.5 miles (1.6-2.4 km) north of crossroads at village of South Gate and 1.9-2.4 miles (3.0-3.8 km) south of intersection of Indiana Route 1 with U.S. Route 52 at Cedar Grove; W¹/₂W¹/₂SE¹/₄ sec. 23, T. 8 N., R. 2 W., Cedar Grove, Indiana, 7.5-minute quadrangle; Franklin County, Indiana; 39°20'08"-39°20'30"N, 84°57'20"-84°57'24"W; 4355853-4356469 m N, 676098-676194 m E, UTM zone 16; elevation at base: 750 ft (228.6 meters)

Units.—Saluda, lower Whitewater, “Brookville formation” (including Liberty Member, Waynesville Shale Member, and “Excello member”); in terms of “traditional nomenclature”: Saluda, lower Whitewater, Liberty, Waynesville, Oregonia Mbr. of Arnheim Fm.

References.—Kirchner (1991), Hay, Kirchner, and Cuffey (paper 12 in this volume), Hay (paper 17 in this volume), Holland (paper 18 in this volume)

IN-FR-0006

Small stream-bank exposures along Whistle Creek where Whistle Creek Road crosses stream, north side of road; center sec. 4, T. 11 N., R. 12 E., Metamora, Indiana, 7.5-minute quadrangle; Franklin County, Indiana

Units.—“trilobite shale unit,” Waynesville

References.—Frey (1987b, loc. 5)

Indiana, Jefferson County

IN-JE-0001 MADISON NORTH ROAD CUT

Alternate designation.—Madison #3 (Holland)

Road cut along U.S. Route 421 0.2-0.7 mile (0.3-1.1 km) south of intersection with Indiana Route 62; massive Saluda Dolomite is nicely exposed 0.5 mile (0.8 km) south of intersection; SW corner sec. 13 and NW¹/₄NW¹/₄ and center NW¹/₄ sec. 24, T. 4 N., R. 10 E., Canaan, Indiana, 7.5-minute quadrangle; Jefferson County, Indiana; 38°46'45"N, 85°21'58"W (Saluda exposure); 4293308 m N, 641928 m E, UTM zone 16 (Saluda exposure)

Units.—Silurian, Whitewater, Saluda, Liberty

References.—Totten and Hay (1987, North Road Cut, A in Fig. 1), Hay, Totten, and Cuffey (paper 6 in this volume), Hay (paper 17 in this volume), Holland (paper 18 in this volume)

IN-JE-0002 MADISON MIDDLE ROAD CUT

Road cut along U.S. Route 421 beginning about 1.1-1.2 miles (1.8-1.9 km) south of intersection with Indiana Route 62; center SW¹/₄ sec. 24, T. 4 N., R. 10 E., Canaan, Indiana, 7.5-minute quadrangle; Jefferson County, Indiana

Units.—“Excello member” of “Brookville formation”

References.—Totten and Hay (1987, Middle Road Cut, B in Fig. 1), Hay, Totten, and Cuffey (paper 6 in this volume), Hay (paper 17 in this volume)

IN-JE-0003 MADISON SOUTH ROAD CUT

Alternate designation.—Madison #1 (Holland)

Road cut along U.S. Route 421 beginning about 3.0 miles (4.7 km) south of intersection with Indiana Route 62 and ending about 0.5 mile (0.8 km) north of Madison; center W¹/₂NE¹/₄ sec. 35, T. 4 N., R. 10 E., Madison West, Indiana-Kentucky, 7.5-minute quadrangle; Jefferson County, Indiana; 38°44'58"N, 85°22'35"W

Units.—“Excello mbr.” of “Brookville fm.” Bellevue

References.—Totten and Hay (1987, South Road Cut, C in Fig. 1), Holland (1990), Diekmeyer (paper 3 in this volume), Hay, Totten, and Cuffey (paper 6 in this volume), Hay (paper 17 in this volume); Holland (paper 18 in this volume)

IN-JE-0004 MADISON-INDIANA ROUTE 56

Four closely spaced road cuts along Indiana Route 56 (formerly also Indiana Route 62) 0.3-1.2 miles (0.5-1.9 km) east (downgrade) from junction with present Indiana Route 62 about 4 miles west of downtown Madison; most conspicuous cut is highwall exposing Saluda Dolomite at sharp curve 0.5-0.6 mile (0.8-1.0 km) east of that junction; NE corner sec. 6 and N¹/₂N¹/₂NW¹/₄ sec. 5, T. 3 N., and S¹/₂SW¹/₄SE¹/₄ sec. 31, T. 4 N., R. 10 E., Madison West, Indiana-Kentucky, 7.5-minute quadrangle; Jefferson County, Indiana; 38°44'12"N, 85°26'28"W (high cut through Saluda); 4288480 m N, 635461 m E, UTM zone 16 (high cut through Saluda)

Units.—upper Whitewater, Saluda, Liberty, Waynesville, Arnheim

References.—Hattin, Nosow, Perkins, Stumm, Mound, and Utgaard (1961, stop 8, p. 328-331, 343-347); Hattin and Cuffey (paper 14 in this volume)

IN-JE-0005 MADISON-RILEY CREEK

Low outcrop on west bank of dry creek (Riley Creek, although name is not on topographic quadrangle) along west side of trailer park on northwest side of historic Madison (down on valley floor near Ohio River); center SW¹/₄NW¹/₄SE¹/₄ sec. 34, T. 4 N., R. 10 E., Madison West, Indiana-Kentucky, 7.5-minute quadrangle; Jefferson County, Indiana; 38°44'36"N, 85°23'42"W; 4289290 m N, 639478 m E, UTM zone 16

Units.—Bellevue (Monticulipora molesta bed)

References.—Patton, Perry, and Wayne (1953, stop 12, p. 26); Dively and Cuffey (1994)

IN-JE-0006 MADISON-HANGING ROCK

Road cuts along east side of Indiana Route 7, 0.8-1.2 miles

(1.3-1.9 km) north of junction with Indiana Route 56 on west side of downtown Madison; Saluda Dolomite forms ledges of small waterfall at sharp bend in road 1.0 mile (1.6 km) north of that junction, ascending out of Ohio River valley; NW $\frac{1}{4}$ NE $\frac{1}{4}$ sec. 34, T. 4 N., R. 10 E., Clifty Falls, Indiana, 7.5-minute quadrangle; Jefferson County, Indiana; 38°45'04"N, 85°23'40"W (Saluda waterfall); 4290186 m N, 639510 m E, UTM zone 16

Units.—Silurian, "Upper Whitewater," Saluda, Liberty
References.—Patton, Perry, and Wayne (1953, stop 12, p. 26-27); Butler & Cuffey (1994, 1996)

Indiana, Ohio County

IN-O-001 RISING SUN

Bank of Indiana Route 56, 2 miles (3.2 km) SW of Rising Sun; base in culvert; incomplete exposure; center SE $\frac{1}{4}$ SW $\frac{1}{4}$ sec. 16, T. 3 N., R. 1 W., Aberdeen, Indiana-Kentucky, 7.5-minute quadrangle; Randolph Twp., Ohio County, Indiana; 38°55'20"N, 84°53'00"W; elevation at base: 630 ft (192.0 meters)

Units.—Fairview, Bellevue, Kope (52 ft/15.8 meters to base of Maysvillian)

References.—Weiss, Edwards, Norman, and Sharp (1965), Diekmeyer (paper 3 in this volume)
OGS measured section no. 14809

IN-O-002 ARNOLD CREEK

Stream exposures in tributary of Arnold Creek, 4.7 km (2.9 miles) west of Rising Sun, near intersection of White Road and Indiana Route 262; extending from NE $\frac{1}{4}$ NE $\frac{1}{4}$ SE $\frac{1}{4}$ to NW $\frac{1}{4}$ SE $\frac{1}{4}$ NE $\frac{1}{4}$ sec. 12, T. 3 N., R. 2 W., Aberdeen, Indiana-Kentucky, 7.5-minute quadrangle; Randolph Twp., Ohio County, Indiana; 38°56'15"N, 84°55'55"W

Units.—Kope
References.—Anstey and Perry (1973), Diekmeyer (paper 3 in this volume)

Indiana, Switzerland County

IN-SW-001 WILEY BRANCH

Stream exposures in Wiley Branch about 0.5 km (0.3 mile) east of Florence; NE $\frac{1}{4}$ SE $\frac{1}{4}$ NE $\frac{1}{4}$ sec. 32 to NE $\frac{1}{4}$ NW $\frac{1}{4}$ NW $\frac{1}{4}$ sec. 33, T. 2 N., R. 1 W., Florence, Indiana-Kentucky, 7.5-minute quadrangle; Switzerland County, Indiana; 38°47'53"N, 84°53'34"W

Units.—Kope
References.—Anstey and Perry (1973), Diekmeyer (paper 3 in this volume)

Indiana, Wayne County

IN-WY-001 RICHMOND-U.S. ROUTE 27

Large road cut on U.S. Route 27, 1.2-1.3 miles (1.9-2.1 km) southwest of junction with Indiana Route 227 on south edge of Richmond; NW $\frac{1}{4}$ SE $\frac{1}{4}$ SW $\frac{1}{4}$ sec. 17, T. 13 N., R. 1 W., Richmond, Indiana, 7.5-minute quadrangle; Wayne Twp., Wayne County, Indiana; 39°47'13"N, 84°54'10"W; 4406000 m N, 679610 m E, UTM zone 16

Units.—Whitewater
References.—Hay, 1977; Frey (1987a, loc. H), Holland (paper 18 in this volume), Hay (paper 17 in this volume), Hay and Cuffey (paper 15 in this volume)

IN-WY-002 SHORT CREEK/STRAIGHT LINE PIKE

Small road cut and adjacent stream cut where Straight Line Pike (Road) crosses Short Creek, 1.0 mile (1.6 km) south of the U.S. 27-Indiana Route 227 junction, Richmond; center E edge SE $\frac{1}{4}$ sec. 17, T. 13 N., R. 1 W., Richmond, Indiana, 7.5-minute quadrangle; Wayne Twp., Wayne County, Indiana

Units.—Whitewater
References.—Hay and Cuffey (paper 15 in this volume)

IN-WY-003 SIM HODGIN PARKWAY

Road cut along Sim Hodgin Parkway in Whitewater Gorge through Richmond, Indiana, Wayne County, Indiana
Units.—Whitewater and/or Elkhorn
References.—Hay and Cuffey (paper 15 in this volume)

IN-WY-004 ELKHORN FALLS

Waterfalls just west of Indiana Route 227 at bridge over Elkhorn Creek (just north of intersection of Indiana Route 227 and Niewohner Road); this site is at east end of Elkhorn Creek locality (IN-WY-0008); SW $\frac{1}{4}$ SE $\frac{1}{4}$ NW $\frac{1}{4}$ sec. 22, T. 13 N., R. 1 W., New Paris, Ohio-Indiana, 7.5-minute quadrangle; Boston Twp., Wayne County, Indiana; NOTE: this is private property; the owner is adamant about trespassing!

Units.—Brassfield, Elkhorn (apparently)
References.—Hay and Cuffey (paper 15 in this volume)

IN-WY-005 SUB RUN

Exposure at intersection of Indiana Route 121 and Pottershop/Abington/Liberty Road $\frac{1}{8}$ mile (0.2 km) southeast of Abington; SE $\frac{1}{4}$ sec. 2, T. 12 N., R. 2 W., Liberty, Indiana, 7.5-minute quadrangle; Wayne County, Indiana; NOTE: this is private property; obtain permission before visiting site
Units.—Liberty Member of "Brookville formation," Whitewater Formation

References.—Hay and Cuffey (paper 15 in this volume)

IN-WY-006 THISTLETHWAITE FALLS

Waterfalls on Waterfall Road, west from U.S. Route 27 North (at first traffic light north of hospital); Thistlethwaite Falls is at northern termination of Whitewater Gorge on West Fork of East Fork Whitewater River; NW $\frac{1}{4}$ SW $\frac{1}{4}$ SE $\frac{1}{4}$ sec. 29, T. 14 N., R. 1 W., Richmond, Indiana, 7.5-minute quadrangle; Wayne Twp., Wayne County, Indiana

Units.—Whitewater
References.—Hay and Cuffey (paper 15 in this volume)

IN-WY-007 MIDDLEFORK RESERVOIR

Outcrop at spillway of reservoir, on Reservoir Road (Cart Road) north of Richmond and east of U.S. Route 27 (turn is north of hospital and north of Waterfall Road); NE $\frac{1}{4}$ sec. 28, T. 14 N., R. 1 W., Richmond, Indiana, 7.5-minute quadrangle; Wayne Twp., Wayne County, Indiana; NOTE: property belongs to Indiana-American Water Company

Units.—Whitewater
References.—Hay and Cuffey (paper 15 in this volume)

IN-WY-008 ELKHORN CREEK

Stream cuts and stream bed between Straight Line Road (Pike) and Indiana Route 227; most of section, including Elkhorn Shale, is exposed between Straight Line Road and Fouts Road; east end of this stretch of creek is Elkhorn Falls (IN-WY-0004); S $\frac{1}{2}$ sec. 21, T. 13 N., R. 1 W. and SW $\frac{1}{4}$ sec. 22, T. 13 N., R. 1 W., Richmond, Indiana, and New Paris, Ohio-Indiana, 7.5-minute quadrangles; Boston Twp., Wayne County, Indiana; NOTE: this is private property; get permission beforehand

Units.—Elkhorn, Whitewater
References.—Hay and Cuffey (paper 15 in this volume)

KENTUCKY

Kentucky, Boone County

KY-BE-001 MALL ROAD

Exposure at south end of Mall Road in Florence, in area now occupied by a shopping plaza; Union, Kentucky, 7.5-minute quadrangle; Boone County, Kentucky

Units.—Bull Fork, Corryville
References.—Meyer and others (1981, first of two entries labeled "stop 5," p. 58-62), Meyer (1990, p. 156)

KY-BE-0002 RABBIT HASH

Road cut on Boone County Road 536 beginning at community of Rabbit Hash and extending about 0.3 mile (0.5 km) east; Rising Sun, Kentucky–Indiana, 7.5-minute quadrangle; Boone County, Kentucky; Carter Coordinates: 20-DD-56, 426.72-1,005.84 m east of west line, 944.88 m south of north line; 38°56'40"N, 84°50'50"W

Units.—Kope

References.—Anstey and Perry (1973), Diekmeyer (paper 3 in this volume)

KY-BE-0003 LAWRENCEBURG

Road cuts on either side of I-275 in Kentucky, 0.7 km (0.4 mile) south of Ohio River across from Lawrenceburg, Indiana; Lawrenceburg, Indiana–Kentucky–Ohio, 7.5-minute quadrangle; Boone County, Kentucky; 39°05'46"N, 84°49'04"W; 4329508 N, 688721 E, UTM zone 16

Units.—Corryville (?), Bellevue, Miamitown, Fairview, Kope

References.—Dattilo (paper 7 in this volume)

Kentucky, Bracken County

KY-BK-0001 BRADFORD

Road cut on Kentucky Route 8 west of Little Snag Creek, 0.25 mile (0.40 km) west of Bradford; Moscow, Ohio–Kentucky, 7.5-minute quadrangle; Bracken County, Kentucky; 38°47'10"N, 84°09'00"W; elevation at base: 504 ft (153.6 meters)

Units.—Point Pleasant (63 ft/19.2 meters); base of Kope estimated by Weiss, Edwards, Norman, and Sharp (1965) to be about 15 ft (4.6 meters) above top of section exposed

References.—Weiss, Edwards, Norman, and Sharp (1965), Harrison (1981 *in* Meyer and others, 1981, stop 1, p. 34, 35, 36), Diekmeyer (paper 3 in this volume)

OGS measured section no. 14932

KY-BK-0002 MELDAHL DAM

Road cut on south side of Kentucky Route 8 about 1.8 miles (2.9 km) west of Meldahl Dam; Moscow, Ohio–Kentucky, 7.5-minute quadrangle; Bracken County, Kentucky

Units.—Kope, Point Pleasant

References.—Harrison (1981, *in* Meyer and others, 1981, stop 2, p. 34, 35, 36)

KY-BK-0003 FOSTER

Bank along Kentucky Route 8 about 0.2 mile (0.3 km) north-west of Foster, starting 8.3 ft (2.53 meters) above end of bridge; Moscow, Ohio–Kentucky, 7.5-minute quadrangle; Bracken County, Kentucky; 38°48'15"N, 84°13'15"W; elevation at base: 512 ft (156.1 meters)

Units.—Kope (10 ft/3.0 meters + another 50 ft/15.2 meters poorly exposed above that), Point Pleasant (60 ft/18.3 meters)

References.—Weiss, Edwards, Norman, and Sharp (1965), Diekmeyer (paper 3 in this volume); see also Foerste (1905, p. 151)

OGS measured section no. 14937

Kentucky, Campbell County

KY-CP-0001 FORT THOMAS

Alternate designations.—445/8 road cut; I-275 East/Kentucky Route 445; Brent

Composite section of two groups of exposures: 1—road cut at intersection of Kentucky Route 8 and Kentucky Route 445; 2—on north side of I-275 just west of Ohio River; Newport, Kentucky–Ohio, 7.5-minute quadrangle; Campbell County, Kentucky; 39°03'15"N, 84°26'00"W

Units.—Kope (59 meters/193.6 ft); Jennette (1986): Fairview (7.2 meters/23.6 ft), Kope (21.3 meters/69.9 ft)

References.—Harrison and Mahan (1981, *in* Meyer and others, 1981, stop 3, p. 36-45), Tobin (1982), Hoge (1985), Jennette

(1986), Diekmeyer (paper 3 in this volume)

KY-CP-0002 NEWPORT SHOPPING CENTER

Alternate designation.—“Newport” (Weiss, Edwards, Norman, and Sharp, 1965)

Cut behind Newport Shopping Center on U.S. Route 27 (Monmouth St./Alexandria Pike) at south edge of Newport, northeast of intersection with Carothers Road; Newport, Kentucky–Ohio, 7.5-minute quadrangle; Campbell County, Kentucky; 39°04'50"N, 84°28'45"W; elevation at base: 575 ft (175.3 meters)

Units.—base of Maysvillian, Kope (164 ft/50.0 meters)

References.—Weiss, Edwards, Norman, and Sharp (1965), Diekmeyer (paper 3 in this volume)

KY-CP-0003 SOUTHGATE

Alternate designation.—I-471/U.S. Route 27

Road cuts on both sides of I-471 at intersection with U.S. Route 27 (interchange 2), E of Southgate; Newport, Kentucky–Ohio, 7.5-minute quadrangle; Campbell County, Kentucky; 39°04'00"N, 84°27'50"W

Units.—Fairview (0.5 meter/1.6 ft), Kope (23.2 meters/76.1 ft)

References.—Jennette (1986), Diekmeyer (paper 3 in this volume)

KY-CP-0004 NEWPORT-GRAND AVENUE

Alternate designation.—I-471/Grand Avenue

Outcrops on west side of I-471 at Grand Avenue, Newport (interchange 4 on quad map); Newport, Kentucky–Ohio, 7.5-minute quadrangle; Campbell County, Kentucky; 39°04'45"N, 84°28'05"W

Units.—Fairview (2.9 meters/9.5 ft), Kope (26 meters/85.3 ft)

References.—Jennette (1986), Diekmeyer (paper 3 in this volume)

KY-CP-0005 JOHNS HILL

Alternate designation.—I-275/Kentucky Route 9

Road cuts on both sides of I-275, ¼ mile (0.4 km) east of interchange with Kentucky Route 9, on flank of Johns Hill; Newport, Kentucky–Ohio, 7.5-minute quadrangle; Campbell County, Kentucky; 39°02'35"N, 84°28'40"W

Units.—Fairview (8.0 meters/26.2 ft), Kope (8.7 meters/28.5 ft)

References.—Jennette (1986), Diekmeyer (paper 3 in this volume)

Kentucky, Gallatin County

KY-GA-0001 SUGAR CREEK

Two exposures: 1—road cut on U.S. Route 42 500 ft (152.4 meters) west of bridge over Sugar Creek; 2—stream exposures in gully ascending hillside, immediately south of U.S. Route 42, about 1 mile (1.6 km) west of bridge over Sugar Creek; Patriot, Kentucky–Indiana, 7.5-minute quadrangle; Gallatin County, Kentucky; Carter Coordinates: 1: 71-BB-57, 1400 ft (426.7 m) east of west line, 200 ft (61.0 m) south of north line; 2: 16-BB-57, 1700 ft (518.2 m) east of west line, 1500 ft (457.2 m) south of north line; 38°47'00"N, 84°48'46"W

Units.—Kope, Lexington Limestone

References.—Anstey and Fowler (1969, localities 4 and 7), Anstey and Fowler (1972), Diekmeyer (paper 3 in this volume)

Kentucky, Kenton County

KY-KE-0001 RIEDLIN ROAD/MASON ROAD

Road cuts along both sides of Riedlin Road/Mason Road, at intersection with east of Kentucky Route 16 (Taylor Mill Road); intersection is 0.4 mile (0.6 km) north of I-275 (exit 79); Covington, Kentucky–Ohio, 7.5-minute quadrangle (Luft, 1971; Ford, 1974); Kenton County, Kentucky; 39°01'15"N, 84°30'30"W

Units.—Bellevue (4.5 meters/14.7 ft), Miamitown (1.0 meter/3.3 ft), Fairview (32.6 meters/107 ft), Kope (8.5 meters/27.9 ft)
References.—Tobin and Pryor (1981, *in* Meyer and others, 1981, stop 4), Tobin (1982, 1986), Davis (1986), Jennette (1986), Frey (1987a, loc. C), Holland (1990), Diekmeyer (1990, 1992, paper 3 in this volume), Dattilo (paper 7 in this volume), Holland (paper 18 in this volume)

KY-KE-0002 THOMAS MORE PARKWAY

Series of small cuts behind "Chancellor Center" office-condominium on Thomas More Parkway, 1.0 km (0.6 mile) east of intersection with Turkeyfoot Road; this site, when cut in 1989, contained three broad benches bounded by 2- to 3-meter (6- to 10-ft), west-facing exposures. Covington, Kentucky–Ohio, 7.5-minute quadrangle (Luft, 1971; Ford, 1974); Kenton County, Kentucky; 39°01'20"N, 84°33'57"W
Units.—Corryville, possibly Bellevue
References.—Goldman (paper 9 in this volume)

KY-KE-0003 ORPHANAGE ROAD

Road cut on north side of Kentucky Route 371 (Orphanage Rd.), 0.4 mile (0.6 km) west of intersection with Kentucky Route 17; Covington, Kentucky–Ohio, 7.5-minute quadrangle (Luft, 1971; Ford, 1974); Kenton County, Kentucky; 39°01'47"N, 84°32'30"W
Units.—Kope (30 meters/98.4 ft)
References.—Frey (1987a, loc. A), Holland (1990, paper 18 in this volume), Diekmeyer (paper 3 in this volume)

KY-KE-0004 COVINGTON

Alternate designation.—I-75/Covington
Cuts along I-75/I-71 0.8 mile (1.3 km) north of interchange with Kentucky Route 1072 (Kyles Lane) (NOTE: since 1992, I-75/I-71 has been slightly re-routed in this stretch); Covington, Kentucky–Ohio, 7.5-minute quadrangle (Luft, 1971; Ford, 1974); Kenton County, Kentucky; 39°03'45"N, 84°31'30"W
Units.—Fairview (3 meters/9.8 ft), Kope (30.5 meters/100.1 ft); Jennette (1986); Fairview (0.3 meter/1.0 ft), Kope (23.3 meters/76.4 ft)
References.—Tobin (1982), Jennette (1986), Diekmeyer (paper 3 in this volume)

KY-KE-0005 I-275/TURKEYFOOT ROAD

Road cut on I-275 east at Turkeyfoot Road (Exit 2); Covington, Kentucky–Ohio, 7.5-minute quadrangle (Luft, 1971; Ford, 1974); Kenton County, Kentucky; 39°01'20"N, 84°34'00"W
Units.—Fairview (0.7 meter/2.3 ft), Kope (12.5 meters/41.0 ft)
References.—Jennette (1986), Diekmeyer (paper 3 in this volume)

KY-KE-0006 WAYNE ROAD

Alternate designation.—Devou Park
Hill above picnic area west of Wayne Road on northwest corner of Prisoner's Lake, in Devou Park in Covington; Covington, Kentucky–Ohio, 7.5-minute quadrangle (Luft, 1971; Ford, 1974); Kenton County, Kentucky; 39°05'00"N, 84°32'14"W; 4328712 N, 713048 E, UTM zone 16
Units.—lower Bellevue, Miamitown, upper Fairview
References.—Dattilo (paper 7 in this volume)

Kentucky, Lewis County

KY-LW-0001 CABIN CREEK

Road cut on both sides of Kentucky Route 10 about 2 miles (3.2 km) east of Tollesboro; Tollesboro, Kentucky, 7.5-minute quadrangle; Lewis County, Kentucky; Carter Coordinate section 3-Y-72
Units.—Brassfield, Preachersville Member of Drakes
References.—Potter and others (1991, stop 2C)

Kentucky, Mason County

KY-MS-0002 MAYSVILLE-KENTUCKY ROUTE 11

Alternate designation.—Maysville (Holland)
Composite section of seven road cuts along Kentucky Route 11 for 2 miles (3.2 km) north of intersection with Kentucky Route 546 (AA Highway), just south of Maysville; Mays Lick, Kentucky, 7.5-minute quadrangle; Mason County, Kentucky; approximately 38°37'N, 83°45'W; Carter Coordinate section 20-Z-69
Units.—Grant Lake, Fairview, Kope
References.—Potter and others (1991, stop 10); Holland (paper 18 in this volume)
OGS measured section no. 16914

KY-MS-0003 SOUTH LICK BRANCH

Two road cuts along Kentucky Route 546 (AA Highway) near its intersection with Kentucky Route 1449 and where Route 546 crosses Stone Lick Branch; Orangeburg, Kentucky, 7.5-minute quadrangle; Mason County, Kentucky; Carter Coordinate sections 22-Z-70 and 23-Z-70
Units.—lower half of Bull Fork (equivalent to Arnheim)
References.—Potter and others (1991, stops 1a and 1b)

KY-MS-0004 MAYSVILLE-U.S. ROUTES 62 & 68

Road cuts along U.S. Routes 62 and 68 just south of Maysville, 0.8 mile (1.3 km) from south end of Maysville-Aberdeen Bridge; Maysville West, Ohio–Kentucky, 7.5-minute quadrangle; Mason County, Kentucky; 38°38'25"N, 83°46'00"W; elevation at base: 162 meters (531 ft)
Units.—Bellevue (22.6 meters/74.1 ft), Fairmount Member of Fairview Formation (21.6 meters/70.9 ft), Mt. Hope Member of Fairview Formation (40.8 meters/133.9 ft), Kope; this locality is one of three co-equal type sections of the Kope Formation (other two are OH-BR-0003 and OH-BR-0004)
References.—Weiss and Sweet (1964), Diekmeyer (paper 3 in this volume)

KY-MS-0005 SLEEPY HOLLOW

Road cut on Kentucky Route 1449, about 6.4 km (4.0 miles) north-northwest of Orangeburg; Orangeburg, Kentucky, 7.5-minute quadrangle; Mason County, Kentucky; 38°37'10"N, 83°42'51"W
Units.—Bull Fork, Grant Lake Limestone (including Corryville Member), Fairview. This locality is the type section of Grant Lake Limestone
References.—Peck (1966), Lee (1974), Diekmeyer (paper 3 in this volume), Goldman (paper 9 in this volume)

KY-MS-0006 MAYSVILLE BRYOZOAN REEF MOUNDS

Road cut on west side of U.S. Route 68, 0.1 mile (0.2 km) south of junction of U.S. Route 68 and U.S. Route 62 at south edge of village of Washington; Mays Lick, Kentucky, 7.5-minute quadrangle (Gibbons, 1968); Mason County, Kentucky; 38°36'36"N, 83°48'39"W; 4277049 m N, 255255 m E, UTM zone 17 (Note: road widened and mounds destroyed between 1992 and 1998.)
Units.—upper member of Grant Lake Limestone (Corryville and Mt. Auburn equivalents)
References.—Cuffey (paper 5 in this volume)

OHIO

Ohio, Adams County

OH-AD-0002 WEST FORK OHIO BRUSH CREEK

Road cut on south side of Ohio Route 32 approximately 2 miles (3.2 km) east of intersection of Ohio Routes 32 and 246; Seaman, Ohio, 7.5-minute quadrangle; Scott Twp., Adams County, Ohio

Units.—Brassfield, Preachersville Member of Drakes Formation, Bull Fork

References.—Schumacher, Shrake, Swinford, Brockman, and Wickstrom (1987, "stop 1," p. 39, 46, 47)

OH-AD-0003 MANCHESTER

Exposure on west-facing hill on east side of Ohio Route 136 near intersection of Route 136 and Brown Hill Road, approximately 1.5 miles (2.4 km) north of Manchester; this road cut is within a few hundred meters of the Isaacs Creek exposure of Sweet (1979a, p. G23, loc. 25). Manchester Islands, Kentucky–Ohio, 7.5-minute quadrangle; Sprigg Twp., Adams County, Ohio; 38°44'07"N, 83°03'09"W

Units.—Bull Fork, Corryville

References.—Lee (1974), Goldman (paper 9 in this volume), Hay (paper 17 in this volume)

OH-AD-0004 WEST UNION

Alternate designation.—West Union #3 (Holland, paper 18 in this volume)

Road cut on west side of Ohio Route 41, 3.2 miles (5.1 km) northeast from intersection with Ohio Route 247 in West Union; West Union, Ohio, 7.5-minute quadrangle; Tiffin Twp., Adams County, Ohio; 38°49'38"N, 83°30'20"W

Units.—Silurian, Elkhorn, Whitewater

References.—Holland (paper 18 in this volume)

Ohio, Brown County

OH-BR-0001 CHICKEN HOLLOW

Two localities: 1—road cut west of U.S. Routes 68 and 62 just north of intersection with Chicken Hollow Road; 2—road cuts on south side of Chicken Hollow Road, 0.1 mile (0.2 km) east of intersection with U.S. Route 62 and 68; Russellville, Ohio–Kentucky, 7.5-minute quadrangle; Union Twp., Brown County, Ohio; 38°46'14"N, 83°48'14"W (at no. 2)

Units.—Grant Lake, Bellevue, Fairview

References.—1: Schumacher, Shrake, Swinford, Brockman, and Wickstrom (1987, stop 2, p. 40, 46, 48, 49, 50); 2: Holland (paper 18 in this volume)

OH-BR-0002 RIPLEY

Along U.S. Routes 68 and 62, 0.2 mile (0.3 km) northeast of intersection with U.S. Route 52; Maysville West, Kentucky–Ohio, 7.5-minute quadrangle; Brown County, Ohio

Units.—Kope

References.—Schumacher, Shrake, Swinford, Brockman, and Wickstrom (1987, stop 3, p. 40, 46, 48-50)

OH-BR-0003 KOPE HOLLOW

Exposures in Kope Hollow, starting about 0.5 km (0.3 mile) east of Levanna, north of U.S. Route 52; mostly Russellville, Ohio–Kentucky, 7.5-minute quadrangle, but also Higginsport, Ohio–Kentucky, 7.5-minute quadrangle; Union Twp., Brown County, Ohio; 38°46'00"N, 83°52'40"W; elevation of base: 151 meters (495 ft), in culvert under U.S. Route 52

Units.—Kope (151 ft/46.0 meters), Point Pleasant (32 ft/9.8 meters); this locality is one of three co-equal type sections of the Kope Formation (other two are KY-MS-0004 and OH-BR-0004)

References.—Weiss and Sweet (1964), Weiss, Edwards, Norman, and Sharp (1965), Diekmeyer (paper 3 in this volume)

OGS measured section no. 13100

OH-BR-0004 RED OAK CREEK

Stream cut 3.2 km (2 miles) north of Ripley; Russellville, Ohio–Kentucky, 7.5-minute quadrangle; Union Twp., Brown

County, Ohio; 38°45'40"N, 83°49'30"W; elevation of base: 175 meters (574 ft)

Units.—Fairview, Kope; this locality is one of three co-equal type sections of the Kope Formation (other two are KY-MS-0004 and OH-BR-0003)

References.—Weiss and Sweet (1964); Diekmeyer (paper 3 in this volume)

OH-BR-0005 EAGLE CREEK

Exposures on unnamed tributary of West Fork Eagle Creek, about 415 meters (1,700 ft) south of junction of Ohio Route 125 and Dr. Faul Road; Russellville, Ohio–Kentucky, 7.5-minute quadrangle; Jefferson Twp., Brown County, Ohio; 38°51'02"N, 83°45'10"W; elevation at base (stream level): 795 ft (242.4 meters)

Units.—Straight Creek (a Mt. Auburn equivalent; 5.5 meters/18.0 ft), Corryville (3.4 meters/11.2 ft)

References.—Goldman (paper 9 in this volume)

OGS measured section no. 16912

OH-BR-0006 GEORGETOWN

Road cuts along both sides of Ohio Route 125 on both sides of White Oak Creek valley, 0.8 mile (1.3 km) west of intersection with U.S. Route 68 in Georgetown; Hamersville, Ohio, and Higginsport, Ohio–Kentucky, 7.5-minute quadrangles; Lewis Twp., Brown County, Ohio; 38°52'33"N, 83°55'53"W

Units.—Bull Fork, Grant Lake (28.8 meters/94.4 ft), Bellevue (14.8 meters/48.6 ft), Fairview (9.2 meters/30.2 ft); Kope exposed in the banks of White Oak Creek south of Ohio Route 125 (Pleasant Twp., 38°52'10"N, 83°55'00"W) and in a road cut on Ohio Route 125 about 0.5 mile (0.8 km) west of city limits of Georgetown (Pleasant Twp., 38°52'15"N, 83°55'00"W)

References.—Lee (1974), Tobin (1982), Holland (1990), Diekmeyer (paper 3 in this volume), Goldman (paper 9 in this volume), Holland (paper 18 in this volume)

OH-BR-0007 GOOSE RUN

Exposure along stream 91 meters (300 ft) west of White Oak Station Road on Goose Run; Hamersville, Ohio, 7.5-minute quadrangle; Scott Twp., Brown County, Ohio; 38°58'47"N, 83°53'38"W

Units.—Straight Creek/Mt. Auburn, Corryville

References.—Goldman (paper 9 in this volume)

OGS measured section no. 16906

OH-BR-0008 RIPLEY #2

Road cut 2.5 miles (4.0 km) north of city limits of Ripley, along west side of U.S. Routes 62/68; Russellville, Ohio–Kentucky, 7.5-minute quadrangle; Union Twp., Brown County, Ohio; 38°46'30"N, 83°49'08"W

Units.—Grant Lake (19.7 meters/64.4 ft), Fairview

References.—Lee (1974), Goldman (paper 9 in this volume), Hay (paper 17 in this volume)

OH-BR-0009 WHITE OAK CREEK

Stream cut along White Oak Creek, approximately 300 meters (1,000 ft) north of New Hope; Hamersville, Ohio, 7.5-minute quadrangle; Scott Twp., Brown County, Ohio; 38°57'57"N, 83°54'36"W; elevation at base (stream level): approximately 860 ft (262 meters)

Units.—Corryville (6.5 meters/21.3 ft)

References.—Goldman (paper 9 in this volume)

OGS measured section no. 16910

OH-BR-0010 WHITE SWAN RUN

Exposures along unnamed creek running south into White Oak Creek near mouth of Cochran Run; Higginsport, Ohio–Kentucky, 7.5-minute quadrangle; Lewis Twp., Brown County, Ohio; 38°50'15"N, 83°56'30"W; elevation at base: 555 ft (169.2 meters)

Units.—Kope (161 ft/49.1 meters), to base of Maysvillian
References.—Weiss, Edwards, Norman, and Sharp (1965),
Diekmeyer (paper 3 in this volume)
OGS measured section no. 13108

Ohio, Butler County

OH-BU-0001 ARMCO STEEL

Hill and road cut on Armco Steel Company property in Middletown; on line between secs. 19 and 13, T. 2, R. 4, Trenton, Ohio, 7.5-minute quadrangle; Butler County, Ohio; elevation: 680-730 ft (207.3-222.5 meters)

Units.—Mt. Auburn (40 ft/12.2 meters), Corryville (10 ft/3.05 meters)

References.—Reinhart (1977a, p. I-35 to I-36, stop 5)

OH-BU-0002 MAUD CUT

Alternate designation.—“Big 4” railroad cut
Exposures along railroad line between Maud and West Chester, east of I-75 viaduct over railroad line and northeast of intersection of I-75 (exit 21) and Cincinnati-Dayton Road (map: Pope and Martin, 1977, page before p. I-1); Glendale, Ohio, 7.5-minute quadrangle; Union Twp., Butler County, Ohio; 39°20'25"N, 84°23'30"W; NOTE: This exposure is difficult to reach because of lack of local paths or roads that lead to tracks. It is also dangerous because the line is active, and trains move through briskly. Do not trespass on railroad property

Units.—Arnheim, Mt. Auburn, Corryville (section: Reinhart, 1977a, p. I-38)

References.—Reinhart (1977a, p. I-37 to I-38, stop 6), Goldman (paper 9 in this volume)

OH-BU-0003 ELK CREEK

Stream exposure along east bank of Elk Creek at first meander bend, 460 meters (1,500 ft) north of Howe Road bridge; Middletown, Ohio, 7.5-minute quadrangle; Madison Twp., Butler County, Ohio; 39°30'20"N, 84°27'36"W

Units.—Corryville (5.5 meters/18.0 ft)

References.—Goldman (paper 9 in this volume)

OGS measured section no. 16893

OH-BU-0004 HUNTS CREEK

Exposure along Hunts Creek, 460 meters (1,500 ft) west of junction of Princeton and Yankee Roads; exposure begins at intersection of a south-flowing stream and Hunts Creek and continues upstream as a series of low cuts on alternating sides of creek; Trenton, Ohio, 7.5-minute quadrangle; Liberty Twp., Butler County, Ohio; 39°23'25"N; 84°23'25"W

Units.—Mt. Auburn (1.3 meters/4.3 ft), Corryville (7.4 meters/24.2 ft)

References.—Goldman (paper 9 in this volume)

OGS measured section no. 16889

OH-BU-0005 MIDDLETOWN

Alternate designation.—“Excello” locality (Hay, 1981)

Road cuts along both sides of Ohio Route 4, just south of bridge across Dicks Creek, about 0.5 km (0.3 mile) south of junction with Ohio Route 73; Trenton, Ohio, 7.5-minute quadrangle; Lemon Twp., Butler County, Ohio; 39°28'40"N, 84°24'58"W

Units.—Mt. Auburn, Corryville; this locality is type section of Hay's (1981) “Excello member” of “Brookville formation”

References.—Hay (1981), Tobin (1982), Goldman (paper 9 in this volume)

OH-BU-0006 WEST MIDDLETOWN

Exposure along unnamed northern tributary of unnamed stream situated 152 meters (500 ft) north of driveway on west side of Trenton-Franklin Road; driveway is 0.8 km (0.5 mile) north of West Middletown; Middletown, Ohio, 7.5-minute quadrangle; Madison Twp., Butler County, Ohio;

39°31'55"N, 84°25'05"W

Units.—Mt. Auburn (5.2 meters/17.1 ft), Corryville (5.0 meters/16.4 ft)

References.—Goldman (paper 9 in this volume)

OGS measured section no. 16885

OH-BU-0007

Stream-bank exposures on south side of Collins (Bull) Run, at Pfeffer Park, just south of Miami University campus; center SE¹/₄, sec. 27, T. 5 N., R. 1 E., Millville, Ohio, 7.5-minute quadrangle; Oxford Twp., Butler County, Ohio

Units.—“trilobite shale unit,” Waynesville

References.—Frey (1987b, loc. 7)

Ohio, Clermont County

OH-CT-0001 BATAVIA #1

Road cut along exit ramp from Ohio Route 32 to Ohio Route 132, 0.5 km (0.3 mile) east of East Fork Little Miami River; Batavia, Ohio, 7.5-minute quadrangle; Batavia Twp., Clermont County, Ohio; 39°05'15"N, 84°10'15"W

Units.—Kope

References.—Hay, Pope, and Frey (1981, stop 1, p. 80-81), Tobin (1982), Meyer and others (1985), Shrake, Schumacher, and Swinford (1988, stop 1-A, p. 31-35, 37), Diekmeyer (paper 3 in this volume)

OH-CT-0002 BATAVIA #2

Road cuts at top of hill on both sides of Ohio Route 32 about 1.2 km (0.7 mile) east of intersection of Ohio Routes 132/222 and Ohio Route 32, east of Batavia; Batavia, Ohio, 7.5-minute quadrangle; Batavia Twp., Clermont County, Ohio; 39°05'10"N, 84°09'55"W

Units.—Bellevue (7.9 meters/25.9 ft), Miamitown (0.9-1.2 meters/3.0-3.9 ft), Fairview (12.1 meters/39.7 ft)

References.—Lee (1974), Hay, Pope, and Frey (1981, stop 2, p. 81-82), Shrake, Schumacher, and Swinford (1988, stop 1-B, p. 31, 32, 35-38), Diekmeyer (paper 3 in this volume), Goldman (paper 9 in this volume)

OH-CT-0004 STONELICK CREEK

Stream cuts along Stonelick Creek upstream and downstream from Ohio Route 131 bridge; Newtonsville, Ohio, plus a bit of the Goshen, Ohio, 7.5-minute quadrangles; Stonelick Twp., Clermont County, Ohio; 39°10'40"N, 84°06'43"W (at Ohio Route 131 bridge)

Units.—Mt. Auburn, Corryville, Bellevue, Fairview; proposed neotype section for Corryville Member of Grant Lake Formation (Schumacher, Swinford, and Shrake, 1991) is part of these Stonelick Creek exposures

References.—Meyer, Osgood, Hinterlong, and Tobin (1981, *in* Meyer and others, 1981, stop 6), Pojeta (1982, 1987), Schumacher (1984), Brandt Velbel (1985, loc. C), Schumacher, Swinford, and Shrake (1991), Diekmeyer (paper 3 in this volume), Goldman (paper 9 in this volume)

OGS measured section nos. 16836, 16837, 16838, 16847, 16848, 16852

OH-CT-0005 WILLIAM LIGHT PAVING COMPANY QUARRY

Abandoned quarry south of Indian Creek, 0.25 mile (0.4 km) east of junction of Ohio Route 232 and Big Indian Road; this intersection is 1.5 miles (2.4 km) east of U.S. Route 52 at Point Pleasant; Laurel, Ohio-Kentucky, 7.5-minute quadrangle; Clermont County, Ohio; 38°53'17"N, 84°12'13"W

Units.—Kope, Point Pleasant

References.—Schumacher, Shrake, Swinford, Brockman, and Wickstrom (1987, stop 4, p. 41, 50-52)

OH-CT-0006 BACKBONE CREEK

Stream cuts along Backbone Creek, 0.27 mile (0.43 km) east of Ohio Route 32 entrance ramp onto Ohio Route 132; Batavia, Ohio, 7.5-minute quadrangle; Batavia Twp., Clermont

County, Ohio; 39°05'20"N, 84°10'10"W

Units.—Corryville, Bellevue (8 meters/26.2 ft), Fairview (30 meters/98.4 ft), Kope (30 meters/98.4 ft)

References.—Meyer, Schumacher, Swinford, Jennette, and Brockman (1985), Jennette (1986), Kepferle, Noger, Meyer, and Schumacher (1987), Diekmeyer (paper 3 in this volume)

OH-CT-0007 EAST FORK RESERVOIR SPILLWAY/W.M. A. HARSHA LAKE

Emergency overflow spillway of East Fork Reservoir (= Wm. A. Harsha Lake); spillway is reached via Slade Road, off Ohio Route 222, 1.2 km (0.7 mile) north of intersection with Ohio Route 125; Batavia, Ohio, 7.5-minute quadrangle; Batavia Twp., Clermont County, Ohio; 39°01'34"N, 84°09'30"W

Units.—Grant Lake, upper Fairview

References.—Diekmeyer (paper 3 in this volume)

OH-CT-0008 O'BANNON CREEK

Gully along east-facing stream exposure 365 meters (1,200 ft) south of Gaynor Farm Road bridge; Goshen, Ohio, 7.5-minute quadrangle; Goshen Twp., Clermont County, Ohio; 39°14'07"N, 84°11'03"W

Units.—Corryville, Bellevue

References.—Goldman (paper 3 in this volume)

OGS measured section no. 16834

OH-CT-0009 TODD RUN

Stream cut 60 meters (200 ft) west of Ohio Route 133 bridge, which is 0.8 mile (1.3 km) south of junction of Ohio Routes 133 and 32; Williamsburg, Ohio, 7.5-minute quadrangle; Williamsburg Twp., Clermont County, Ohio; 39°02'23"N, 84°02'38"W

Units.—Corryville

References.—Goldman (paper 9 in this volume)

OGS measured section no. 16853

OH-CT-0010 BEAR CREEK

Quarry on east side of Bear Creek, adjacent to U.S. Route 52; Moscow, Ohio—Kentucky, 7.5-minute quadrangle; Washington Twp., Clermont County, Ohio; 38°48'00"N, 84°09'20"W; elevation: 497 ft (151.5 meters)

Units.—Eden (70 ft/21.3 meters), Point Pleasant (69 ft/21 meters, basal 12 ft/3.7 meters covered)

References.—Weiss, Edwards, Norman, and Sharp (1965), Diekmeyer (paper 3 in this volume)

OH-CT-0011 BOAT RUN

Stream exposures above Clermontville Road bridge (at east edge of New Richmond 7.5-minute quadrangle) and up left fork toward Mt. Zion Church; Laurel, Ohio—Kentucky, 7.5-minute quadrangle; Monroe Twp., Clermont County, Ohio; 38°55'25"N, 84°14'30"W; elevation at base: 488 ft (149 meters)

Units.—Kope (72 ft/21.9 meters), Point Pleasant (52 ft/15.8 meters)

References.—Weiss, Edwards, Norman, and Sharp (1965), Diekmeyer (paper 3 in this volume)

OGS measured section no. 14934

OH-CT-0012 CHILO

Stream exposures along creek parallel to Ohio Route 222, 0.2 mile (0.32 km) from U.S. Route 52; Moscow, Ohio—Kentucky, 7.5-minute quadrangle; Franklin Twp., Clermont County, Ohio; 38°47'45"N, 84°08'15"W; elevation at base: 590 ft (179.8 meters)

Units.—Kope (30 ft/9.1 meters), Point Pleasant (11 ft/3.4 meters)

References.—Weiss, Edwards, Norman, and Sharp (1965); Diekmeyer (paper 3 in this volume)

OGS measured section no. 13098

OH-CT-0013 HAPPY HOLLOW

Stream exposures in Happy Hollow between U.S. Route 50 and Ohio Route 28 east of Milford; Madeira, Ohio, 7.5-minute quadrangle; Miami Twp., Clermont County, Ohio; 39°10'45"N, 84°16'00"W; elevation at base: 558 ft (170.1 meters)

Units.—Weiss, Edwards, Norman, and Sharp (1965); Kope (87 ft/26.5 meters) from below middle bridge to base of Maysville; Jennette (1986); Kope (16.25 meters/53.3 ft), Fairview (1.0 meter/3.3 ft)

References.—Weiss, Edwards, Norman, and Sharp (1965); Jennette (1986); Diekmeyer (paper 3 in this volume)

OGS measured section no. 13105

OH-CT-0014 INDIAN CREEK

Composite section consisting of a 21-ft (6.4-meter) cutbank at junction of North Fork with Indian Creek and an upper 59 ft (18.0 meters) beginning below Ohio Route 743 bridge over Indian Creek; Laurel, Ohio—Kentucky, and Moscow, Ohio—Kentucky, 7.5-minute quadrangles; Washington Twp., Clermont County, Ohio; part 1: 38°53'20"N, 84°10'42"W; part 2: 38°52'15"N, 84°09'40"W; elevation at base: 622 ft (189.6 meters)

Units.—part 1 is lower Kope; part 2, which may overlap part 1, is middle Kope

References.—Weiss, Edwards, Norman, and Sharp (1965); Diekmeyer (paper 3 in this volume)

OGS measured section nos. 13097 (parts 1 and 2), 13112 (part 2)

OH-CT-0015 NINEMILE CREEK

Two exposures: 1—upper exposure along creek from bridge at fork of Ninemile-Tobasco Road and Bradbury [Bradburg on quad] Road 1.75 miles (2.8 km) SSW of Withamsville, Ohio; 2—cutbank on lower Ninemile Creek opposite junction of Ninemile-Tobasco Road and Nordyke Road; Withamsville, Ohio—Kentucky, 7.5-minute quadrangle; Pierce Twp., Clermont County, Ohio; 1: 39°02'30"N, 84°17'30"W; 2: 39°02'10"N, 84°18'30"W; elevation at base: 1 = 575 ft (175.3 meters); 2 = 490 ft (149.4 meters)

Units.—1: Kope (154 ft/46.9 meters to base of Maysville); 2: lower Kope (31 ft/9.4 meters, overlaps upper exposure by about 8 ft/2.4 meters)

References.—Weiss, Edwards, Norman, and Sharp (1965); Diekmeyer (paper 3 in this volume)

OGS measured section nos. 13094 (2), 13138 (1)

OH-CT-0016 NORTH POINT PLEASANT

Unnamed ravine 0.3 mile (0.5 km) northwest of Opossum Hollow, 1.1 miles (1.7 km) west from Point Pleasant; base adjacent to U.S. Route 52; Laurel, Ohio—Kentucky, 7.5-minute quadrangle; Monroe Twp., Clermont County, Ohio; 38°54'25"N, 84°14'35"W; elevation at base: 505 ft (153.9 meters)

Units.—Kope (65 ft/19.8 meters), Point Pleasant (53 ft/16.2 meters)

References.—Weiss, Edwards, Norman, and Sharp (1965); Diekmeyer (paper 3 in this volume); see also Foerste (1905, p. 151)

OGS measured section no. 13096

OH-CT-0017 SLICKAWAY RUN

Stream exposures along creek from Cedron to Felicity beginning with lowest rock in run; Felicity, Ohio—Kentucky, 7.5-minute quadrangle; Franklin Twp., Clermont County, Ohio; 39°48'15"N to 39°49'45"N, 84°03'00"W to 84°04'50"W; elevation at base: 551 ft (167.9 meters)

Units.—Kope (171 ft/52.1 meters), Point Pleasant (4 ft/1.2 meters); upper Point Pleasant also exposed on Bullskin Creek below mouth of Slickaway Run

References.—Weiss, Edwards, Norman, and Sharp (1965); Diekmeyer (paper 3 in this volume)

OGS measured section no. 13099

OH-CT-0018 TWELVEMILE CREEK

Stream exposures along creek beginning at bridge of Ohio Route 132 where Fagin Run branches to the north; New Richmond, Kentucky–Ohio, and Laurel, Ohio–Kentucky, 7.5-minute quadrangles; Ohio Twp., Clermont County, Ohio; 38°58'00"N, 84°15'40"W; elevation at base: 496.5 ft (151.3 meters)

Units.—Kope (108.5 ft/33.1 meters), Point Pleasant (6.5 ft/2.0 meters)

References.—Weiss, Edwards, Norman, and Sharp (1965)
OGS measured section no. 13095

OH-CT-0019 BECKJORD

Road cuts on north side of eastbound U.S. Route 52, just west of Beckjord Coal Plant, 0.6 mile (1 km) east of Ohio Route 749; Withamsville, Ohio–Kentucky, 7.5-minute quadrangle; Clermont County, Ohio; 39°00'07"N, 84°17'49"W

Units.—Kope, Point Pleasant

References.—Holland (paper 18 in this volume)

OH-CT-0020 POINT PLEASANT

Stream exposures on north side of U.S. Route 52, 0.3 mile (0.5 km) west of western city limits of Point Pleasant; Laurel, Ohio–Kentucky, 7.5-minute quadrangle; Clermont County, Ohio; 38°54'02"N, 84°14'13"W

Units.—Kope, Point Pleasant

References.—Holland (paper 18 in this volume)

Ohio, Clinton County

OH-CN-0001 COWAN LAKE

Exposures in spillway of Cowan Lake and on north bank of lake adjacent to spillway; Clarksville, Ohio, 7.5-minute quadrangle; Vernon Twp., Clinton County, Ohio; 39°23'22"N, 83°55'38"W

Units.—Whitewater, Liberty, Bull Fork

References.—Frey (1987a, loc. I), Holland (paper 18 in this volume)

OH-CN-0002

Stream-bank exposures along Stony Hollow, just north of Todds Fork, immediately north of Clarksville; Clarksville, Ohio, 7.5-minute quadrangle; Vernon Twp., Clinton County, Ohio; 39°24'30"N, 83°59'25"W

Units.—"*Treptoceras duseri* shale unit," Waynesville

References.—Frey (1987b, loc. 3)

Ohio, Greene County

OH-GR-0001

Road cut on west side of U.S. Route 68, 0.6 km (0.4 mile) south of Goes Station; Yellow Springs, Ohio, 7.5-minute quadrangle; Greene County, Ohio; 39°45'30"N, 83°55'30"W

Units.—Preachersville Member of Drakes

References.—Frey (1987a, loc. K)

Ohio, Hamilton County

OH-HA-0001 FAIRVIEW PARK

Exposure in city park, reached via a one-way drive off McMillan Street west of intersection with Ravine Street; drive debouches into Ravine Street just south of intersection with Warner Street; city of Cincinnati, Cincinnati West, Ohio, and Covington, Kentucky–Ohio, 7.5-minute quadrangles (Luft, 1971; Ford, 1974); Hamilton County, Ohio; 39°07'30"N, 84°31'50"W

Units.—Bellevue, Miamitown, Fairview; name of Fairview Formation was taken from this locality

References.—Diekmeyer (paper 3 in this volume)

OH-HA-0002 EMMING STREET

Road cut (old quarry face) on north side of Emming Street,

west of Clifton Avenue, between Wheeler and Stratford, and just below the transmitting tower; sec. 19, city of Cincinnati, Covington, Kentucky–Ohio, 7.5-minute quadrangle (Luft, 1971, Ford, 1974); Hamilton County, Ohio; 39°07'22"N, 84°31'21"W

Units.—Corryville, Bellevue, Miamitown, Fairview

References.—Diekmeyer (paper 3 in this volume), Dattilo (paper 7 in this volume), Goldman (paper 9 in this volume)

OH-HA-0003 BELLEVUE HILL

Alternate designation.—Clifton Hill (Ford, 1967)

Exposure between top of cliff in Bellevue Hill Park and Clifton Avenue; sec. 13, city of Cincinnati, Covington, Kentucky–Ohio, 7.5-minute quadrangle (Luft, 1971; Ford, 1974); Hamilton County, Ohio; 39°07'17"N, 84°31'14"W; elevation at base: 645 ft (196.6 meters); NOTE: steep (not for children or the clumsy)

Units.—Corryville, Bellevue (22 ft/6.7 meters), Miamitown (5 ft/1.5 meters), Fairview (108 ft/32.9 meters), Kope (37 ft/11.3 meters). NOTE: Bellevue Limestone was named for the Bellevue House, a drinking establishment that was located on the hilltop above this exposure. This exposure is Ford's (1967) neotype section for Fairview Formation. The Bellevue-Corryville contact is set at a prominent limestone bed that forms a small ledge near the cliff edge

References.—Ford (1967), Diekmeyer (paper 3 in this volume), Dattilo (paper 7 in this volume), Goldman (paper 9 in this volume)

OH-HA-0004 RICE AND GAGE STREETS

Alternate designation.—Christ Hospital

Road cut at intersection of Rice and Gage Streets, Cincinnati, on grounds of Christ Hospital, under hospital heliport; opposite 2227 Gage Street, Cincinnati, Ohio; this locality is on a spur of the hill called Mount Auburn; the Jackson Hill Park locality (OH-HA-0018) is situated on the next spur to the south; sec. 13, Fractional R. 2, T. 3, city of Cincinnati, Covington, Kentucky–Ohio, 7.5-minute quadrangle (Luft, 1971; Ford, 1974); Hamilton County, Ohio; 39°07'15"N, 84°30'50"W; 4332909 N, 714976 E, UTM zone 16; elevation at base: 663 ft (202.1 meters). NOTE: because this locality is on the grounds of Christ Hospital, permission must be obtained from the hospital before this site is visited, or security guards will help you complete your visit in a prompt manner

Units.—Bellevue, Miamitown, Fairview; this exposure was designated the neotype section of the Bellevue Limestone by Ford (1967)

References.—Ford (1965, 1967), Diekmeyer (paper 3 in this volume), Dattilo (paper 7 in this volume)
OGS measured section no. 15372

OH-HA-0005 BLUE ROCK ROAD

Road cut 0.3 mile (0.5 km) southeast of junction of I-275 (Exit 31) and Blue Rock Road; just north of exit ramp from the Cross County Highway; Colerain Twp., T. 2, R. 1, Cincinnati West, Ohio, 7.5-minute quadrangle (Ford, 1974); Hamilton County, Ohio; 39°13'47"N, 84°37'16"W

Units.—Sunset, Mt. Auburn, Corryville (mostly buried, at base of cut); measured section in Krumpolz (1980, p. 170-175)

References.—Ford (1967), Krumpolz (1980, loc. 1), Meyer, Krumpolz, & Tobin (1981, in Meyer and others, 1981, stop 7, p. 68-70), Tobin (1982), Rassman (1984), Schumacher (1984), Meyer (1990), Goldman (paper 9 in this volume), Holland (paper 18 in this volume)

OH-HA-0006 GRAND AVENUE

East-facing cliff (quarry) just west of Grand Avenue, Cincinnati (opposite 2196 Grand Avenue); near center of north edge of topographic sheet; city of Cincinnati, Covington, Kentucky–Ohio, 7.5-minute quadrangle (Ford, 1974); Hamilton County, Ohio; 39°07'10"N, 84°33'20"W; elevation at base:

585 ft (178.3 meters)

Units.—Kope (94 ft/28.7 meters from quarry floor to Maysville); type section of Grand Avenue Member of Kope

References.—Weiss, Edwards, Norman, and Sharp (1965), Ford (1967), Diekmeyer (paper 3 in this volume)

OGS measured section no. 13277

OH-HA-0007 HARRISON

Road cut on Marvin Road on north side of I-74 at New Bidinger Road; sec. 18, T. 2 N., R. 1 E., Harrison, Ohio–Indiana, 7.5-minute quadrangle; Harrison Twp., Hamilton County, Ohio

Units.—Fairmount and Mt. Hope Members of Fairview

References.—Reinhart (1977a, stop 7, p. I-40)

OH-HA-0008 MIAMITOWN WEST

Alternate designation.—“H’way cuts W. of Miamitown” (Ford, 1967, fig. 6, p. 926)

Originally a road cut on I-74 about 1 mile (1.6 km) west of Miamitown, Ohio; subsequently an interchange between I-74 and I-275 was constructed at the location; locality consists of six road cuts surrounding the various interchange roads; the measured section “highway cuts west of Miamitown” of Ford (1967, fig. 10) was a composite that ran essentially all the way from this locality to that designed herein as “Miamitown/Hamilton-Cleves exit” (OH-HA-0009); sec. 35, Addyston, Ohio–Kentucky, 7.5-minute quadrangle (Ford, 1972); Whitewater Twp., Hamilton County, Ohio; 39°13'14"N, 84°44'10"W; 4343499 N, 695446 E, UTM zone 16. NOTE: as with all interstate highway cuts, permission from the Ohio Department of Transportation is required to stop at this site

Units.—Miamitown, Fairview (including North Bend Tongue), Kope (including Grand Avenue Member and Wesselman Tongue); this locality is the type section of the Miamitown Shale

References.—Ford (1965, 1967), Dattilo (paper 7 in this volume)

OH-HA-0009 MIAMITOWN/HAMILTON-CLEVES EXIT

Road cut on exit ramp from I-74/I-275 east, leading to U.S. Route 52/Ohio Route 128 (Hamilton-Cleves Road), 0.8 mile (1.3 km) SSW of crossroads in center of Miamitown; the measured section “highway cuts west of Miamitown” of Ford (1967) was a composite that ran essentially all the way from this locality to that designed herein as “Miamitown West” (OH-HA-0008); center SE¹/₄SE¹/₄ sec. 1, T. 1 N., R. 1 E., Addyston, Ohio–Kentucky 7.5-minute quadrangle (Ford, 1972); Whitewater Twp., Hamilton County, Ohio; 39°12'21"N, 84°42'39"W; 4341928 m N, 697649 m E, UTM zone 16; elevation at base: 550 ft (167.6 meters)

Units.—Grand Avenue Member and Wesselman Tongue of Kope; Jennette (1986); Fairview (0.2 meter/0.7 ft), Kope or McMicken (19.9 meters/65.3 ft)

References.—Ford (1965, 1967), Jennette (1986), Diekmeyer (paper 3 in this volume), Cuffey (paper 4 in this volume)

OGS measured section no. 15378

OH-HA-0010 NORTH BEND

South-facing bank of a southwest-flowing stream slightly more than 1 mile (1.6 km) ENE of North Bend; Miami Twp., Addyston, Ohio–Kentucky, 7.5-minute quadrangle (Ford, 1972); Hamilton County, Ohio; 39°09'15"N, 84°44'00"W; elevation: 618-630 ft (188-192 meters)

Units.—North Bend Tongue of Fairview; this locality is the type section of the North Bend Tongue

References.—Ford (1965; 1967, p. 929); Diekmeyer (paper 3 in this volume)

OH-HA-0011 WESSELMAN

East-facing embankment of an unnamed creek that is followed by Wesselman Road, about 400 ft (122 meters) south of Zion

Hill Road; NE¹/₄NE¹/₄NE¹/₄ sec. 4, Addyston, Ohio–Kentucky, 7.5-minute quadrangle (Ford, 1972); Miami and Green Twps., Hamilton County, Ohio; between 39°10'45"N and 39°12'30"N, and 84°40'45"W and 84°41'15"W

Units.—Wesselman Tongue of Kope, bounded above and below by rock of the Fairview; this locality is the type section of the Wesselman Tongue

References.—Ford (1965; 1967, p. 928), Diekmeyer (paper 3 in this volume)

OH-HA-0012 AIKEN SCHOOL

Alternate designation.—Aiken High School

Composite of three sites: 1—north cut of athletic field of Aiken High School (sec. 30); 2—excavation west of school on Belmont Avenue (sec. 30); 3—excavation at Hammond North Apartments, east of Hamilton Avenue (sec. 29); secs. 29 and 30, Fractional R. 2, T. 3, city of Cincinnati, Cincinnati West, Ohio, 7.5-minute quadrangle (Ford, 1974); Hamilton County, Ohio; 39°11'33"N, 84°32'58"W; elevation: at base of 1, 735 ft (224.0 meters); at base of 2, 854 ft (260.3 meters); at base of 3, 813 ft (247.8 meters)

Units.—Bellevue (23 ft/7.0 meters), Miamitown (5 ft/1.5 meters), Fairview (26 ft/7.9 meters)

References.—Ford (1965; 1967, p. 919, 927), Tobin (1982), Diekmeyer (paper 3 in this volume), Holland (paper 18 in this volume)

OGS measured section nos. 15375 (1), 15456 (3), 15457 (2) (originally all three measured sections were no. 15375)

OH-HA-0013 BALD KNOB

Hill in Cincinnati, Ohio; quarry west of State Avenue; road cut 0.9 mile (1.5 km) west of intersection of State Avenue and Lehman Road; Covington, Kentucky–Ohio 7.5-minute quadrangle (Luft, 1971; Ford, 1974); Hamilton County, Ohio; 39°06'00"N, 84°33'05"W; elevation at base: 630 ft (192 meters). NOTE: prior to the 1930's Bald Knob was taller than it now is; the top was removed to provide fill material for the site of Cincinnati's Union Terminal, in order to bring that site above flood stage

Units.—Weiss, Edwards, Norman, and Sharp (1965): 22 ft (6.7 meters) from toe of south end of bluff to base of Maysvillian; Ford (1967, fig. 6): Miamitown, Fairview, and Kope (including Grand Avenue Member); Jennette (1986): Fairview (9.7 meters/31.8 ft), Kope (7.6 meters/24.9 ft)

References.—Weiss, Edwards, Norman, and Sharp (1965), Ford (1967), Sweet (1979), Tobin (1982), Jennette (1986), Diekmeyer (paper 3 in this volume)

OGS measured section nos. 14807, 14808

OH-HA-0014 BRIARLY CREEK

Stream cut of Briarly Creek, from creek bed to top of southeast bank, southeast of bend in Sheed Road; sec. 30, Fractional R. 2, T. 2, Addyston, Ohio–Kentucky, 7.5-minute quadrangle (Ford, 1972); Green Twp., Hamilton County, Ohio; 39°12'20"N, 84°38'45"W; elevation of base: 602 ft (183.5 meters)

Units.—Miamitown, Fairview, Wesselman Tongue of Kope (5 ft/1.5 meters), North Bend Tongue of Fairview (11 ft/3.4 meters), other Kope (19 ft/5.8 meters)

References.—Ford (1965, 1967), Diekmeyer (paper 3 in this volume)

OGS measured section no. 15382

OH-HA-0015 CAMP CLAYBANKS

Cliff opposite Camp Claybanks, by spillway on West Fork Mill Creek; sec. 17, R. 1, T. 3, Glendale, Ohio, 7.5-minute quadrangle; Hamilton County, Ohio; elevation of base: 624 ft (190.2 meters); 39°15'35"N, 84°29'30"W

Units.—Bellevue (3 ft/0.9 meter), Miamitown (8 ft/2.4 meters), Fairview (84 ft/25.6 meters), Kope (12 ft/3.7 meters)

References.—Ford (1965, 1967), Diekmeyer (paper 3 in this volume)

OGS measured section no. 15374

OH-HA-0016 CONGRESS RUN

Stream cuts of Congress Run and tributary to Congress Run along Galbraith Road east of Winton Road; sec. 14, R. 1, T. 3, Cincinnati East and Cincinnati West, Ohio, 7.5-minute quadrangles (Ford, 1974); Springfield Twp., Hamilton County, Ohio; 39°13'00"N, 84°30'15"W; elevation at base: 603 ft (183.8 meters)

Units.—Miami town, Fairview, Wesselman Tongue of Kope, North Bend Tongue of Fairview, plus (Ford, 1967, fig. 6): Kope

References.—Ford (1967), Diekmeyer (paper 3 in this volume), Dattilo (paper 7 in this volume)

OGS measured section no. 15373

OH-HA-0017 CROSBY ROAD

Road cuts on west and east sides of Crosby Road; sec. 13, T. 2 N., R. 1 E., Shandon, Ohio, 7.5-minute quadrangle; Crosby Twp., Hamilton County, Ohio; 39°16'00"N, 84°42'45"W; elevation at base: 627 ft (191 meters)

Units.—Miami town (21 ft/6.4 meters), Fairview (63 ft/19.2 meters), Kope (32 ft/9.7 meters)

References.—Ford (1965, 1967), Diekmeyer (paper 3 in this volume), Dattilo (paper 7 in this volume)

OGS measured section no. 15379

OH-HA-0018 JACKSON HILL PARK

Hillside opposite 42 Mulberry Street, beneath crest of Jackson Hill Park, Cincinnati, Ohio; Jackson Hill is a spur of the hill called Mount Auburn; Rice and Gage Streets locality (OH-HA-0004) is located on next spur to north; Fractional R. 2, T. 3, Covington, Kentucky–Ohio, 7.5-minute quadrangle (Luft, 1971; Ford, 1974); Hamilton County, Ohio; 39°07'05"N, 84°30'50"W; elevation at base: 761 ft (232.0 meters)

Units.—Bellevue (6.7 meters/22 ft), Miami town (1.2 meters/4.0 ft), Fairview (33.2 meters/109 ft), Kope (6.1 meters/20 ft)

References.—Ford (1967, p. 927, 929); Diekmeyer (paper 3 in this volume)

OH-HA-0019 MIAMI STATION

Highway cut south of "eng. sta. 480" (OGS measured section no. 15381), I-74 east, due south of Miami Station; sec. 1, R. 1, T. 1, Addyston Ohio–Kentucky, 7.5-minute quadrangle (Ford, 1972); Colerain Twp., Hamilton County, Ohio; 39°12'40"N, 84°41'25"W; elevation at base: 554 ft (168.9 meters)

Units.—Miami town, Fairview (29 ft/8.8 meters), Wesselman Tongue of Kope (27 ft/8.2 meters), North Bend Tongue of Fairview (8 ft/2.4 meters), Kope (53 ft/16.2 meters), Grand Avenue Member of Kope (11 ft/3.4 meters)

References.—Ford (1965; 1967, p. 919, 925, 926), Diekmeyer (paper 3 in this volume)

OGS measured section no. 15381

OH-HA-0020 MUDDY CREEK

Composite of a number of individual exposures: 1—stream cut just south of Muddy Creek Road bridge 1.0 km (0.6 mile) east of junction of Muddy Creek Road and Ebenezer Road, in sec. 19; 2—creek bed and banks of Muddy Creek in sec. 25; 3—cliff section east of Muddy Creek in sec. 25; 4—cliff section north of Muddy Creek in sec. 19; 5—exposure 0.1 mile (0.2 km) south of the intersection of Muddy Creek Road and Devils Backbone Road, in sec. 25; secs. 19 and 25, T. 2, Fractional R. 2, Addyston, Ohio–Kentucky, 7.5-minute quadrangle (Ford, 1972); Green Twp., Hamilton County, Ohio; 39°07'45" to 39°08'00"N, 84°38'28" to 84°40'00"W; elevation at base: 533 ft (162.5 meters). NOTE: this creek is polluted and may contain raw sewage

Units.—Corryville, Bellevue (29 ft/8.8 meters), Miami town (5 ft/1.5 meters), Fairview (67 ft/20.4 meters), Wesselman Tongue of Kope (4 ft/1.2 meters), North Bend Tongue of Fairview (20 ft/6.1 meters), Grand Avenue Member of Kope

(12 ft/3.7 meters), Kope (106 ft/32.3 meters)

References.—Ford (1965; 1967, p. 919, 925, 926, 929), Jennette (1986), Diekmeyer (paper 3 in this volume), Dattilo (paper 7 in this volume), Goldman (paper 9 in this volume), Holland (paper 18 in this volume)

OGS measured section nos. 15376 (2), 15458 (3), 15459 (1)

OH-HA-0021 RIVERSIDE

Exposure in west branch of unnamed stream that enters Ohio River at Riverside, Cincinnati; base of section in stream bed north of intersection of Hillside Avenue and Tyler Street; sec. 10, Fractional R. 1, T. 3, Covington, Kentucky–Ohio 7.5-minute quadrangle (Luft, 1971; Ford, 1974); Delhi Twp., Hamilton County, Ohio; 39°04'35"N, 84°36'30"W; elevation at base: 606 ft (184.7 meters)

Units.—Fairview (39 ft/11.9 meters), Kope (81 ft/24.7 meters), Grand Avenue Mbr. of Kope (11 ft/3.4 meters)

References.—Ford (1965; 1967, p. 919, 925, 927), Diekmeyer (paper 3 in this volume)

OGS measured section no. 15384

OH-HA-0022 SEKITAN ROAD

Cliff section north of Sekitan Road and drainage channel 700 ft (213.4 meters) due east of first section; sec 8, Fractional R. 2, T. 1, Addyston, Ohio–Kentucky, 7.5-minute quadrangle (Ford, 1972); Hamilton County, Ohio; 39°08'40"N, 84°43'00"W; elevation at base: 566 ft (172.5 meters)

Units.—Miami town, Fairview (26 ft/7.9 meters), Wesselman Tongue of Kope (14 ft/4.3 meters), North Bend Tongue of Kope (10 ft/3.0 meters), Kope (62 ft/18.9 meters), Grand Avenue Member of Kope

References.—Ford (1965; 1967, p. 919, 926); Diekmeyer (paper 3 in this volume)

OGS measured section no. 15377

OH-HA-0023 SHEITS ROAD

Exposures along two branches of an unnamed stream: 1—branch that follows Sheits Road, just west of intersection of stream and I-275 near Dornbusch, Ohio; 2—west-flowing branch and a portion of the branch of #1 crossing Sheits Road downstream of #1; secs. 21 and 15, R. 1, T. 1, Cincinnati West, Ohio, 7.5-minute quadrangle (Ford, 1974); Colerain Twp., Hamilton County, Ohio; 39°14'50"N, 84°37'24"W; elevation at base: 698 ft (212.8 meters)

Units.—Corryville (18.4 ft/5.6 meters), Bellevue (42 ft/12.8 meters), Miami town (14 ft/4.3 meters), Fairview (38 ft/11.6 meters)

References.—Ford (1967, p. 919, 926), Diekmeyer (paper 3 in this volume), Dattilo (paper 7 in this volume), Goldman (paper 9 in this volume)

OGS measured section nos. 15380, 16872

OH-HA-0024 WEST FORK

Creek bed and banks of West Fork, Mt. Airy Forest; secs. 33 and 34, Fractional R. 2, T. 3, sec. 4, Fractional R. 2, T. 2, Cincinnati West, Ohio, 7.5-minute quadrangle (Ford, 1974); Hamilton County, Ohio; 39°09'30"N, 84°34'45"W; elevation at base: 530 ft (161.5 meters)

Units.—Miami town, Fairview (43 ft/13.1 meters), Wesselman Tongue of Kope (11 ft/3.4 meters), North Bend Tongue of Fairview (19 ft/5.8 meters), Kope (110 ft/33.5 meters), Grand Avenue Member of Kope

References.—Ford (1965; 1967, p. 919, 925, 927), Diekmeyer (paper 3 in this volume)

OGS measured section no. 15383

OH-HA-0025 WYNBROOK APARTMENTS

Exposure behind (north of) Wynbrook Apartments east of Winton Road just north of intersection with Dutch Colony Drive, Cincinnati; Dutch Colony Drive is about 2.0 miles (3.2 km) north of intersection of Spring Grove Avenue and Winton Road; Cincinnati West, Ohio, 7.5-minute quadrangle (Ford,

1974); Cincinnati, Hamilton County, Ohio; 39°11'25"N, 84°31'10"W

Units.—Mt. Auburn, Corryville, Bellevue, Miamitown

References.—Diekmeyer (paper 3 in this volume); Goldman (paper 9 in this volume)

OH-HA-0026 DORNBUSCH

Stream cut in west-flowing stream just north of town of Dornbusch and south-southeast of stream's intersection with I-275; Cincinnati West, Ohio, 7.5-minute quadrangle (Ford, 1974); Colerain Twp., Hamilton County, Ohio; 39°14'52"N, 84°36'30"W

Units.—Corryville (11.2 meters/36.7 ft)

References.—Goldman (paper 9 in this volume)

OGS measured section no. 16873

OH-HA-0027 McMILLAN STREET

Exposure in empty lot west of 16 McMillan Street and just northwest of junction of McMillan and Vine Streets, Cincinnati; Cincinnati West, Ohio, 7.5-minute quadrangle (Ford, 1974); Cincinnati, Hamilton County, Ohio; 39°07'39"N, 84°30'52"W

Units.—Corryville; this locality is in the type area of the Corryville

References.—Goldman (paper 9 in this volume)

OH-HA-0028 RAVINE STREET

Poor, mostly buried exposure behind Seminole Apartments southwest of intersection of Ravine and McMillan Streets, Cincinnati; Cincinnati West, Ohio, 7.5-minute quadrangle (Ford, 1974); Hamilton County, Ohio; 39°07'43"N, 84°31'36"W

Units.—Corryville

References.—Goldman (paper 9 in this volume)

OH-HA-0029 RUMPKE LANDFILL

Inaccessible (buried) exposure in field east of U.S. Route 27 N (Colerain Avenue), 2.5 km (1.5 miles) north of I-275; NW¹/₄ sec. 11, R. 1, T. 2, Greenhills, Ohio, 7.5-minute quadrangle; Colerain Twp., Hamilton County, Ohio; 39°16'33"N, 84°36'03"W

Units.—Corryville, Mt. Auburn (now buried as a result of landfill)

References.—Tobin (1982), Brandt Velbel (1985, loc. D), Goldman (paper 9 in this volume)

OH-HA-0030 SHARONVILLE

Exposure in upper half of field at Sharonville industrial park, situated at end of 800-foot (243-meter) lane (tram way) north of Hauck Road and 0.4 mile (0.7 km) east of intersection of Hauck and Reading Roads; Glendale, Ohio, 7.5-minute quadrangle; Sycamore Twp., Hamilton County, Ohio; 39°17'38"N, 84°24'07"W

Units.—Corryville, Miamitown, Bellevue, Fairview

References.—Goldman (paper 9 in this volume)

OH-HA-0031 WESTWOOD-NORTHERN BOULEVARD

Exposure in gully northwest of the junction of Westwood-Northern Boulevard and Gobel Street, Cincinnati; north end of Gobel Street has been closed off to traffic; Cincinnati West, Ohio, 7.5-minute quadrangle (Ford, 1974); Hamilton County, Ohio; 39°08'50"N, 84°39'45"W

Units.—? upper Corryville

References.—Goldman (paper 9 in this volume)

OH-HA-0032 GALBRAITH ROAD

Stream cut along Kugler Mill Road, the eastward continuation of Galbraith Road; SE¹/₄ sec. 25, R. 1, Twp. 5, Between the Miamis land survey, Madeira, Ohio, 7.5-minute quadrangle; Symmes Twp., Hamilton County, Ohio; 39°11'23"N, 84°18'32"W; elevation at base: 648 ft (197.5 meters)

Units.—Kope (2 ft/0.6 meter to base of Fairview)

References.—Ford (1965), Diekmeyer (paper 3 in this volume) OGS measured section no. 13104

OH-HA-0033 GALBRAITH ROAD, CLIFF SECTION

Stream bed and banks of south tributary to Congress Run parallel to Hollyhock Drive, in city of Wyoming; sec. 14, R. 1, T. 3, Cincinnati West, Ohio, 7.5-minute quadrangle (Ford, 1974); Springfield Twp., Hamilton County, Ohio; 39°13'30"N, 84°30'15"W; elevation at base: 665 ft (202.7 meters)

Units.—Bellevue (20 ft/6.1 meters), Miamitown (7 ft/2.1 meters), Fairview (73 ft/22.3 meters), North Bend Tongue of Fairview (18 ft/5.5 meters), Kope (34 ft/10.4 meters)

References.—Ford (1965), Diekmeyer (paper 3 in this volume)

OH-HA-0034 EIGHTMILE CREEK

Stream exposures along Eightmile Creek near U.S. Route 52 up right fork subparallel to road to Cherry Grove; Withamsville, Ohio—Kentucky, 7.5-minute quadrangle; Anderson Twp., Hamilton County, Ohio; 39°33'20"N, 84°19'45"W; elevation at base: 476 ft (145.1 meters)

Units.—Weiss, Edwards, Norman, and Sharp (1965): Eden (140 ft/44.7 meters exposed below Maysville contact, below that another 50 ft/15.2 meters, mostly covered, to a 6-ft/1.8-meter ledge of Cynthiana Formation); Jennette (1986): Fairview (1.0 meter/3.3 ft), Kope (34.9 meters/114.5 ft)

References.—Weiss, Edwards, Norman, and Sharp (1965); Diekmeyer (paper 3 in this volume); Jennette (1986)

OGS measured section no. 13093

OH-HA-0035 MILLVALE

South end of bluff behind Millvale Court housing project on Beekman Street, Cincinnati; W¹/₂SW¹/₄ sec. 27, Fractional R. 2, T. 3, Between the Miamis land survey, Cincinnati West, Ohio, 7.5-minute quadrangle (Ford, 1974); Hamilton County, Ohio; 39°08'30"N, 84°33'15"W; elevation: 545 ft (166.1 meters)

Units.—Kope (106 ft/32.3 meters) in middle and upper parts
References.—Weiss, Edwards, Norman, and Sharp (1965), Diekmeyer (paper 3 in this volume)

OH-HA-0036 RAPID RUN

Stream cuts north of U.S. Route 50 (base near ford close to highway) and 0.25-km (0.16-mile) cut along Bender Road northeast of U.S. Route 50; secs. 23, 24, 29, Fractional R. 1, T. 3, Between the Miamis land survey, Burlington, Kentucky—Ohio, 7.5-minute quadrangle (Ford, 1972); Delhi Twp., Hamilton County, Ohio; 39°05'50"N, 84°40'15"W; elevation at base: 475 ft (144.8 meters)

Units.—Kope (202 ft/61.6 meters to base of Maysvillian)

References.—Ford (1965), Weiss, Edwards, Norman, and Sharp (1965), Diekmeyer (paper 3 in this volume)

OGS measured section no. 13106

OH-HA-0037 PABCO/TIMBERVIEW APARTMENTS

Two exposures: 1—outcrop behind Pabco Fluid Power Co., intersection of U.S. Route 50 and Bender Road; 2—outcrop behind Timberview Apartments, 5564 Hillside Avenue; sec. 29, Fractional R. 1, T. 3, Between the Miamis land survey, Burlington, Kentucky—Ohio, 7.5-minute quadrangle (Ford, 1972); Delhi Twp., Hamilton County, Ohio; 1—39°06'00"N, 84°40'20"W; 2—39°06'15"N, 84°40'40"W

Units.—Fairview (0.5 meter/1.6 ft), Kope (29.4 meter/96.5 ft)

References.—Jennette (1986), Diekmeyer (paper 3 in this volume)

OH-HA-0038 MONTANA AVENUE

Alternative name.—I-74/Montana Avenue

Road cut along eastbound entrance ramp onto I-74 from Montana Avenue; Cincinnati West, Ohio, 7.5-minute quadrangle (Ford, 1974); Cincinnati, Hamilton County, Ohio; 39°09'15"N, 84°33'45"W

Units.—Fairview (0.3 meter/1.0 ft), Kope 15.6 meters/51.2 ft)

References.—Jennette (1986), Diekmeyer (paper 3 in this volume)

OH-HA-0039 MT. AIRY FOREST

Series of large road cuts on either side of I-74, between mile 15.6 and mile 16.8, where highway passes through Mt. Airy Forest; sec. 4, Fractional R. 2, T. 2, Cincinnati West, Ohio, 7.5-minute quadrangle (Ford, 1974), Green Twp., Hamilton County, Ohio; 39°10'05"N, 84°34'52"W; 438000 N, 709000 E, UTM zone 16; NOTE: this is an interstate cut; permission from the Ohio Department of Transportation is required for close examination and sampling. Cut is steep and climbing it may be risky both to the geologist and to motorists
Units.—Corryville, Bellevue, Miamitown, Fairview, Kope
References.—Dattilo (paper 7 in this volume)

OH-HA-0040 DELHI PIKE

Exposure on north side of Delhi Pike opposite intersection with Mt. Alverno Road in sec. 5 (note that Mt. Alverno Road also intersects Delhi Pike in sec. 11); SW¹/₄ sec. 5, Fractional R. 1, T. 3, Covington, Kentucky–Ohio, 7.5-minute quadrangle (Luft, 1971; Ford, 1974), Delhi Twp., Hamilton County, Ohio; 39°05'38"N, 84°35'35"W; 4329736 N, 708168 E, UTM zone 16
Units.—Miamitown, Fairview
References.—Dattilo (paper 7 in this volume)
OGS measured section no. 16879

OH-HA-0041 ASHTREE DRIVE AND HAMILTON AVENUE

Alternate designations.—Ashtree Shopping Center; Hamilton Avenue
Cut northwest of intersection of Ashtree Drive and Hamilton Avenue, Cincinnati; flat area in front of cut once was occupied by the Ashtree Shopping Center; for a time the building was the Southern Ohio College Technical Center, and a more recent tenant was the MTA School; W¹/₂NE¹/₄SW¹/₄ sec. 29, Fractional R. 2, T. 3, Cincinnati West, Ohio, 7.5-minute quadrangle (Ford, 1974); Cincinnati, Hamilton County, Ohio; 39°10'45"N, 84°32'50"W
Units.—lower Bellevue, Miamitown, Fairview
References.—Dattilo (paper 7 in this volume)

OH-HA-0042 BOUDINOT AVENUE

Construction exposure, now covered, two blocks south of intersection of Boudinot Avenue and Westwood-Northern Boulevard, Cincinnati; NE¹/₄ sec. 9, Fractional R. 2, T. 2, Cincinnati West, Ohio, 7.5-minute quadrangle (Ford, 1974); Hamilton County, Ohio; 39°09'38"N, 84°36'07"W
Units.—Liberty (Brandt Velbel, 1985, fig. 2), Waynesville and Arnheim (Stephen H. Felton, personal commun., 1998)
References.—Brandt Velbel (1985, loc. G)

Ohio, Warren County

OH-WA-0001 CAESAR CREEK EMERGENCY OVERFLOW SPILLWAY

Emergency overflow spillway of Caesar Creek Reservoir; Oregonia, Ohio, 7.5-minute quadrangle; Warren County, Ohio; 39°28'49"N, 84°03'25"W; 4373976 m N, 753156 m E, UTM zone 16; elevation of uppermost exposed strata: 925 ft (281.9 meters); elevation of Clarksville Road in middle of spillway: 883 ft (269.1 meters)
Units.—Whitewater, Liberty, Waynesville
References.—Staursky (1981), Schumacher and Ausich (1983), Brandt Velbel (1985, loc. B), Schumacher, Shrake, Swinford, Brockman, and Wickstrom (1987, stop 5, p. 44, 52-55), Shrake (1992), Shrake, Schumacher, and Swinford (1988, p. 39-44), University of Dayton Department of Geology (1989, stop 4, p. 8-10); Shrake and Schumacher (1989), Schumacher, Shrake, Swinford, Rockwell, and Cuffey (paper 13 in this volume), Holland (paper 18 in this volume)

OH-WA-0002 CAESAR CREEK GORGE

Alternate name.—Caesar Creek Dam
Cut in gorge below dam of Caesar Creek Reservoir; Oregonia, Ohio, 7.5-minute quadrangle; Warren County, Ohio; 39°29'05"N, 84°03'49"W; 4374458 m N, 752555 m E, UTM zone 16; elevation at base: 755 ft (230.1 meters)
Units.—Waynesville, Arnheim
References.—Schumacher and Ausich (1983), Shrake (1992), Schumacher, Shrake, Swinford, Brockman, and Wickstrom (1987, p. 52-55), Shrake, Schumacher, and Swinford (1988, p. 39-44), University of Dayton Department of Geology (1989, stop 3, p. 7-8); Schumacher, Shrake, Swinford, Rockwell, and Cuffey (paper 13 in this volume), Holland (paper 18 in this volume)

OH-WA-0003 FLAT FORK CREEK

Stream bed and vertical cut banks of Flat Fork Creek (southeast of Caesar Creek Reservoir); Oregonia, Ohio, 7.5-minute quadrangle; Massie Twp., Warren County, Ohio; 39°28'47"N, 84°03'00"W
Units.—lower Whitewater, Liberty, topmost Waynesville; before Caesar Creek was dammed in 1978, the complete section from the Arnheim to the Whitewater was exposed in Flat Fork Creek
References.—Wolford (1930), Schumacher, Shrake, Swinford, Brockman, and Wickstrom (1987, p. 52-55)

OH-WA-0004 LICK RUN

Stream exposures along Lick Run west of Ohio Route 132 bridge and 760 meters (2500 ft) north of Blackhawk, at junction of Ohio Routes 132 and 123; Pleasant Plain, Ohio, 7.5-minute quadrangle; Harlan Twp., Warren County, Ohio; 39°20'02"N; 84°04'41"W
Units.—Mt. Auburn, Corryville; measured section in Krumpolz (1980, p. 182-189)
References.—Krumpolz (1980, loc. 3), Goldman (paper 9 in this volume)

OH-WA-0006 SECOND CREEK

Stream cuts along Second Creek below and west of Cozzandale Road bridge; Pleasant Plain, Ohio, 7.5-minute quadrangle; Harlan Twp., Warren County, Ohio; 39°19'25"N, 84°05'31"W
Units.—Mt. Auburn, Corryville
References.—Osgood (1970), Goldman (paper 9 in this volume)

OH-WA-0007

Stream-bank exposures along north fork of Harpers Run, off Strout Road at Camp Whip-Poor-Will G.S.A. outdoor center; Oregonia, Ohio, 7.5-minute quadrangle; Washington Twp., Warren County, Ohio; 39°23'N, 84°05'W
Units.—"*Treptoceras duseri* shale unit," Waynesville
References.—Frey (1987a, loc. E; 1987b, loc. 2)

OH-WA-0008

South-facing road cut along U.S. Route 42, 3 km (1.9 miles) northeast of Waynesville; SW¹/₄SW¹/₄ sec. 26, T. 4, R. 5, Waynesville, Ohio, 7.5-minute quadrangle; Washington Twp., Warren County, Ohio
Units.—"*Treptoceras duseri* shale unit," Waynesville
References.—Frey (1987a, loc. F); see also Frey (1987b, loc. 1)

ALPHABETICAL LIST BY LOCALITY NAME

Aiken School (OH-HA-0012)
Armco Steel (OH-BU-0001)
Arnold Creek (IN-O-0002)
Ashtree Drive and Hamilton Avenue (OH-HA-0041)
Ashtree Shopping Center (see Ashtree Drive and Hamilton Avenue, OH-HA-0041)
Aurora (IN-D-0001)

- Backbone Creek (OH-CT-0006)
 Bald Knob (OH-HA-0013)
 Batavia #1 (OH-CT-0001)
 Batavia #2 (OH-CT-0002)
 Bear Creek (OH-CT-0010)
 Beckjord (OH-CT-0019)
 Bellevue Hill (OH-HA-0003)
 "Big 4" railroad cut (see Maud cut, OH-BU-0002)
 Blue Rock Road (OH-HA-0005)
 Boat Run (OH-CT-0011)
 Bon Well Hill (IN-FR-0001)
 Boudinot Avenue (OH-HA-0042)
 Bradford (KY-BK-0001)
 Brent (KY-CP-0001)
 Briarly Creek (OH-HA-0014)
 Brookville Dam spillway (IN-FR-0002)
 Brookville North (see Garr Hill, IN-FR-0003)
- Cabin Creek (KY-LW-0001)
 Caesar Creek emergency overflow spillway (OH-WA-0001)
 Caesar Creek gorge (OH-WA-0002)
 Camp Claybanks (OH-HA-0015)
 Causeway Road (see Fairfield Causeway, IN-FR-0004)
 Chicken Hollow (OH-BR-0001)
 Chilo (OH-CT-0012)
 Christ Hospital (see Rice and Gage Streets, OH-HA-0004)
 Clifton Hill (see Bellevue Hill, OH-HA-0003)
 Congress Run (OH-HA-0016)
 Covington (KY-KE-0004)
 Cowan Lake (OH-CN-0001)
 Crosby Road (OH-HA-0017)
- Delhi Pike (OH-HA-0040)
 Devou Park (see Wayne Road, KY-KE-0006)
 Dornbusch (OH-HA-0026)
- Eagle Creek (OH-BR-0005)
 East Fork Reservoir spillway/Wm. A. Harsha Lake (OH-CT-0007)
 Eightmile Creek (OH-HA-0034)
 Elk Creek (OH-BU-0003)
 Elkhorn Creek (IN-WY-0008)
 Elkhorn Falls (IN-WY-0004)
 Emming Street (OH-HA-0002)
 Excello (see Middletown, OH-BU-0005)
- Fairfield Causeway (IN-FR-0004)
 Fairfield Road (see Fairfield Causeway, IN-FR-0004)
 Fairview Park (OH-HA-0001)
 Flat Fork Creek (OH-WA-0003)
 Fort Thomas (KY-CP-0001)
 Foster (KY-BK-0003)
- Galbraith Road (OH-HA-0032)
 Galbraith Road, cliff section (OH-HA-0033)
 Garr Hill (IN-FR-0003)
 Georgetown (OH-BR-0006)
 Goose Run (OH-BR-0007)
 Grand Avenue (OH-HA-0006) (see also Newport-Grand Avenue, KY-CP-0004)
- Hamilton Avenue (see Ashtree Drive and Hamilton Avenue, OH-HA-0041)
 Happy Hollow (OH-CT-0013)
 Harrison (OH-HA-0007)
 Harsha, Wm. A., Lake (see East Fork Reservoir spillway, OH-CT-0007)
 Hunts Creek (OH-BU-0004)
- I-75/Covington (see Covington, KY-KE-0004)
 I-275/Kentucky Route 9 (see Johns Hill, KY-CP-0005)
 I-275/Turkeyfoot Road (KY-KE-0005)
- I-275 East/Kentucky Route 445 (see Fort Thomas, KY-CP-0001)
 I-471/Grand Avenue (see Newport-Grand Avenue, KY-CP-0004)
 I-471/U.S. Route 27 (see Southgate, KY-CP-0003)
 Indiana Route 1 (see Southgate Hill, IN-FR-0005)
 Indian Creek (OH-CT-0014)
 Isaacs Creek (see Manchester, OH-AD-0003)
- Jackson Hill Park (OH-HA-0018)
 Johns Hill (KY-CP-0005)
- Kope Hollow (OH-BR-0003)
- Lawrenceburg (KY-BE-0003)
 Lick Run (OH-WA-0004)
 Light, William, Paving Company quarry (OH-CT-0005)
 Madison-Hanging Rock (IN-JE-0006)
 Madison-Indiana Route 56 (IN-JE-0004)
 Madison Middle road cut (IN-JE-0002)
 Madison North road cut (IN-JE-0001)
 Madison-Riley Creek (IN-JE-0005)
 Madison South road cut (IN-JE-0003)
 Mall Road (KY-BE-0001)
 Manchester (OH-AD-0003)
 Maud cut (OH-BU-0002)
 Maysville-Kentucky Route 11 (KY-MS-0002)
 Maysville-U.S. Routes 62 & 68 (KY-MS-0004)
 Maysville bryozoan reef mounds (KY-MS-0006)
 McMillan Street (OH-HA-0027)
 Meldahl Dam (KY-BK-0002)
 Miami Station (OH-HA-0019)
 Miamitown/Hamilton-Cleves exit (OH-HA-0009)
 Miamitown West (OH-HA-0008)
 Middlefork Reservoir (IN-WY-0007)
 Middletown (OH-BU-0005) (see also West Middletown, OH-BU-0006)
 Millvale (OH-HA-0035)
 Montana Avenue (OH-HA-0038)
 Mt. Airy Forest (OH-HA-0039)
 Muddy Creek (OH-HA-0020)
- Newport-Grand Avenue (KY-CP-0004)
 Newport Shopping Center (KY-CP-0002)
 Ninemile Creek (OH-CT-0015)
 North Bend (OH-HA-0010)
 North Point Pleasant (OH-CT-0016)
- O'Bannon Creek (OH-CT-0008)
 Ohio Brush Creek, West Fork (OH-AD-0002)
 Orphanage Road (KY-KE-0003)
- Pabco/Timberview Apartments (OH-HA-0037)
 Point Pleasant (OH-CT-0020) (see also North Point Pleasant, OH-CT-0016)
- Rabbit Hash (KY-BE-0002)
 Rapid Run (OH-HA-0036)
 Ravine Street (OH-HA-0028)
 Red Oak Creek (OH-BR-0004)
 Rice and Gage Streets (= Christ Hospital) (OH-HA-0004)
 Richmond-U.S. Route 27 (IN-WY-0001)
 Riedlin Road/Mason Road (KY-KE-0001)
 Ripley (OH-BR-0002)
 Ripley #2 (OH-BR-0008)
 Rising Sun (IN-O-0001)
 Riverside (OH-HA-0021)
 Rumpke Landfill (OH-HA-0029)
- St. Leon (IN-D-0002)
 Second Creek (OH-WA-0006)
 Sekitan Road (OH-HA-0022)
 Sharonville (OH-HA-0030)

Sheits Road (OH-HA-0023)	Turkeyfoot Road (see I-275/Turkeyfoot Road, KY-KE-0005)
Short Creek/Straight Line Pike (IN-WY-0002)	Twelvemile Creek (OH-CT-0018)
Sim Hodgins Parkway (IN-WY-0003)	
Sleepy Hollow (KY-MS-0005)	Wayne Road (KY-KE-0006)
Slickaway Run (OH-CT-0017)	Wesselman (OH-HA-0011)
Southgate (KY-CP-0003)	West Fork (OH-HA-0024)
South Gate Hill (= Indiana Route 1) (IN-FR-0005)	West Fork Ohio Brush Creek (OH-AD-0002)
South Lick Branch (KY-MS-0003)	West Harrison (IN-D-0004)
Stonelick Creek (OH-CT-0004)	West Middletown (OH-BU-0006)
Sub Run (IN-WY-0005)	West Union (OH-AD-0004)
Sugar Creek (KY-GA-0001)	Westwood-Northern Boulevard (OH-HA-0031)
	White Oak Creek (OH-BR-0009)
Tanners Creek (IN-D-0003)	White Swan Run (OH-BR-0010)
Thistlethwaite Falls (IN-WY-0006)	Wiley Branch (IN-SW-0001)
Thomas More Parkway (KY-KE-0002)	William Light Paving Company quarry (OH-CT-0005)
Timberview Apartments (see Pabco/Timberview Apartments, OH-HA-0037)	Wynbrook Apartments (OH-HA-0025)
Todd Run (OH-CT-0009)	
	445/8 road cut (see Fort Thomas, KY-CP-0001)

Addresses of authors: Richard Arnold Davis, Biology Department, College of Mount St. Joseph, 5701 Delhi Road, Cincinnati, OH 45233-1670; Sharon C. St. Louis Diekmeyer, Department of Geology, University of Cincinnati, Cincinnati, OH 45221-0013; Benjamin F. Dattilo, Geoscience Department, Weber State University, 2507 University Circle, Ogden, UT 84408-2507; Lawrence I. Goldman, MINEsoft, Ltd., 165 S. Union Blvd, #510, Lakewood, CO 80228; Steven M. Holland, Geology Department, University of Georgia, Athens, GA 30602; Roger J. Cuffey, Department of Geosciences (Deike Building), Pennsylvania State University, University Park, PA 16802.

APPENDIX B.—BIBLIOGRAPHY ON THE TYPE-CINCINNATIAN

by Richard Arnold Davis and Richard A. Spohn

INTRODUCTION

The literature relating to the type-Cincinnatian is massive and far flung. It ranges from huge monographs to casual mentions, *en passant*, in works devoted to other topics. We long ago gave up hope of compiling a truly comprehensive bibliography. We have tried to list all of the major and semi-major works, but we have not ignored minor mentions when they came our way. Certainly we have missed some works. We hope that none of these are your favorites nor those truly critical to your research.

One known source of information on the type-Cincinnatian has not been included here: the 1:24,000-scale open-file maps of the Ohio Division of Geological Survey. Reconnaissance-level bedrock-geology, bedrock-topography, and structure-contour maps are available for all of the 7.5-minute quadrangles in southwestern Ohio, and, in fact, for all of Ohio.

An especial concern has been unpublished theses and similar works. In the end we opted to include such items, but not without some discomfort. First of all, such materials are more difficult for the potential reader to obtain than are published works. Second, it is harder even to learn of the existence of such items—hence, we certainly missed many of them, so the bibliography is far from comprehensive in the coverage of unpublished material.

We have endeavored to verify every citation with the actual work cited. In some instances this has not been possible, particularly with unpublished works. Hence, we apologize, in advance, for any errors in the citations.

Some readers may find it frustrating that authors are not cited in a consistent format. We deliberately decided to try to list each author as he or she was listed in the actual publication. Thus, in some cases, the first name of an author is spelled out, whereas, in other instances, only initials are given.

Compiling a bibliography such as this one can be aided greatly by colleagues willing to share their carefully hoarded lists of references. Douglas L. Shrake, of the Ohio Division of Geological Survey, generously made available the ongoing compilation of references on the Cincinnatian that he and others at the Survey have put together over the years. Steven M. Holland, of the University of Georgia, likewise provided a list of references he compiled. Richard Fuchs, of the Cincinnati Dry Dredgers, passed on citations to references he has located in his long-term study of Cincinnati scolecodonts. Merriane Hackathorn, of the Ohio Division of Geological Survey, added some references and corrected many others, particularly adding missing information for theses and dissertations. Stig M. Bergström, of The Ohio State University, provided a list of additional references. We are grateful especially to those individuals, but also to the many others who, with the simple phrase, “Have you seen the paper . . .?”, helped make this compilation better than it might have been. It would be unfair not to give recognition to the many, many folks who have helped make the various printed comprehensive bibliographies of geology and GeoRef the extraordinarily valuable tools they are to all geologists. Hats off to them all!

BIBLIOGRAPHY

- Alberstadt, Leonard P., 1979, The brachiopod genus *Platystrophia*: U.S. Geological Survey Professional Paper 1066-B, 20 p., 7 pls.
- Alexander, Richard R., 1972, The autecology of the Cincinnatian (Upper Ordovician) brachiopod *Rafinesquina*; a biometric analysis (abs.): Geological Society of America Abstracts with Programs, v. 4, no. 5, p. 305-306.
- Alexander, Richard R., 1975, Phenotypic lability of the brachiopod *Rafinesquina alternata* (Ordovician) and its correlation with the sedimentologic regime: *Journal of Paleontology*, v. 49, no. 4, p. 607-618.
- Alexander, Richard R., 1986, Resistance to and repair of shell breakage induced by durophages in Late Ordovician brachiopods: *Journal of Paleontology*, v. 60, no. 2, p. 273-285.
- Alexander, Richard R., and Scharpf, Carl D., 1990, Epizoans on Late Ordovician brachiopods from southeastern Indiana: *Historical Biology*, v. 4, no. 3, p. 179-202.
- Ali, Syed Afaq, 1967, Kope and Fairview sedimentary structures: Brown and Adams Counties, Ohio: M.S. thesis (unpub.), The Ohio State University, 130 p.
- Ali, Syed A., and Weiss, Malcolm P., 1971, Kope and Fairview ripple marks: Brown and Adams Counties, Ohio: *Ohio Journal of Science*, v. 71, no. 3, p. 129-149.
- Amsden, Thomas W., 1983, The Late Ordovician brachiopod genera *Lepidocyclus* and *Hiscobeccus*: *Oklahoma Geological Survey Bulletin* 132, p. 36-44, pls. 5-7.
- Anstey, Robert Leland, 1970, The trepostome bryozoan fauna of the Eden Shale (Ordovician) in southeastern Indiana and adjacent areas in Kentucky and Ohio: Ph.D. dissertation (unpub.), Indiana University, 216 p. (abstract: *Dissertations Abstracts International*, v. 31, no. 4, p. 2157B).
- Anstey, Robert L., 1981, Zooid orientation structures and water flow patterns in Paleozoic bryozoan colonies: *Lethaia*, v. 14, no. 4, p. 287-302.
- Anstey, Robert L., 1986, Bryozoan provinces and patterns of generic evolution and extinction in the Late Ordovician of North America: *Lethaia*, v. 19, no. 1, p. 33-51.
- Anstey, Robert L., 1987a, Astogeny and phylogeny: evolutionary heterochrony in Paleozoic bryozoans: *Paleobiology*, v. 13, no. 1, p. 20-43.
- Anstey, R. L., 1987b, Colony patterning and functional morphology of water flow in Paleozoic stenolaemate Bryozoa, in Ross, J. R. P., ed., *Bryozoa: past and present*: Western Washington University, p. 1-8.
- Anstey, Robert L., and Bartley, John W., 1984, Quantitative stereology: an improved thin section biometry for bryozoans and other colonial organisms: *Journal of Paleontology*, v. 58, no. 3, p. 612-625.
- Anstey, Robert L., and Chase, Terry L., 1974, Geographic diversity of Late Ordovician corals and bryozoans in North America: *Journal of Paleontology*, v. 48, no. 6, p. 1141-1148.
- Anstey, Robert L., and Fowler, Michael L., 1969, Lithostratigraphy and depositional environment of the Eden Shale (Ordovician) in the tri-state area of Indiana, Kentucky, and Ohio: *Journal of Geology*, v. 77, no. 6, p. 668-682.
- Anstey, Robert L., and Pachut, Joseph F., 1980, Fourier packing ordinate: A univariate size-independent measurement of the polygonal packing variation in Paleozoic bryozoans: *Mathematical Geology*, v. 12, no. 2, p. 139-156.
- Anstey, Robert L., Pachut, Joseph F., and Prezbindowski, Dennis R., 1976, Morphogenic gradients in Paleozoic bryozoan colonies: *Paleobiology*, v. 2, no. 2, p. 131-146.
- Anstey, Robert L., and Perry, T. G., 1969, Redescription of cotypes of *Peronopora vera* Ulrich, a Cincinnatian (Late Ordovician) ectoproct species: *Journal of Paleontology*, v. 43, no. 2, p. 245-251, pls. 31-32.
- Anstey, Robert L., and Perry, T. G., 1972 (1973), Eden Shale bryo-

- zoans: a numerical study (Ordovician, Ohio Valley): Michigan State University, Publications of the Museum, Paleontological Series, v. 1, no. 1, 80 p., 22 pls. (title page bears the date 1972, but the verso of the title page states "Published 14 January 1973").
- Anstey, Robert L., and Rabbio, S. Frank, 1989, Regional bryozoan biostratigraphy and taphonomy of the Edenian stratotype (Kope Formation, Cincinnati area): graphic correlation and gradient analysis: *Palaios*, v. 4, no. 6, p. 574-584.
- Anstey, R. L., Rabbio, S. F., and Tuckey, M. E., 1987, Bryozoan bathymetric gradients within a Late Ordovician epeiric sea: *Paleoceanography*, v. 2, no. 2, p. 165-176.
- Anstey, Robert L., and Wilson, Mark A., 1996, Phylum Bryozoa, in Feldmann, Rodney M., and Hackathorn, Merrienne, eds., *Fossils of Ohio: Ohio Division of Geological Survey Bulletin 70*, p. 196-209.
- Anthony, John G., 1838, New trilobite, *Ceratocephala ceralepta*: *American Journal of Science*, v. 34, no. 2, p. 379-380.
- Anthony, John G., 1839a, Fossil encrinite: *American Journal of Science*, v. 35, no. 2, p. 359-360.
- Anthony, John G., 1839b, Description of a new fossil (*Calymene bucklandii*): *American Journal of Science*, v. 36, no. 1, p. 106-107.
- Anthony, John Gould, 1847, On an impression of the soft parts of an *Orthoceras* [Cincinnati, Ohio]: *Quarterly Journal of the Geological Society of London*, v. 3, p. 255-257, 1 illustration.
- Anthony, John G., 1848, On an impression of the soft parts of an *Orthoceras*: *American Journal of Science*, ser. 2, v. 6, no. 16, p. 132-133, 1 illustration (this actually is a discussion of the previous paper; author is not given).
- Aronoff, Steven Martin, 1973, A reexamination of the type material of some Upper Ordovician nautiloid cephalopods: *Orthocerida and Actinocerida of the Cincinnati, Ohio area*: M.S. thesis (unpub.), University of Cincinnati, 79 p., 11 pls.
- Aronoff, Steven M., 1979, Orthoconic nautiloid morphology and the case of *Treptoceras* vs. *Orthonybyoceras*: *Neues Jahrbuch für Geologie und Paläontologie, Abhandlungen*, v. 158, no. 1, p. 100-122.
- Aronoff, Steven Martin, 1981, Phenon and interspecific variation in selected Paleozoic orthoconic nautiloids (Mollusca, Cephalopoda): Ph.D. dissertation (unpub.), University of Iowa, 382 p.
- Aronoff, Steven M., 1983, The taxonomy of smooth orthoconic nautiloid cephalopods; a reappraisal (abs.): *Geological Society of America Abstracts with Programs*, v. 15, no. 6, p. 517.
- Association of Geology Teachers. See [National] Association of Geology Teachers [East-Central Section], 1958.
- Astrova, G. G., 1973, Polymorphism and its development in the trepostomatous Bryozoa, in Larwood, G. P., ed., *Living and fossil Bryozoa*: London and New York, Academic Press, p. 1-10, pls. 1-2.
- Atkins, Frank Pearce, Jr., 1940, The geology of the Higginsport quadrangle (north of the Ohio River): M.A. thesis (unpub.), University of Cincinnati, 69 p.
- Ausich, William I., 1981, The regional paleontology and stratigraphy of Ohio: Wright State University Department of Geology Technical Report PA8103WA, 100 p.
- Ausich, William I., 1996, Phylum Echinodermata, in Feldmann, Rodney M., and Hackathorn, Merrienne, eds., *Fossils of Ohio: Ohio Division of Geological Survey Bulletin 70*, p. 242-261.
- Austin, George M., 1927, Richmond fossil zones in Warren and Clinton Counties, Ohio: *Proceedings of the U.S. National Museum*, v. 70, art. 22 (no. 2671), 18 p.
- Axon, A. G., 1986, Paleocology of a Cincinnati (Ordovician) crinoid garden from southwestern Ohio (abs.): *Geological Society of America Abstracts with Programs*, v. 18, no. 4, p. 278.
- Axon, Allan G., 1987, Preliminary paleoecologic study of a Maysvillian (Cincinnati) crinoid garden: M.S. thesis (unpub.), University of Kentucky, 145 p.
- Axon, Allan G., 1992, Ordovician crinoid paleoecology and community paleoecology: Ph.D. dissertation (unpub.), University of Kentucky, 416 p.
- Babcock, Loren E., 1996a, Phylum Porifera, stromatoporoids, in Feldmann, Rodney M., and Hackathorn, Merrienne, eds., *Fossils of Ohio: Ohio Division of Geological Survey Bulletin 70*, p. 64-65.
- Babcock, Loren E., 1996b, Phylum Conulariida, in Feldmann, Rodney M., and Hackathorn, Merrienne, eds., *Fossils of Ohio: Ohio Division of Geological Survey Bulletin 70*, p. 66-69.
- Babcock, Loren E., 1996c, Phylum Cnidaria, in Feldmann, Rodney M., and Hackathorn, Merrienne, eds., *Fossils of Ohio: Ohio Division of Geological Survey Bulletin 70*, p. 70-89.
- Babcock, Loren E., 1996d, Phylum Arthropoda, Class Trilobita, in Feldmann, Rodney M., and Hackathorn, Merrienne, eds., *Fossils of Ohio: Ohio Division of Geological Survey Bulletin 70*, p. 90-113.
- Baird, Gordon C., Brett, Carleton E., and Frey, Robert C., 1989, "Hitchhiking" epizoans on orthoconic cephalopods: preliminary review of the evidence and its implications: *Senckenbergiana Lethaea*, v. 69, no. 5/6, p. 439-465, 4 pls.
- Baptista, Braulio M., 1969, Petrology of limestones of the Arnheim Formation (Cincinnati Series) from western Butler County, Ohio: M.S. thesis (unpub.), Miami University, 93 p., 1 pl.
- Barbour, George B., Bernhagen, Ralph J., Caster, Kenneth E., Durrell, Richard H., Koucky, Frank L., and Lewis, Robert C., 1961, Geology of the Cincinnati region: Guide to the Thirty-sixth Annual Field Conference of the Ohio Academy of Science, April 22 & 23, 1961, 21 p., 6 pls.
- Barnes, C. R., 1992, The uppermost series of the Ordovician System: in Webby, B. D., and Laurie, J. R., eds., *Global perspectives on Ordovician geology: Proceedings of the Sixth International Symposium on the Ordovician System*, University of Sydney, Australia, 15-19 July 1991, Rotterdam, The Netherlands, A. A. Balkema, p. 185-192.
- Barnes, C. R., and Bergström, S. M., 1988, Conodont biostratigraphy of the uppermost Ordovician and lowermost Silurian: *Bulletin of the British Museum (Natural History), Geology*, v. 43, p. 325-343.
- Bartley, John W., and Anstey, Robert L., 1983, Cyclic growth in lower Paleozoic stenolaemate bryozoans (abs.): *Geological Society of America Abstracts with Programs*, v. 15, no. 6, p. 523.
- Bartley, John W., Anstey, Robert L., and Brunner, Carol M., 1982, Nested growth rhythms in Paleozoic bryozoans (abs.): *Geological Society of America Abstracts with Programs*, v. 14, no. 5, p. 254.
- Bassarab, D. R., and Huff, W. D., 1969, Clay mineralogy of Kope and Fairview Formations (Cincinnati) in the Cincinnati area: *Journal of Sedimentary Petrology*, v. 39, no. 3, p. 1014-1022.
- Bassler, R. S., 1903, The structural features of the bryozoan genus *Homotrypa*, with descriptions of species from the Cincinnati Group: *Proceedings of the U.S. National Museum*, v. 26, no. 1323, p. 565-591, pls. 20-25.
- Bassler, Ray S., 1906, A study of the James types of Ordovician and Silurian Bryozoa: *Proceedings of the U.S. National Museum*, v. 30, no. 1442, p. 1-66, pls. 1-6.
- Bassler, R. S., 1913, *Bryozoa*, in Eastman, Charles P., ed., *Zittel's Text-book of paleontology* (2nd ed.): London, Macmillan and Co., v. 1, p. 314-355.
- Bassler, Ray S., 1915, Bibliographic index of American Ordovician and Silurian fossils: *U.S. National Museum Bulletin 92*, 2 v., 1,521 p., 4 pls.
- Bassler, Ray S., 1953, Part G, *Bryozoa*, in Moore, Raymond C., ed., *Treatise on Invertebrate Paleontology: Geological Society of America and University of Kansas Press*, 253 p.
- Bassler, R. S., and Duncan, H., 1955, Proposed use of the plenary powers to standardize the current use of "*Monticulipora*" D'Orbigny, 1849 (Class Bryozoa, Order Cyclostomata or Trepostomata, Family Monticuliporidae Nicholson): *Bulletin of Zoological Nomenclature*, v. 11, pt. 3, p. 90-92.
- Bather, F. A., 1920, [Review of McEwan, Eula Davis, 1919, A study of the brachiopod genus *Platystrophia*]: *Geological Magazine*, v. 57, p. 88-90.
- Bayha, David C., 1965, The petrology of selected limestones of the

- McMillan Formation from the Trenton quadrangle, Butler County, Ohio: M.S. thesis (unpub.), Miami University, 128 p., 25 pls.
- Becker, Karl, 1938, Cincinnati area, the Mother of Geologists: *Compass*, v. 19, no. 1, p. 188-196.
- Bell, Bruce McConnell, 1966, An introduction to the Cincinnati edrioasteroids (Echinodermata): M.S. thesis (unpub.), University of Cincinnati, 168 p., 2 pls.
- Bell, Bruce McConnell, 1972, A study of North American Edrioasteroidea: Ph.D. dissertation (unpub.), University of Cincinnati, 2 v., 1,022 p., 63 pls.
- Bell, Bruce M., 1976, A study of North American Edrioasteroidea: *New York State Museum and Science Service Memoir* 21, 447 p.
- Bell, Bruce M., 1979, Edrioasteroids (Echinodermata): U.S. Geological Survey Professional Paper 1066-E, 7 p., 2 pls.
- Bell, Thomas J., 1882, History of the water supply of the world, arranged in a comprehensive form from eminent authorities, containing a description of the various methods of water supply, pollution and purification of waters, and sanitary effects, with analysis of potable waters, also geology and water strata of Hamilton County, Ohio, statistics of the Ohio River, proposed water supply of Cincinnati: Cincinnati, Peter G. Thompson, 134 p.
- Berdan, Jean M., 1984, Leperditicopid ostracodes from Ordovician rocks of Kentucky and nearby states and characteristic features of the order Leperditicopida: U.S. Geological Survey Professional Paper 1066-J, 40 p., 11 pls.
- Bergstrand, P. M., Colby, N. D., Hodnett, M. L., Lasemi, Z., and Stone, G. B., 1989, Limestone nodules from the Oregonia and Elk Creek beds, Cincinnati Series, southwestern Ohio (abs.): *Ohio Journal of Science*, v. 89, no. 2, p. 9.
- Bergström, Stig M., 1971, Conodont biostratigraphy of the Middle and Upper Ordovician of Europe and eastern North America, in Sweet, Walter C., and Bergström, Stig M., eds., *Symposium on conodont biostratigraphy*: Geological Society of America Memoir 127, p. 83-157., 2 pls.
- Bergström, Stig M., 1980, Relations between North American, Baltic, and British standard series and stages as suggested by conodont and graptolite evidence (abs.): 26th International Geological Congress, Paris, 1980, Abstracts, v. 1, p. 204.
- Bergström, Stig M., 1982, Biostratigraphic relations between Ordovician conodont and graptolite zones (abs.): Geological Society of America Programs with Abstracts, v. 14, no. 5, p. 255.
- Bergström, Stig M., 1986, Biostratigraphic integration of Ordovician graptolite and conodont zones—a regional review, in Hughes, C. P., and Rickards, R. B., eds., *Palaeoecology and biostratigraphy of graptolites*: Geological Society Special Publication 20, p. 61-78.
- Bergström, Stig M., 1990, Relations between conodont provincialism and the changing palaeogeography during the early Palaeozoic: Geological Society Memoir 12, p. 105-121.
- Bergström, Stig M., 1991, The Sebree Trough Project. 3. Biostratigraphic and biogeographic significance of graptolites from the Ordovician Utica Shale in the subsurface of Ohio and Indiana (abs.): Geological Society of America Abstracts with Programs, v. 23, no. 3, p. 4.
- Bergström, Stig M., 1996a, Phylum Hemichordata, Class Graptolithina, in Feldmann, Rodney M., and Hackathorn, Merrienne, eds., *Fossils of Ohio*: Ohio Division of Geological Survey Bulletin 70, p. 270-281.
- Bergström, Stig M., 1996b, Tentaculitoids, in Feldmann, Rodney M., and Hackathorn, Merrienne, eds., *Fossils of Ohio*: Ohio Division of Geological Survey Bulletin 70, p. 282-287.
- Bergström, Stig M., 1997, The oldtimers were right: new data on the relations between the type Cincinnati and the Upper Ordovician in the Upper Mississippi Valley, Oklahoma, Texas, New Mexico, and the Western Interior (abs.): Geological Society of America Abstracts with Programs, v. 29, no. 4, p. 5.
- Bergström, Stig M., 1998, The base of the *Amorphognathus ordovicicus* Zone as a potential global stage boundary in a revised chronostratigraphic classification of the Ordovician System (abs.): Geological Society of America Abstracts with Programs, v. 30, no. 7, p. A-337.
- Bergström, S. M., and Boucot, A. J., 1988, The Ordovician-Silurian boundary in the United States: *Bulletin of the British Museum (Natural History), Geology*, v. 43, p. 273-284.
- Bergström, Stig M., Carnes, John B., Ethington, Raymond L., Votaw, Robert B., and Wigley, Perry B., 1974, *Appalachignathus*, a new multielement conodont genus from the Middle Ordovician of North America: *Journal of Paleontology*, v. 48, no. 2, p. 227-235, 1 pl.
- Bergström, Stig M., Evans, Robert D., Knabe, Keith A., and Miller, Merrell A., 1981, Middle and Upper Ordovician chitinozoan and graptolite biostratigraphy in the Cincinnati region (Ohio, Kentucky, Indiana) (abs.): Geological Society of America Abstracts with Programs, v. 13, no. 7, p. 438.
- Bergström, Stig M., Kolata, Dennis, R., and Huff, Warren D., 1991, Geologic significance of gigantic ash falls in the Ordovician of North America and northwestern Europe (abs.): *Ohio Journal of Science*, v. 91, no. 2, p. 34.
- Bergström, Stig M., and Mitchell, Charles E., 1985, Occurrence, morphology, and biostratigraphic significance of graptolites in the Cincinnati Series, the reference standard of the North American Upper Ordovician (abs.): Graptolite Working Group of the International Palaeontological Association, Third International Conference, Copenhagen, Denmark, 1985, Abstracts, p. 3.
- Bergström, Stig M., and Mitchell, Charles E., 1986, The graptolite correlation of the North American Upper Ordovician Standard: *Lethaia*, v. 19, no. 3, p. 247-266.
- Bergström, Stig M., and Mitchell, Charles E., 1987, Geology of Middle to Upper Ordovician Sebree Trough northwest of Cincinnati Arch, Ohio, Indiana, and Kentucky (abs.): *American Association of Petroleum Geologists Bulletin*, v. 71, p. 1101.
- Bergström, Stig M., and Mitchell, Charles E., 1989, Relations between the Ordovician Trenton Group and associated strata in the Northern Appalachian foreland basin-shelf region and their lateral equivalents in the eastern Midcontinent: *Appalachian Basin Industrial Associates, Program for Spring Meeting*, v. 15, p. 7-41.
- Bergström, Stig M., and Mitchell, Charles E., 1990a, Trans-Pacific graptolite faunal relations: the biostratigraphic position of the base of the Cincinnati Series (Upper Ordovician) in the standard Australian graptolite zone succession: *Journal of Paleontology*, v. 64, no. 6, p. 992-997.
- Bergström, Stig M., and Mitchell, Charles E., 1990b, The Utica Shale of northern Ohio and its relationship to the Utica Shale of the northern Appalachian Basin and lithologically similar rocks in the central Great Lakes region: *Appalachian Basin Industrial Associates, Program for Fall Meeting*, v. 17, p. 2-39.
- Bergström, S. M., and Mitchell, C. E., 1992, The Ordovician Utica Shale in the eastern Midcontinent region: age, lithofacies, and regional relationships: *Oklahoma Geological Survey Bulletin* 145, p. 67-89.
- Bergström, Stig M., and Mitchell, Charles E., 1994, Regional relationships between late Middle and early Late Ordovician standard successions in New York and Quebec and in the Cincinnati region in Ohio, Indiana, and Kentucky: *New York State Museum Bulletin* 481, p. 5-20.
- Bergström, Stig M., Mitchell, Charles E., and Goldman, D., 1989, Relations between the Mohawkian and Cincinnati North American Standard series: new biostratigraphic data from the type areas in the Cincinnati region and New York State (abs.): Geological Society of America Abstracts with Programs, v. 21, no. 4, p. 4.
- Bergström, Stig M., Schumacher, Gregory A., and Mitchell, Charles E., 1991, Recent scientific core drilling in western Ohio: significance of the Oxford and Elkhorn cores for the interpretation and nature of the Ordovician Sebree Trough (abs.): *Ohio Journal of Science*, v. 91, no. 2, p. 30.
- Bergström, Stig M., and Sweet, W. C., 1966, Conodonts from the Lexington Limestone (Middle Ordovician) of Kentucky and its lateral equivalents in Ohio and Indiana: *Bulletins of American*

- Paleontology, v. 50, no. 229, p. 271-441, pls. 28-35.
- Bergström, Stig M., and Sweet, W. C., 1969, Ordovician conodont localities, southwestern Ohio: Field trip guidebook, 3rd annual meeting, North-Central Section, Geological Society of America, Ohio Division of Geological Survey, p. 3-1 to 3-13.
- Berry, William B. N., 1966, *Orthograptus truncatus richmondensis* (Ruedemann) in the Arnheim Formation (Ordovician) in Indiana: *Journal of Paleontology*, v. 40, no. 6, p. 1392-1394.
- Blackwell, W. H., 1985, Evidence of algal source of micrite in a Saluda coral zone in southeastern Indiana: *Proceedings of the Indiana Academy of Science for 1984*, v. 94, p. 391-394.
- Blackwell, Will H., Marak, Joe H., and Powell, Martha J., 1982, The identity and reproductive structures of a misplaced *Solenopora* (Rhodophycophyta) from the Ordovician of southwestern Ohio and eastern Indiana: *Journal of Phycology*, v. 18, p. 477-482.
- Blackwell, Will H., Marak, Joe H., Powell, Martha J., and Martin, Wayne D., 1984, *Girvanella* (Cyanochloronta) in an Ordovician oncolite zone in eastern Indiana: *Bulletin of the Torrey Botanical Club*, v. 111, no. 2, p. 165-170.
- Blackwell, W. H., and Martin, A. J., 1986, Bacterially-like (filamentous) structures associated with pyritized burrow linings, Arnheim Formation (Upper Ordovician), southeastern Indiana (abs.): *American Association of Petroleum Geologists Bulletin*, v. 70, no. 5, p. 566.
- Blake, D. B., 1975, The order Cryptostomata resurrected, in David, L., ed., *Bryozoan 1974*, Documents Laboratoires Geologie Faculté des Sciences Lyon, H. S. 3, fasc. 1, p. 211-223, 3 pls.
- Blake, D. B., 1979, The Arthrostylidae and articulated growth habits in Paleozoic bryozoans, in Larwood, G. P., and Abbott, M. B., eds., *Advances in Bryozoology*: London, Academic Press, p. 337-344.
- Blake, Daniel B., 1990, Paleobiological implications of some Upper Ordovician juvenile asteroids (Echinodermata): *Lethaia*, v. 23, no. 4, p. 347-357.
- Blake, Daniel B., and Guensburg, Thomas E., 1988, The water vascular system and functional morphology of Paleozoic asteroids: *Lethaia*, v. 21, no. 3, p. 189-206.
- Blake, D. B., and Guensburg, T. E., 1992, Caught in the act: a Late Ordovician asteroid and its pelecypod prey (abs.): *Geological Society of America Abstracts with Programs*, v. 24, no. 4, p. 6.
- Blake, Daniel B., and Guensburg, Thomas E., 1994, Predation by the Ordovician asteroid *Promopalaeaster* on a pelecypod: *Lethaia*, v. 27, no. 3, p. 235-239.
- Bliss, Franklin E., 1984, The "concrete layer" (a unit resembling the Hitz Bed) of the Saluda Formation (Upper Ordovician) in Franklin County, Indiana: M.S. thesis (unpub.), Miami University, 121 p.
- Boardman, Richard S., 1960, A revision of the Ordovician bryozoan genera *Batostoma*, *Anaphragma*, and *Amplexopora*: *Smithsonian Miscellaneous Collections*, v. 140, no. 5, 28 p., 7 pls.
- Boardman, R. S., 1968, Colony development and convergent evolution of budding pattern in "rhombotrypid" Bryozoa: *Atti della Società di Scienze Naturali e del Museo Civico di Storia Naturale di Milano*, v. 108, p. 179-184.
- Boardman, Richard S., 1971, Mode of growth and functional morphology of autozooids in some Recent and Paleozoic tubular Bryozoa: *Smithsonian Contributions to Paleobiology*, no. 8, 51 p., 11 pls.
- Boardman, R. S., Cheetham, A. H., Blake, D. B., Utgaard, John, Karlkins, O. L., Cook, P. L., Sandberg, P. A., Lutaud, Genevieve, and Wood, T. S., 1983, *Bryozoa*, Part G, revised, volume 1: Introduction, Order Cystoporata, Order Cryptostomata, in Robison, R. A., ed., *Treatise on Invertebrate Paleontology*: Geological Society of America and University of Kansas Press, 625 p.
- Boardman, Richard S., and Cheetham, Alan H., 1969, Skeletal growth, intracolony variation, and evolution in Bryozoa: a review: *Journal of Paleontology*, v. 43, no. 2, p. 205-233, pls. 27-30.
- Boardman, R. S., Cheetham, A. H., and Cook, P. L., 1970, Intracolony variation and the genus concept in Bryozoa: *North American Paleontological Convention, Chicago, 1969, Proceedings C*, p. 294-320.
- Boardman, Richard S., and McKinney, Frank K., 1976, Skeletal architecture and preserved organs of four-sided zooids in convergent genera of Paleozoic Trepostomata (Bryozoa): *Journal of Paleontology*, v. 50, no. 1, p. 25-78, 16 pls.
- Boardman, Richard S., and Utgaard, John, 1964, Modifications of study methods for Paleozoic Bryozoa: *Journal of Paleontology*, v. 38, no. 4, p. 768-770.
- Boardman, Richard S., and Utgaard, John, 1966, A revision of the Ordovician bryozoan genera *Monticulipora*, *Peronopora*, *Heterotrypa*, and *Dekayia*: *Journal of Paleontology*, v. 40, no. 5, p. 1082-1108, pl. 133-142.
- Bodenbender, Brian E., Wilson, Mark A., and Palmer, Timothy J., 1989, Paleocology of *Sphenothallus* on an Upper Ordovician hardground: *Lethaia*, v. 22, no. 2, p. 217-225.
- Bolton, T. E., 1988, Stromatoporoida from the Ordovician rocks of central and eastern Canada: *Geological Survey of Canada Bulletin* 379, p. 17-45.
- Bolton, T. E., and Ross, J. R. P., 1985, The cryptostomate bryozoan *Sceptropora* (Rhabdomesina, Arthrostylidae) from Upper Ordovician rocks of southern Mackenzie Mountains, District of Mackenzie: *Geological Survey of Canada Paper* 85-1A, p. 29-45, pls. 5.1-5.7.
- Booth, James S., and Osborne, Robert H., 1971, The American Upper Ordovician Standard. XV. Clay mineralogy of insoluble residues from Cincinnati limestones, Hamilton County, Ohio: *Journal of Sedimentary Petrology*, v. 41, no. 3, p. 840-843.
- Borella, Peter E., and Osborne, Robert H., 1978, Late Middle and early Late Ordovician history of the Cincinnati Arch province, central Kentucky to central Tennessee: *Geological Society of America Bulletin*, v. 89, no. 10, p. 1559-1573.
- Bowsher, A. L., 1955, Origin and adaptation of platyceratid gastropods: *University of Kansas Paleontological Contributions, Mollusca*, art. 5, p. 1-11.
- Bradshaw, Lael E., 1989, E. O. Ulrich and the development of American stratigraphy: Ph.D. dissertation (unpub.), University of Cincinnati (Department of History), 329 p.
- Brandt, Danita S., 1980a, Paleocology of an Upper Ordovician "trilobite shale" (abs.): *Geological Society of America Abstracts with Programs*, v. 12, no. 5, p. 220.
- Brandt, Danita S., 1980b, Phenotypic variation and paleocology of *Flexicalymene* (Arthropoda: Trilobita) in the Cincinnati Series (Upper Ordovician) near Cincinnati, Ohio: M.S. thesis (unpub.), University of Cincinnati, 148 p., 4 pls.
- Brandt Velbel, Danita, 1985, Ichnologic, taphonomic, and sedimentologic clues to the deposition of Cincinnati shales (Upper Ordovician), Ohio, in Curran, H. Allen, ed., *Biogenic structures: their use in interpreting depositional environment*: *Society of Economic Paleontologists and Mineralogists Special Publication* 35, p. 299-307.
- Brandt, Danita S., 1986, Preservation of event beds through time: *Palaios*, v. 1, no. 1, p. 92-96.
- Brandt, Danita S., 1991, Aspects of *Flexicalymene* (Trilobita) paleobiology (abs.): *Geological Society of America Abstracts with Programs*, v. 23, no. 3, p. 4.
- Brandt, Danita S., 1993, Ecdysis in *Flexicalymene meeki* (Trilobita): *Journal of Paleontology*, v. 67, no. 6, p. 999-1005.
- Brandt, Danita S., 1996, Epizoans on *Flexicalymene* (Trilobita) and implications for trilobite paleocology: *Journal of Paleontology*, v. 70, no. 3, p. 442-449.
- Brandt, Danita S., Meyer, David L., and Lask, Peter B., 1995, *Isotelus* (Trilobita) "hunting burrow" from Upper Ordovician strata, Ohio: *Journal of Paleontology*, v. 69, no. 6, p. 1079-1083.
- Branson, E. B., Mehl, M. G., and Branson, C. C., 1951, Richmond conodonts of Kentucky and Indiana: *Journal of Paleontology*, v. 25, no. 1, p. 1-17, pls. 1-4.
- Branstrator, J. Wayne, 1972, *Lanthanaster cruciformis*, a new Upper Ordovician sea star from Cincinnati, Ohio: *Journal of Paleontology*, v. 46, no. 1, p. 66-69, 1 pl.
- Branstrator, Jon W., 1975a, Paleobiology and revision of the Ordovician Asteriadiina (Echinodermata: Asteroidea) of the Cincinnati area: Ph.D. dissertation (unpub.), University of Cincinnati, 269 p., 18 pls.

- Branstrator, J. Wayne, 1975b, Podial efficacy of some Ordovician asteroids (Echinodermata) of North America: *Bulletins of American Paleontology* v. 67, no. 287, p. 57-69, 1 pl.
- Branstrator, J. W., 1979, Asteroidea (Echinodermata): U.S. Geological Survey Professional Paper 1066-F, 7 p., 3 pls.
- Braun, E. Lucy, 1912, The Cincinnati Series and its brachiopods in the vicinity of Cincinnati: Master's thesis (unpub.), University of Cincinnati, 48 p.
- Braun, E. Lucy, 1916, The Cincinnati Series and its brachiopods in the vicinity of Cincinnati: *Journal of the Cincinnati Society of Natural History*, v. 22, no. 1, p. 18-44, + fold-out chart.
- Bretsky, Peter, 1970, Late Ordovician benthic marine communities in north-central New York: *New York State Museum and Science Service Bulletin* 414, 34 p.
- Briggs, C., Jr., 1838, Report of Mr. Briggs: Geological Survey of Ohio Second Annual Report, p. 109-154, + 1 section.
- Brower, James C., 1995, Dendrocrinid crinoids from the Ordovician of northern Iowa and southern Minnesota: *Journal of Paleontology*, v. 69, no. 5, p. 939-960.
- Brown, George D., Jr., and Anstey, Robert L., 1968, Lexington Limestone-Kope Formation contact (Ordovician) in southwestern Indiana: *American Association of Petroleum Geologists Bulletin*, v. 52, no. 3, p. 488-493.
- Brown, George D., Jr., and Daly, Edward J., 1981, Evolutionary trends in *Parvohallopora* Singh in the Dillsboro Formation (Upper Ordovician) of southeastern Indiana, in Larwood, Gilbert P., and Nielsen, Claus, eds., Recent and fossil Bryozoa: 5th International Conference of the International Bryozoology Association, Durham, U.K., September 1980: University of Durham, Department of Geological Sciences, p. 29-38.
- Brown, George D., Jr., and Daly, Edward J., 1985a, Analysis of evolution of structural characters of *Parvohallopora* Singh from the Dillsboro Formation, Cincinnati Series, Indiana, in Nielsen, Claus, and Larwood, Gilbert P., eds., Bryozoa, Ordovician to Recent: papers presented at the 6th International Conference on Bryozoa, Vienna, 1983: Fredensborg, Denmark, Olsen and Olsen, p. 51-58.
- Brown, George D., Jr., and Daly, Edward J., 1985b, Trepastome Bryozoa from the Dillsboro Formation (Cincinnati Series) of southeastern Indiana: *Indiana Geological Survey Special Report* 33, 95 p., 12 pls.
- Brown, George D., Jr., and Daly, Edward J., 1987, Trepastome bryozoan distributions in the Dillsboro Formation (Cincinnati Series), Indiana: water depth controls?, in Ross, June R. P., ed., *Bryozoa: present and past*: Western Washington University, p. 41-47.
- Brown, George D., Jr., and Lineback, Jerry A., 1966, Lithostratigraphy of the Cincinnati Series (Upper Ordovician) in southeastern Indiana: *American Association of Petroleum Geologists Bulletin*, v. 50, no. 5, p. 1018-1023.
- Browne, Ruth G., 1964, The coral horizons and stratigraphy of the upper Richmond Group in Kentucky west of the Cincinnati Arch: *Journal of Paleontology*, v. 38, no. 2, p. 385-392.
- Browne, Ruth G., 1965, Some upper Cincinnati (Ordovician) colonial corals of north-central Kentucky: *Journal of Paleontology*, v. 39, no. 6, p. 1177-1191, pls. 146-152.
- Brownstein, Jack M., 1960, Analysis of the non-clay fractions of the mudstones from a lower Cincinnati Series core, Butler County, Ohio: M.S. thesis (unpub.), Miami University, 53 p.
- Bruns, Richard Harte, 1953, The role of storm winnowing in producing the characteristic stratification of the Cincinnati rocks in the Cincinnati region: M.S. thesis (unpub.), University of Cincinnati, 45 p.
- Bucher, Walter H., 1917, Large current-ripples as indicators of paleogeography: *Proceedings of the U.S. National Academy of Sciences*, v. 3, p. 285-291.
- Bucher, W. H., 1919, On ripples and related sedimentary surface forms and their indicators of paleogeographic interpretation: *American Journal of Science*, ser. 4, v. 47, no. 279, p. 149-210; no. 280, p. 241-269.
- Bucher, W. H., 1938, A shell-boring gastropod in a Dalmanella bed of upper Cincinnati age: *American Journal of Science*, ser. 5, v. 36, no. 211, p. 1-7 [note: Dalmanella was not italicized in original].
- Bucher, Walter H., Caster, Kenneth E., and Jones, Stewart, 1939, Elementary description of Cincinnati fossils and strata and plates of commoner fossils in the vicinity of Cincinnati, Ohio: University of Cincinnati, 13 p., 8 pls.
- Bucher, Walter H., Caster, Kenneth E., and Jones, Stewart M., 1945, Elementary guide to the fossils and strata in the vicinity of Cincinnati, Ohio: Cincinnati Museum of Natural History, 31 p.
- Butler, John C., and Scotford, David M., 1973, Re-examination of upper Cincinnati shale petrology: *Geological Society of America Bulletin*, v. 84, no. 7, p. 2431-2434.
- Butler, Katharine L., and Cuffey, Roger J., 1994, Bryodiversity and paleostress in the Saluda dolomitic shelf lagoon (latest Ordovician, southeastern Indiana) (abs.): *Geological Society of America Abstracts with Programs*, v. 26, no. 5, p. 7.
- Butler, Katharine L., and Cuffey, Roger J., 1996, Reduced bryozoan diversity and paleoenvironmental stress in the Saluda Dolomite (uppermost Ordovician, southeastern Indiana), in Gordon, Dennis P., Smith, Abigail M., and Grant-Mackie, Jack A., eds., *Bryozoans in space and time*: Wellington, New Zealand, Proceedings of the 10th International Bryozoology Conference (1995), National Institute of Water & Atmospheric Research, Ltd., p. 55-61.
- Butler, Robert E., 1954, Paleontology and stratigraphy of the Cynthiana Formation: M.S. thesis (unpub.), Miami University, 59 p.
- Calvert, Warren L., Noger, Martin C., and Branson, E. R., 1968, Geological aspects of the Maysville-Portsmouth region, southern Ohio and northeastern Kentucky: *Guidebook for the Ohio Geological Society-Geological Society of Kentucky Joint Field Conference*, May 17-18, 1968, Kentucky Geological Survey, 86 p., 1 pl.
- Carpenter, Gene C., 1965, The lower dolomite member of the Ordovician Chazy Limestone and the St. Peter Sandstone of north-central Kentucky and southwestern Ohio: *Ohio Journal of Science*, v. 65, no. 2, p. 85-94.
- Carpenter, James W., and Ory, Thomas R., 1961, The American Upper Ordovician Standard. VI. The Covington Sequence at Maysville, Kentucky: *Ohio Journal of Science*, v. 61, no. 6, p. 372-378.
- Carriker, Melbourne R., and Yochelson, Ellis L., 1968, Recent gastropod boreholes and Ordovician cylindrical borings: U.S. Geological Survey Professional Paper 593-B, 26 p., 5 pls.
- Caster, Kenneth E., 1942, Two siphonophores from the Paleozoic: *Palaeontographica Americana*, v. 3, no. 14, 34 p., 2 pls.
- Caster, Kenneth E., 1952, Concerning *Enoploura* of the Upper Ordovician and its relation to other carpoid echinoderms: *Bulletins of American Paleontology*, v. 34, no. 141, 56 p., 4 pls.
- Caster, Kenneth E., 1961, Sunday all-day field excursion in the Cincinnati region, field trip 7: *Guidebook for field trips, Cincinnati meeting*, 1961, Geological Society of America, p. 253-257.
- Caster, Kenneth E., 1981, Cincinnati contributions to knowledge of the Lophora, in Broadhead, T. W., ed., *Lophophorates*, notes for a short course: University of Tennessee, Department of Geological Sciences, *Studies in Geology* 5, p. 237-251.
- Caster, Kenneth E., 1982, The Cincinnati "School" of paleontology: *Earth Sciences History*, v. 1, no. 1, p. 23-28.
- Caster, Kenneth E., Dalvé, Elizabeth A., and Pope, John K., 1955, Elementary guide to the fossils and strata of the Ordovician in the vicinity of Cincinnati, Ohio: Cincinnati Museum of Natural History, 47 p., 8 pls. [commonly cited as 1961, from a reprint dated that year; the original printing has a yellow cover; the first reprint has a gray cover; subsequent reprintings each have a blue cover].
- Caster, Kenneth E., and Kjellesvig-Waering, Erik N., 1964, Upper Ordovician eurypterids of Ohio: *Palaeontographica Americana*, v. 4, no. 32, p. 301-358, pls. 43-53.
- Caster, Kenneth E., and Macke, William B., 1952, An aglaspid merostome from the Upper Ordovician of Ohio: *Journal of Paleontology*, v. 26, no. 5, p. 753-757, pl. 109.

- Caster, K. E., Durrell, R. H., and Jenks, W. F., 1970, Cincinnati strata from Oregonia to the Ohio River, with notes on Pleistocene geology along the route (Warren and Clermont Counties): Guide to the forty-fifth annual field conference of the Section of Geology of the Ohio Academy of Science, April 18, 1970, 19 p., 9 figs.
- Chappars, Michael Stephen, 1935, Stratigraphy of Cincinnati and vicinity: *Compass*, v. 15, no. 3, p. 135-142.
- Chappars, Michael Stephen, 1936, Catalog of the type specimens of fossils in the University of Cincinnati Museum: *Ohio Journal of Science*, v. 36, no. 1, p. 1-45.
- Chimney, Peter J., 1977, Paleoecology of the trepostome ectoprocts of the Richmond Group (Upper Ordovician) southwest Ohio-southeast Indiana: M.S. thesis (unpub.), Miami University, 147 p.
- Clarke, J. M., and Schuchert, Charles, 1899, The nomenclature of the New York series of geological formations: *Science*, new ser., v. 10, p. 874-878.
- Committee on Geological Nomenclature, 1879. See Miller, S. A., and others, 1879.
- Conklin, B. J., 1977, Stratigraphy and petrography of the Madison, Indiana, U.S. 421 road cut, Upper Ordovician through Middle Silurian: B.S. thesis (unpub.), Earlham College, 75 p.
- Conrad, T. A., 1842, Observations of the Silurian and Devonian systems of the U.S., with descriptions of new organic remains: *Journal of the Academy of Natural Sciences of Philadelphia*, v. 8, no. 2, p. 228-235.
- Coogan, Alan H., 1996, Ohio's surface rocks and sediments, in Feldmann, Rodney M., and Hackathorn, Merrienne, eds., *Fossils of Ohio: Ohio Division of Geological Survey Bulletin 70*, p. 31-50.
- Copper, Paul, 1977, *Zygospira* and some related Ordovician and Silurian atrypoid brachiopods: *Palaeontology*, v. 20, pt. 2, p. 295-335, pls. 37-40.
- Corneliussen, Eric F., and Perry, T. G., 1970, The ectoproct *Batostoma? cornula* (Cumings & Galloway) and its enigmatic intrazoecial spines [Fort Atkinson Limestone (Cincinnatian), Wilmington, Illinois]: *Journal of Paleontology*, v. 44, no. 6, p. 997-1008, pls. 133-135.
- Corneliussen, Eric F., and Perry, T. G., 1973, *Monotrypa*, *Hallopora*, *Amplexopora*, and *Hennigopora* (Ectoprocta) from the Brownsport Formation (Niagaran), western Tennessee: *Journal of Paleontology*, v. 47, no. 2, p. 151-220, 10 pls.
- Cressman, E. R., 1964, Geology of the Tyrone quadrangle, Kentucky: U.S. Geological Survey Geologic Quadrangle Map GQ-303, scale 1:24,000.
- Cross, Aural T., Taggart, Ralph E., and Gillespie, William H., 1996, The fossil plants of Ohio: introduction, overview, and nonvascular plants, in Feldmann, Rodney M., and Hackathorn, Merrienne, eds., *Fossils of Ohio: Ohio Division of Geological Survey Bulletin 70*, p. 370-395.
- Crowther, P. R., and Bergström, S. M., 1980, Cortical bandages in an Upper Ordovician glyptograptid graptolite from Ohio: *Bollettino della Società Paleontologica Italiana*, v. 19, p. 250-253.
- Cuffey, Roger J., Butler, Katharine L., and Rockwell, Lynn A., 1996, Bryozoan diversity and salinity in the uppermost Ordovician (Saluda and Whitewater Formations) across southeastern Indiana and southwestern Ohio (abs.): *Geological Society of America Abstracts with Programs*, v. 28, no. 6, p. 34-35.
- Cuffey, Roger J., and Kamandulis, Michael A., 1985, Trepostome bryozoan reef-mounds in the Upper Ordovician near Maysville, Kentucky (abs.): *Geological Society of America Abstracts with Programs*, v. 17, no. 5, p. 283.
- Cuffey, Roger J., and Perry, T. G., 1964, Redescription of the syntypes of the bryozoan species *Rhombotrypa quadrata* (Rominger): *University of Michigan Museum of Paleontology Contributions*, v. 19, no. 4, p. 37-45, pls. 1-2.
- Cuffey, Roger J., and Pursell, Benjamin F., 1995, A possible causal explanation for the Maysville bryozoan reefs (Upper Ordovician, north-central Kentucky) (abs.): *Geological Society of America Abstracts with Programs*, v. 27, no. 3, p. 45.
- Cumings, Edgar R., 1901a, Notes on the Ordovician rocks of southern Indiana: *Indiana Academy of Science Proceedings* for 1900, p. 200-215.
- Cumings, Edgar R., 1901b, A section of the Upper Ordovician at Vevay, Indiana: *American Geologist*, v. 28, no. 6, p. 361-379, pls. 34-35.
- Cumings, Edgar R., 1902, A revision of the bryozoan genera *Dekayia*, *Dekayella*, and *Heterotrypa* of the Cincinnati Group: *American Geologist*, v. 29, no. 4, p. 197-217, pls. 9-12.
- Cumings, Edgar Roscoe, 1903, The morphogenesis of *Platystrophia*: a study of the evolution of a Paleozoic brachiopod: *American Journal of Science*, ser. 4, v. 15, no. 85, p. 1-48; no. 86, p. 121-136, fig. 27 [fold-out plate].
- Cumings, Edgar Roscoe, 1904, Development of some Paleozoic Bryozoa: *American Journal of Science*, ser. 4, v. 17, no. 97, p. 49-78.
- Cumings, E. R., 1908, The stratigraphy and paleontology of the Cincinnati Series of Indiana: *Indiana Department of Geology and Natural Resources, Thirty-second Annual Report for 1907*, p. 605-1189, 55 pls.
- Cumings, E. R., 1912, Development and systematic position of the monticuliporoids: *Geological Society of America Bulletin*, v. 23, p. 357-370, pls. 19-22.
- Cumings, E. R., 1922, Nomenclature and description of the geological formations of Indiana, in Logan, W. N., and others, *Handbook of Indiana geology: Indiana Department of Conservation Publication 21*, p. 403-570.
- Cumings, E. R., and Galloway, J. J., 1912, A note on the *Batostomas* of the Richmond Series: *Proceedings of the Indiana Academy of Science for 1911*, p. 147-167, pls. 1-7.
- Cumings, E. R., and Galloway, J. J., 1913, The stratigraphy and paleontology of the Tanner's Creek section of the Cincinnati Series in Indiana: *Indiana Department of Geology and Natural Resources, Thirty-seventh Annual Report for 1912*, p. 353-479, 20 pls.
- Cumings, E. R., and Galloway, J. J., 1915, Studies of the morphology and history of the Trepostomata or monticuliporoids: *Geological Society of America Bulletin*, v. 26, p. 349-374, pls. 10-15.
- Cumings, Edgar R., and Mauck, Abram V., 1902, A quantitative study of variation in the fossil brachiopod *Platystrophia lynx*: *American Journal of Science*, ser. 4, v. 14, no. 79, p. 9-16, pls. 2-3.
- Cutler, John Frederick, 1968, Morphology, taxonomy, and evolution of the bryozoan *Constellaria* from the Cincinnati Arch: Ph.D. dissertation (unpub.), Columbia University, 297 p.
- Dalvé, Elizabeth, compiler, 1948, The fossil fauna of the Ordovician in the Cincinnati region: University Museum, Department of Geology and Geography, University of Cincinnati, 56 p.
- Daly, Edward Joseph, 1979, The trepostomatous bryozoan genus *Hallopora* Bassler in the Dillsboro Formation (Cincinnatian) of southeastern Indiana: M.S. thesis (unpub.), Boston College, 93 p.
- Daly, Edward, and Brown, George D., Jr., 1979, The trepostomatous bryozoan genus *Hallopora* Bassler in the Dillsboro Formation (Cincinnatian) of southeastern Indiana (abs.): *Geological Society of America Abstracts with Programs*, v. 11, no. 1, p. 9.
- Dattilo, Benjamin F., 1991, The Miamitown Shale (Upper Ordovician): a facies after all (abs.): *Geological Society of America Abstracts with Programs*, v. 23, no. 3, p. 9.
- Dattilo, Benjamin F., 1992, Unravelling Cincinnatian stratigraphy: high-resolution ecologic and bio-event stratigraphy of the Miamitown Shale (Ordovician) (abs.): *Geological Society of America Abstracts with Programs*, v. 24, no. 4, p. 11.
- Dattilo, Benjamin F., 1996, A quantitative paleoecological approach to high-resolution cyclic and event stratigraphy: the Upper Ordovician Miamitown Shale in the type Cincinnatian: *Lethaia*, v. 29, no. 1, p. 21-37.
- Davies, Rhys J., 1958, A correlation study of the Cynthiana, Gallatin County, Kentucky: M.S. thesis (unpub.), Miami University, 74 p.
- Davis, Paul E., Lerch, Norbert, Tornes, Larry, Steiger, Joseph, Smeck, Neil, Andrus, Howard, Trimmer, John, and Bottrell, George, 1976, Soil survey of Montgomery County, Ohio: U.S. Department of Agriculture, Soil and Conservation Service, 107

- p., 83 pls. + index map.
- Davis, R. A., ed., 1981, Cincinnati fossils, an elementary guide to the Ordovician rocks and fossils of the Cincinnati, Ohio, region: Cincinnati Museum of Natural History, 58 p., 8 pls.
- Davis, R. A., ed., 1985, Cincinnati fossils, an elementary guide to the Ordovician rocks and fossils of the Cincinnati, Ohio, region: Cincinnati Museum of Natural History, 61 p., 8 pls.
- Davis, R. A., 1986, Cincinnati region: Ordovician stratigraphy near the southwest corner of Ohio, *in* Neathery, Thornton L., ed., Southeastern Section of the Geological Society of America: Geological Society of America, Centennial Field Guide, v. 6, p. 21-24.
- Davis, R. A., ed., 1992, Cincinnati fossils, an elementary guide to the Ordovician rocks and fossils of the Cincinnati, Ohio, region: Cincinnati Museum of Natural History, 61 p., 8 pls.
- Davis, Richard Arnold, and Mapes, Royal H., 1996, Phylum Mollusca, Class Cephalopoda, *in* Feldmann, Rodney M., and Hackathorn, Merrienne, eds., Fossils of Ohio: Ohio Division of Geological Survey Bulletin 70, p. 166-195.
- Decker, Charles E., 1952, Graptolites from Kentucky and Kaukauna, Wisconsin: American Association of Petroleum Geologists Bulletin, v. 36, no. 1, p. 160-163.
- Delmet, Dale A., and Anstey, Robert L., 1974, Fourier analysis of morphological plasticity within an Ordovician bryozoan colony: *Journal of Paleontology*, v. 48, no. 2, p. 217-226.
- Dick, Jeffrey C., and Shakoor, Abdul, 1993, Clay mineralogy of Ohio mudrocks (abs.): *Ohio Journal of Science*, v. 93, no. 2, p. 40.
- Diekmeyer, Sharon C., 1990, Quantitative analysis of cyclical faunal patterns in the Upper Ordovician Kope through Bellevue sequence, Cincinnati, Ohio: M.S. thesis (unpub.), University of Cincinnati, 168 p., 15 pls.
- Diekmeyer, Sharon C., 1992, Bedrock geology and paleontology (stop #7), *in* Haneberg, William C., Riestenberg, Mary M., Pohana, Richard E., and Diekmeyer, Sharon C., Cincinnati's geologic environment: a trip for secondary-school science teachers: Ohio Division of Geological Survey Guidebook 9, p. 13-18.
- Dively, C. Gordon, and Cuffey, Roger J., 1994, *Monticulipora molestata*—intraspecific?? variability (Bryozoa, Upper Ordovician, southeastern Indiana) (abs.): Geological Society of America Abstracts with Programs, v. 26, no. 3, p. 15.
- Donovan, Stephen K., Kallmeyer, Jack W., and Veltkamp, Cornelius J., 1995, Functional morphologies of the columns of Upper Ordovician *Xenocrinus* and *Dendrocrinus*: *Lethaia*, v. 28, no. 4, p. 309-315.
- Douglas, Dale, 1970, Midwest fossils—pt. VIII, worm fossils, conodonts: *Earth Science*, v. 23, no. 3, p. 127-134.
- Drake, Daniel, 1825, Geological account of the Valley of the Ohio: *Transactions of the American Philosophical Society*, new ser., v. 2, p. 124-139.
- Droste, John B., and Shaver, Robert H., 1983, Atlas of early and middle Paleozoic paleogeography of the southern Great Lakes area: Indiana Geological Survey Special Report 32, 32 p.
- Dry Dredgers. See Martin, Wayne, and Shideler, W. H., 1958.
- Dunn, Paul Heaney, 1924, The Cynthiana Formation of north central Kentucky: M.A. thesis, The Ohio State University, 58 p.
- Durrell, Richard H., 1967, Invasion by sea and ice: *The Explorer*, v. 9, no. 3, p. 12-16.
- Durrell, Richard H., 1973, Collecting sea shells by the seashore: *Earth Science*, v. 26, no. 4, p. 175-178.
- Durrell, Richard H., 1977, A recycled landscape: Cincinnati Museum of Natural History Quarterly, v. 14, no. 2, p. 8-15 (reprinted as Popular Publication Series 9, 9 p.).
- Dyche, D. T. D., 1892a, On termination of columns of crinoids from the Cincinnati Group: *American Geologist*, v. 10, no. 2, p. 130.
- Dyche, D. T. D., 1892b, The crinoid *Heterocrinus subcrassus*: *Science*, v. 20, no. 459, p. 66.
- Dyche, D. T. D., 1892c, Remarks on the stems and roots of crinoids from near Lebanon, Ohio: *Journal of the Cincinnati Society of Natural History*, v. 15, no. 2, p. 101.
- Dyer, William S., 1923, Stratigraphy and paleontology of the Credit River section of the upper Cincinnatian Series of Ontario: Ph.D. dissertation (unpub.), University of Toronto, 62 p., 8 pls.
- Eckert, J. D., 1988, Late Ordovician extinction of North American and British crinoids: *Lethaia*, v. 21, no. 2, p. 147-167.
- Edwards, William R., 1957, Petrography and petrology of the Cynthiana Limestone and lower Eden strata of southwestern Ohio and northern Kentucky: M.S. thesis (unpub.), The Ohio State University, 200 p., 4 pls.
- Elias, Maxim K., 1954, *Cambroporella* and *Coeloclema*, Lower Cambrian and Ordovician bryozoans: *Journal of Paleontology*, v. 28, no. 1, p. 52-58, pls. 9-10.
- Elias, Robert J., 1980a, Borings in solitary rugose corals of the Selkirk Member, Red River Formation (late Middle or Upper Ordovician), southern Manitoba: *Canadian Journal of Earth Sciences*, v. 17, no. 2, p. 272-277.
- Elias, Robert J., 1980b, Upper Ordovician solitary rugose corals of the Cincinnati Arch region, Ohio-Indiana-Kentucky: Geological Society of American Abstracts with Programs, v. 12, no. 5, p. 224-225.
- Elias, Robert J., 1981, Late Ordovician solitary rugose corals of the Upper Mississippi Valley region (abs.): Geological Society of America Abstracts with Programs, v. 13, no. 6, p. 276.
- Elias, Robert J., 1982, Latest Ordovician solitary rugose corals of eastern North America: *Bulletins of American Paleontology*, v. 81, no. 314, 116 p., 15 pls.
- Elias, Robert J., 1983a, Late Ordovician solitary rugose corals of the Stony Mountain Formation, southern Manitoba, and its equivalents: *Journal of Paleontology*, v. 57, no. 5, p. 924-956.
- Elias, Robert J., 1983b, Middle and Late Ordovician solitary rugose corals of the Cincinnati Arch region: U.S. Geological Survey Professional Paper 1066-N, 13 p., 2 pls.
- Elias, Robert J., 1984a, Paleobiologic significance of fossulae in North American Late Ordovician solitary rugose corals: *Paleobiology*, v. 10, no. 1, p. 102-114.
- Elias, Robert J., 1984b, Paleobiology of solitary rugose corals, Late Ordovician of North America: *Palaeontographica Americana*, no. 54, p. 533-537.
- Elias, Robert J., 1984c, Taphonomy of Late Ordovician solitary rugose corals: paleoecologic and sedimentologic significance (abs.): Geological Society of America Abstracts with Programs, v. 16, no. 3, p. 136.
- Elias, Robert J., 1986a, New Late Ordovician solitary rugose coral with perforate septa: *Journal of Paleontology*, v. 60, no. 1, p. 14-25.
- Elias, Robert J., 1986b, Symbiotic relationships between worms and solitary rugose corals in the Late Ordovician: *Paleobiology*, v. 12, no. 1, p. 32-45.
- Elias, R. J., 1989a, Extinctions and origins of solitary rugose corals, latest Ordovician to earliest Silurian in North America, *in* Jell, P. A., and Pickett, J. W., Proceedings of the Fifth International Symposium on Fossil Cnidaria Including Archaeocyathata and Spongiomorphs, Brisbane, Queensland, July 25-29, 1988: Association of Australasian Palaeontologists Memoir 8, p. 319-326.
- Elias, Robert J., 1989b, Extinctions and origins of solitary rugose corals, latest Ordovician to earliest Silurian in North America (abs.): Geological Society of America Abstracts with Programs, v. 21, no. 4, p. 10.
- Elias, Robert J., 1992, Environmental cycles in the Upper Ordovician of cratonic North America (abs.): Geological Society of America Abstracts with Programs, v. 24, no. 4, p. 14.
- Elias, Robert J., Brandt, Danita S., and Clark, T. H., 1990, Late Ordovician solitary rugose corals of the St. Lawrence Lowland, Québec: *Journal of Paleontology* v. 64, no. 3, p. 340-352.
- Elias, Robert J., Zeilstra, Ronald G., and Bayer, Thomas N., 1988, Paleoenvironmental reconstruction based on horn corals, with an example from the Late Ordovician of North America: *Palaios*, v. 3, no. 1, p. 22-34.
- Errett, Douglas H., and Cuffey, Roger J., 1989, Bryozoan mode of occurrence relative to individual beds in the type Cincinnatian (Upper Ordovician; Ohio, Indiana, Kentucky) (abs.): Geological Society of America Abstracts with Programs, v. 21, no. 4, p. 11.
- Ethington, R. L., Droste, J. B., and Rexroad, C. B., 1986, Conodonts from subsurface Champlainian (Ordovician) rocks of eastern

- Indiana: Indiana Geological Survey Special Report 37, 32 p., 2 pls.
- Ewers, Ralph O., 1981, A guidebook for student tour of Cincinnati geology: Sponsored by Geological Society of America in cooperation with Cincinnati Public Schools, Cincinnati Museum of Natural History, and the Geology Department of Eastern Kentucky University, 34 p. (unnumbered).
- Faber, Chas. L., 1886, Remarks on some fossils of the Cincinnati Group: Journal of the Cincinnati Society of Natural History, v. 9, no. 1, p. 14-20, pl. I.
- Farber, Steve L., 1968, The petrology of selected limestones of the Southgate and McMicken Members, Latonia Formation, Cincinnati Series, in southern Hamilton County, Ohio: M.S. thesis (unpub.), Miami University, 121 p., 11 pls.
- Felton, Stephen H., and Morris, Robert W., 1990, Symbiotic association of crinoids, platycerid gastropods, and *Cornulites* in the Upper Ordovician (Cincinnati) of the Cincinnati region (abs.): Ohio Journal of Science, v. 90, no. 2, p. 12.
- Fenneman, Nevin M., 1916, Geology of Cincinnati and vicinity: Ohio Division of Geological Bulletin 19, 207 p.
- Fillion, D., and Pickerill, R. K., 1984, On *Arthraria antiquata* Billings, 1872 and its relationship to *Diplocraterion* Torell, 1870 and *Bifungites* Desio, 1940: Journal of Paleontology, v. 58, no. 3, p. 683-696.
- Finney, Stanley C., 1986, Graptolite biofacies and correlation of eustatic, subsidence, and tectonic events in the Middle to Upper Ordovician of North America: Palaios, v. 1, no. 5, p. 435-461.
- Fleming, R. W., 1985, Geologic perspectives—the Cincinnati example, in Slope stability and landslides: Proceedings of the 6th Ohio Valley Soils Seminar, Kentucky Geotechnical Group, University of Kentucky, and University of Louisville, p. 1-22.
- Fleming, Robert W., and Johnson, Arvid M., 1994, Landslides of the Cincinnati, Ohio, area, Part B of Landslides in colluvium: U.S. Geological Survey Bulletin 2059-B, 24 p., 1 pl.
- Fleming, Robert W., Johnson, Arvid M., Hough, James E., Göcke, A. Önder, and Lion, Thomas, 1981, Engineering geology of the Cincinnati area, in Roberts, Thomas G., ed., GSA Cincinnati '81 Field Trip Guidebooks: American Geological Institute, v. 3, p. 543-570.
- Fleming, R. W., and Taylor, F. A., 1980, Estimating the cost of landslide damage in the United States: U.S. Geological Survey Circular 832, 21 p.
- Flood, Cathleen A., and Cuffey, Roger J., 1983, Trepostome bryozoan species from the upper Reedsville Shale (Late Ordovician) in central Pennsylvania—biostratigraphic and paleoecologic implications of a skeletally preserved faunule (abs.): Geological Society of America Abstracts with Programs, v. 15, no. 3, p. 198-199.
- Flower, Rousseau H., 1939, Structure and taxonomic position of *Troedssonoceras* Foerste: Journal of Paleontology, v. 13, no. 5, p. 481-484, pl. 50.
- Flower, Rousseau H., 1943a, Ordovician cephalopods of the Cincinnati region, Pt. 1 (abs.): Geological Society of America Bulletin, v. 54, no. 12, p. 1830-1831.
- Flower, Rousseau H., 1943b, Structure and relationship of Cincinnati Cyrtocera: Ohio Journal of Science, v. 43, no. 2, p. 51-64, 2 pls.
- Flower, Rousseau H., 1946, Ordovician cephalopoda of the Cincinnati region. Part I: Bulletins of American Paleontology, v. 29, no. 116, p. 83-738, 50 pls.
- Flower, Rousseau H., 1955, Trails and tentacular impressions of orthoconic cephalopods: Journal of Paleontology, v. 29, no. 5, p. 857-867.
- Fluegeman, Richard H., Jr., 1979, The New Point Tongue of the Brainard Shale (Upper Ordovician) in southeastern Indiana, stratigraphy and community succession: M.S. thesis (unpub.), Miami University, 104 p., 4 pls., 16 text-figs.
- Fluegeman, Richard H., Jr., 1986, Guidebook to collecting localities in the Dillsboro, Saluda, and Whitewater Formations (Ordovician; Richmondian) of southeastern Indiana: National Association of Geology Teachers, East Central Section, Annual Field Trip, Ball State University, Department of Geology, 15 p.
- Fluegeman, Richard H., Jr., and Pope, John K., 1983, Brainard Shale outliers (Upper Ordovician, Maquoketa Group) in southeastern Indiana (abs.): Geological Society of America Abstracts with Programs, v. 15, no. 6, p. 574.
- Foerste, Aug. F., 1887, The Clinton Group of Ohio, Part III: Bulletin of the Scientific Laboratories of Denison University, v. 2, pt. 2, p. 149-176, pls. 15-17.
- Foerste, Aug. F., 1888a, Notes on a geological section at Todd's Fork, Ohio: American Geologist, v. 2, no. 6, p. 412-419.
- Foerste, Aug. F., 1888b, Notes on Paleozoic fossils: Bulletin of the Scientific Laboratories of Denison University, v. 3, art. 1, p. 117-137, pl. 13.
- Foerste, Aug. F., 1896, An account of the Middle Silurian rocks of Ohio and Indiana: Journal of the Cincinnati Society of Natural History, v. 18, nos. 3 and 4, p. 161-200.
- Foerste, Aug. F., 1900, A general discussion of Middle Silurian rocks of the Cincinnati anticlinal region, with their synonymy: Indiana Department of Geology and Natural Resources, Twenty-fourth Annual Report for 1899, p. 41-80.
- Foerste, Aug. F., 1902, The Cincinnati Anticline of southern Kentucky: American Geologist, v. 30, no. 6, p. 359-369, 1 pl.
- Foerste, Aug. F., 1903, The Richmond Group along the western side of the Cincinnati Anticline in Indiana and Kentucky: American Geologist, v. 31, no. 6, p. 333-361, pls. 20-22.
- Foerste, Aug. F., 1904a, Ordovician, in Hopkins, T. C., A short description of the topography of Indiana, and of the rocks of the different geologic periods; to accompany the geological map of the state: Indiana Department of Geology and Natural Resources, Twenty-eighth Annual Report for 1903, p. 21-27.
- Foerste, Aug. F., 1904b, The Ordovician-Silurian contact in the Ripley Island area of southern Indiana, with notes on the age of the Cincinnati geanticline: American Journal of Science, ser. 4, v. 18, no. 107, p. 321-342, 1 pl.
- Foerste, Aug. F., 1904c, Variations in thickness of the subdivisions of the Ordovician of Indiana: American Geologist, v. 34, no. 2, p. 87-102, pl. 5.
- Foerste, August F., 1905a, The classification of the Ordovician rocks of Ohio and Indiana: Science, new ser., v. 22, no. 553, p. 149-152.
- Foerste, Aug. F., 1905b, Note on the distribution of Brachiopoda in the Arnheim and Waynesville beds: American Geologist, v. 36, no. 4, p. 244-250.
- Foerste, Aug. F., 1906, The Silurian, Devonian and Irvine formations of east-central Kentucky, with an account of their clays and limestones: Kentucky Geological Survey Bulletin 7, 369 p.
- Foerste, Aug. F., 1909a, Preliminary notes on Cincinnati fossils: Bulletin of the Scientific Laboratories of Denison University, v. 14, art. 13, p. 209-231, pl. 4.
- Foerste, Aug. F., 1909b, Preliminary notes on Cincinnati and Lexington fossils: Bulletin of the Scientific Laboratories of Denison University, v. 14, art. 17, p. 289-333, pls. 7-11.
- Foerste, Aug. F., 1910, Preliminary notes on Cincinnati and Lexington fossils of Ohio, Indiana, Kentucky, and Tennessee: Bulletin of the Scientific Laboratories of Denison University, v. 16, art. 2, p. 17-100, pls. 1-6.
- Foerste, Aug. F., 1912a, The Arnheim Formation within the areas traversed by the Cincinnati Geanticline: Ohio Naturalist, v. 12, no. 3, p. 429-456.
- Foerste, Aug. F., 1912b, *Strophomena* and other fossils from Cincinnati and Mohawkian horizons, chiefly in Ohio, Indiana and Kentucky: Bulletin of the Scientific Laboratories of Denison University, v. 17, art. 2, p. 17-172, 18 pls.
- Foerste, Aug. F., 1914a, Notes on Agelacrinidae and Lepadocystinae, with descriptions of *Thresherodiscus* and *Brockocystis*: Bulletin of the Scientific Laboratories of Denison University, v. 17, art. 14, p. 399-487, 4 pls. (no italics in original).
- Foerste, Aug. F., 1914b, The Rogers Gap fauna of central Kentucky: Journal of the Cincinnati Society of Natural History, v. 21, no. 4, p. 109-156, pls. 1-4.
- Foerste, Aug. F., 1916, Notes on Cincinnati fossil types: Bulletin of the Scientific Laboratories of Denison University, v. 18, art. 4, p. 285-355, pls. 1-7.
- Foerste, Aug. F., 1917a, Notes on Richmond and related fossils:

- Journal of the Cincinnati Society of Natural History, v. 22, no. 2, p. 42-55, 3 pls.
- Foerste, Aug. F., 1917b, Notes on Silurian fossils from Ohio and other central states [concluded from April number]; Appendix: Two species of Ordovician fossils: Ohio Journal of Science, v. 17, no. 7, p. 233-267, pls. 8-12.
- Foerste, Aug. F., 1919a, Notes on *Isotelus*, *Acrolichas*, *Calymene*, and *Encrinurus*: Bulletin of the Scientific Laboratories of Denison University, v. 19, art. 1, p. 65-82, pls. 14-18.
- Foerste, Aug. F., 1919b, Silurian fossils from Ohio, with notes on related species from other horizons: Ohio Journal of Science, v. 19, no. 7, p. 367-404, pls. 16-19.
- Foerste, Aug. F., 1920, The generic relations of the American Ordovician Lichadidae: American Journal of Science, v. 49, no. 289, p. 26-50, pls. I-IV.
- Foerste, Aug. F., 1924a, Notes on American Paleozoic cephalopods: Journal of the Scientific Laboratories of Denison University, v. 20, art. 10, p. 193-268, pls. 21-42.
- Foerste, A. F., 1924b, Upper Ordovician faunas of Ontario and Quebec: Geological Survey of Canada Memoir 138, 255 p., 46 pls.
- Foerste, Aug. F., 1926, Actinosiphonate, trochoceroid and other cephalopods: Journal of the Scientific Laboratories of Denison University, v. 21, art. 5, p. 285-383, pls. 32-53.
- Foerste, Aug. F., 1927, Ordovician and Silurian cephalopods of the Hudson Bay area: Journal of the Scientific Laboratories of Denison University, v. 22, art. 1, p. 1-107, pls. 1-24.
- Foerste, Aug. F., 1928a, American arctic and related cephalopods: Journal of the Scientific Laboratories of Denison University, v. 23, art. 1, p. 1-110, pls. 1-29.
- Foerste, Aug. F., 1928b, A restudy of American orthoconic Silurian cephalopods: Journal of the Scientific Laboratories of Denison University, v. 23, art. 7, p. 236-320, pls. 48-75.
- Foerste, Aug. F., 1928c, Some hitherto unfigured Ordovician cephalopods from Anticosti Island: Transactions of the Royal Society of Canada, third ser., v. 22, sec. 4, p. 223-234, pls. 1-11.
- Foerste, Aug. F., 1929a, The Ordovician and Silurian of American arctic and subarctic regions: Journal of the Scientific Laboratories of Denison University, v. 24, art. 2, p. 27-79, pls. 2, 3.
- Foerste, Aug. F., 1929b, The cephalopods of the Red River Formation of southern Manitoba: Journal of the Scientific Laboratories of Denison University, v. 24, art. 7, p. 129-235, pls. 11-39.
- Foerste, Aug. F., 1930, Three studies of cephalopods: Journal of the Scientific Laboratories of Denison University, v. 24, art. 10, p. 265-381, pls. 41-63 (date nominally December 1929, but published the next month).
- Foerste, Aug. F., 1932, Black River and other cephalopods from Minnesota, Wisconsin, Michigan, and Ontario (Part I): Journal of the Scientific Laboratories of Denison University, v. 27, art. 2, p. 47-136, pls. 7-37.
- Foerste, Aug. F., 1933, Black River and other cephalopods from Minnesota, Wisconsin, Michigan, and Ontario (Part II): Journal of the Scientific Laboratories of Denison University, v. 28, art. 1, p. 1-146.
- Foerste, Aug. F., 1935, Big Horn and related cephalopods: Journal of the Scientific Laboratories of Denison University, v. 30, art. 1, p. 1-96, pls. 1-22.
- Foerste, Aug. F., 1936, The cephalopods of the Maquoketa Shale of Iowa: Journal of the Scientific Laboratories of Denison University, v. 30, art. 5, p. 231-257, pls. 27-37 (date nominally December 1935, but published the next month).
- Foerste, Aug. F., and Cox, Ian H., 1936, Cephalopods and a *Beatricea* from Akpatok Island: Geological Magazine, v. 73, no. 865, p. 289-307, pl. 4.
- Foerste, Aug. F., and Teichert, Curt, 1930, Actinoceroids of east-central North America: Journal of the Scientific Laboratories of Denison University, v. 25, art. 6, p. 201-296, pls. 27-49.
- Foord, A. H., 1884, On three new species of monticuliporoid corals: Annals and Magazine of Natural History, series 5, v. XII, p. 338-342, pl. 12.
- Ford, John Philip, 1965, Bedrock geology in southwest Hamilton County, Ohio: Ph.D. dissertation (unpub.), The Ohio State University, 103 p., 3 pls.
- Ford, J. P., 1967, Cincinnati geology in southwest Hamilton County, Ohio: American Association of Petroleum Geologists Bulletin, v. 51, no. 6, p. 918-936.
- Ford, J. P., 1968a, Cincinnati geology in southwest Hamilton County, Ohio (abs.), in Abstracts for 1966: Geological Society of America Special Paper 101, p. 357.
- Ford, J. P., 1968b, Upper Ordovician stratigraphic relations between Covington and Clays Ferry, Kentucky: American Association of Petroleum Geologists Bulletin, v. 52, no. 9, p. 1779-1791.
- Ford, John P., 1972, Bedrock geology of the Addyston quadrangle and part of the Burlington quadrangle, Hamilton County, Ohio: Ohio Division of Geological Survey Report of Investigations 83, 1 sheet.
- Ford, John P., 1974, Bedrock geology of the Cincinnati West quadrangle and part of the Covington quadrangle, Hamilton County, Ohio: Ohio Division of Geological Survey Report of Investigations 93, 1 sheet.
- Forsyth, Jane Louise, 1946, The Eden and Maysville groups of the Cincinnati series at Cincinnati, Ohio: M.A. thesis (unpub.), University of Cincinnati, 122 p.
- Fox, William T., 1962, Stratigraphy and paleoecology of the Richmond Group in southeastern Indiana: Geological Society of America Bulletin, v. 73, no. 5, p. 621-641.
- Frank, Glenn W., ed., 1969, Ohio intercollegiate field trip guides 1950-51 to 1969-70: Kent State University Printing Service, p. i-vii, 1-1 through 20-27 (field trip 9, p. 9-1 through 9-15, and field trip 17, p. 17-1 through 17-40, include Cincinnati stops).
- Freas, Robert C., 1968, The faunal distribution of the Blanchester Member of the Waynesville Formation in Indiana, Ohio, and Kentucky: M.S. thesis (unpub.), Miami University, 121 p.
- Freeman, L. B., 1953, Regional subsurface stratigraphy of the Cambrian and Ordovician in Kentucky and vicinity: Kentucky Geological Survey Bulletin, ser. IX, no. 12, 352 p., 10 pls.
- Frest, T. J., Strimple, H. L., and Kelly, S. M., 1976, A new Ordovician camerate crinoid from Kentucky: Southeastern Geology, v. 17, no. 3, p. 139-148.
- Frey, Robert C., 1976, The biostratigraphy of the Richmond Group (Upper Ordovician), Franklin County, Indiana: M.S. thesis (unpub.), Miami University, 87 p.
- Frey, Robert C., 1980, *Vanuxemia waynesvillensis*, a new species of cyrtodontid pelecypod from the Upper Ordovician of southwest Ohio: Journal of Paleontology, v. 54, no. 4, p. 740-744, 1 pl.
- Frey, Robert C., 1981, *Nartheoceras* (Cephalopoda) from the Upper Ordovician (Richmondian) of southwest Ohio: Journal of Paleontology, v. 55, no. 6, p. 1217-1224.
- Frey, Robert Charles, 1983, The paleontology and paleoecology of the *Treptoceras duseri* shale unit (Late Ordovician, Richmondian) of southwestern Ohio: Ph.D. dissertation (unpub.), Miami University, 2 v., 719 p., 23 pls.
- Frey, Robert C., 1985a, Late Ordovician pelecypod faunas from the Cincinnati, Ohio area (abs.): Geological Society of America Abstracts with Programs, v. 17, no. 7, p. 587.
- Frey, Robert C., 1985b, A well-preserved specimen of *Schuchertoceras* (Cephalopoda, Ascocerida) from the Upper Ordovician (basal Richmondian) of southwest Ohio: Journal of Paleontology, v. 59, no. 6, p. 1506-1511.
- Frey, Robert C., 1987a, The occurrence of pelecypods in Early Paleozoic epeiric-sea environments, Late Ordovician of the Cincinnati, Ohio area: Palaios, v. 2, no. 1, p. 3-23.
- Frey, Robert C., 1987b, The paleoecology of a Late Ordovician shale unit from southwest Ohio and southeastern Indiana: Journal of Paleontology, v. 61, no. 2, p. 242-267.
- Frey, Robert C., 1988a, Paleoecology of *Treptoceras duseri* (Murchelinoceeratida, Proteoceratidae) from Late Ordovician of southwestern Ohio: New Mexico Bureau of Mines and Mineral Resources Memoir 44, p. 79-102, 4 pls.
- Frey, R. C., 1988b, A well-preserved nautiloid assemblage from a Late Ordovician shale unit in southwestern Ohio (abs.): Society of Economic Paleontologists and Mineralogists Annual Midyear Meeting Abstracts, v. 5, p. 20.
- Frey, Robert C., 1989, Paleoecology of a well-preserved nautiloid assemblage from a Late Ordovician shale unit, southwestern

- Ohio: *Journal of Paleontology*, v. 63, no. 5, p. 604-620.
- Frey, Robert C., 1995, Middle and Upper Ordovician nautiloid cephalopods of the Cincinnati Arch region of Kentucky, Indiana, and Ohio: U.S. Geological Survey Professional Paper 1066-P, 126 p., 22 pls.
- Fritz, Madeleine A., 1926, The stratigraphy and paleontology of the Workman's Creek section of the Cincinnati Series of Ontario: Ph.D. dissertation (unpub.), University of Toronto, 44 p., 4 pls.
- Fürsich, Franz T., 1974, On *Diplocraterion* Torell, 1870 and the significance of morphological features in vertical spreiten-bearing U-shaped trace fossils: *Journal of Paleontology*, v. 48, no. 5, p. 952-962.
- Galloway, Jesse James, 1913, The stratigraphy and paleontology of the Tanner's Creek section of the Cincinnati Series of Indiana: Ph.D. dissertation (unpub.), Indiana University, 235 p. 20 pls.
- Galloway, J. J., and St. Jean, J., Jr., 1956, Bibliography of the order Stromatoporoidea: *Journal of Paleontology*, v. 30, no. 1, p. 170-185.
- Galloway, J. J., and St. Jean, J., Jr., 1961, Ordovician Stromatoporoidea of North America: *Bulletins of American Paleontology*, v. 43, no. 194, 102 p., 13 pls.
- Garner, Dale E., Reeder, Neil E., Ernst, James E., and Lerch, Norbert K., 1973, Soil survey of Warren County, Ohio: U.S. Department of Agriculture, Soil Conservation Service, 115 p., 30 maps + index map.
- Garrison, Gene, 1954, The lithology of the Saluda Formation: M.S. thesis (unpub.), Miami University, 110 p.
- Geitgey, J. E., 1985, Correspondence between Ordovician conodont diversity and isotopically inferred marine paleotemperatures (abs.): *Geological Society of America Abstracts with Programs*, v. 17, no. 2, p. 92.
- Gibbons, A. B., 1968, Geologic map of the Mays Lick quadrangle, Mason County, Kentucky: U.S. Geological Survey Geologic Quadrangle Map GQ-784, scale 1:24,000.
- Gibbons, A. B., 1971, Geologic map of the Alexandria quadrangle, Campbell and Kenton Counties, Kentucky: U.S. Geological Survey Geologic Quadrangle Map GQ-926, scale 1:24,000.
- Gibbons, A. B., 1972, Geologic map of parts of the Burlington and Addyston quadrangles, Boone County, Kentucky: U.S. Geological Survey Geologic Quadrangle Map GQ-1025, scale 1:24,000.
- Gibbons, A. B., 1973, Geologic map of parts of the Newport and Withamsville quadrangles, Campbell and Kenton Counties, Kentucky: U.S. Geological Survey Geologic Quadrangle Map GQ-1072, scale 1:24,000.
- Gibbons, A. B., 1975, Geologic map of the Worthville quadrangle, north-central Kentucky: U.S. Geological Survey Geologic Quadrangle Map GQ-1265, scale 1:24,000.
- Gibbons, A. B., Kohut, Joseph J., and Weiss, Malcolm P., 1975, Geologic map of the New Richmond quadrangle, Kentucky-Ohio: U.S. Geological Survey Geologic Quadrangle Map GQ-1228, scale 1:24,000.
- Gibbons, A. B., and Weiss, Malcolm P., 1972, Geologic map of the Maysville West quadrangle, Kentucky-Ohio: U.S. Geological Survey Geologic Quadrangle Map GQ-1005, scale 1:24,000.
- Giuseffi, David F., 1982, Paleocology and community analysis of selected shale intervals in the Mt. Hope Member of the Fairview Formation (Cincinnati Series, Ordovician), with a comparison to a shale unit in the Ft. Ancient Member of the Waynesville Formation: M. S. thesis (unpub.), Miami University, 145 p.
- Goldman, Daniel, 1995, Taxonomy, evolution, and biostratigraphy of the *Orthograptus quadrimucronatus* species group (Ordovician, Graptolithina): *Journal of Paleontology*, v. 69, no. 3, p. 516-540.
- Goldman, Daniel, and Bergström, Stig M., 1997, Late Ordovician graptolites from the North American midcontinent: *Palaeontology*, v. 40, no. 4, p. 965-1010, 3 pls.
- Goldman, D., and Mitchell, C. E., 1991, Revision of the Upper Ordovician graptolite *Diplograptus (Amplexograptus) recurvens richmondensis* Ruedemann: *Journal of Paleontology*, v. 65, no. 6, p. 1016-1017.
- Gotautas, Vito A., 1951, The Ostracoda of the Saluda Formation: M.S. thesis (unpub.), Miami University, 50 p.
- Gopinath, T., Lee, G., and Martin, W., 1975, Multivariate analysis of limestones within the Cincinnati Series (Upper Ordovician) of southwestern Ohio and north central Kentucky (abs.): American Association of Petroleum Geologists and Society of Economic Paleontologists and Mineralogists Annual Meeting Abstracts, p. 29-30.
- Grabau, Amadeus W., and Shimer, Hervey Woodburn, 1909, North American index fossils, invertebrates: New York, A. G. Seiler and Company, v. 1, 853 p., v. 2, 909 p.
- Graese, Anne M., 1988, Facies distribution within the Maquoketa and Galena-Platteville Groups and their relationship to Ordovician structural history of northeastern Illinois (abs.): *Geological Society of America Abstracts with Programs*, v. 20, no. 5, p. 345.
- Grahn, Yngve, and Bergström, Stig M., 1985, Chitinozoans from the Ordovician-Silurian boundary beds in the eastern Cincinnati region in Ohio and Kentucky: *Ohio Journal of Science*, v. 85, no. 4, p. 175-183.
- Gray, Henry H., 1972, Lithostratigraphy of the Maquoketa Group (Ordovician) in Indiana: Indiana Geological Survey Special Report 7, 31 p.
- Gray, H. H., Brown, G. D., Jr., and Lineback, J. A., 1966, Physical techniques of correlation applied to Upper Ordovician rocks of southeastern Indiana (abs.): American Association of Petroleum Geologists Bulletin, v. 50, no. 3, p. 615.
- Gray, Henry H., Droste, John B., Patton, John B., Rexroad, Carl B., and Shaver, Robert H., 1985, Correlation chart showing Paleozoic stratigraphic units of Indiana: Indiana Geological Survey supplement to Miscellaneous Map 48.
- Gray, Henry H., Forsyth, Jane L., Schneider, Allan F., and Gooding, Ansel M., 1972, Geologic map of the 1° by 2° Cincinnati quadrangle, Indiana and Ohio, showing bedrock and unconsolidated deposits: Indiana Geological Survey Regional Geologic Map No. 7.
- Gray, Jane, and Boucot, A. J., 1972, Palynological evidence bearing on the Ordovician-Silurian paraconformity in Ohio: *Geological Society of America Bulletin*, v. 83, no. 5, p. 1299-1314.
- Green, Darsie A., 1957, Trenton structure in Ohio, Indiana and northern Illinois: American Association of Petroleum Geologists Bulletin, v. 41, no. 4, p. 627-642.
- Gries, Paul, 1934, Upper Ordovician scolecodonts from the Cincinnati area (abs.): *Geological Society of America Proceedings*, 1933, p. 341.
- Grinnell, Geo. Bird, 1877, Notice of a new genus of annelids from the Lower Silurian: *American Journal of Science*, ser. 3, v. 14 (whole number 114), no. 81, p. 229-230.
- Guber, Albert Lee, 1962, Some Richmond (Ordovician) ostracodes from Indiana and Ohio: Ph.D. dissertation (unpub.), University of Illinois, 124 p.
- Guber, Albert L., 1971, Problems of sexual dimorphism, population structure and taxonomy of the Ordovician genus *Tetradella* (Ostracoda): *Journal of Paleontology*, v. 45, no. 1, p. 6-22, pls. 1-4.
- Gutstadt, Allan M., 1958, Upper Ordovician stratigraphy in eastern interior region: American Association of Petroleum Geologists Bulletin, v. 42, no. 3, p. 513-547.
- Hall, Donald D., 1960, Dalmanellidae of the Cincinnati: M.S. thesis (unpub.), University of Cincinnati, 112 p., 5 pls.
- Hall, Donald D., 1962, Dalmanellidae of the Cincinnati: *Paleontographica Americana*, v. 4, no. 29, p. 127-165, pls. 19-21.
- Hall, James, 1842, Notes upon the geology of the western states: *American Journal of Science*, v. 42, no. 1, p. 51-62.
- Hall, James, 1845, Description of some microscopic shells from the decomposing marl slate of Cincinnati: *American Journal of Science*, v. 48, no. 2, p. 292-295.
- Hall, James, 1861, Contributions to palaeontology, being the results of some of the investigations made during the years 1859 and 1860; observations upon some new and other species of fossils, from the rocks of the Hudson River group of Ohio and the western states; with descriptions: Fourteenth Annual Report

- of the Regents of the University of the State of New York, on the condition of the State Cabinet of Natural History, and the historical and antiquarian collection annexed thereto, appendix C, p. 89-92.
- Hall, James, 1872a, Descriptions of new species of Crinoidea and other fossils from strata of the age of the Hudson River Group and Trenton limestone: New York State Museum, 24th Annual Report, p. 205-232, pls. 7-8.
- Hall, James, 1872b, Descriptions of new species of fossils from the Hudson River Group in the vicinity of Cincinnati, Ohio: New York State Museum, 24th Annual Report, p. 225-232.
- Hall, James, 1883, Paleontology: Indiana Department of Geology and Natural Resources, Twelfth Annual Report for 1882, p. 239-400, 38 pls.
- Hall, James, and Whitfield, R. P., 1875, Descriptions of invertebrate fossils, mainly from the Silurian System: Geological Survey of Ohio, v. 2, pt. 2, p. 65-180, pls. 1-13.
- Hannibal, Joseph T., 1996a, Phylum Arthropoda: phyllocarids, millipedes, insects, and other less common forms, in Feldmann, Rodney M., and Hackathorn, Merrienne, eds., Fossils of Ohio: Ohio Division of Geological Survey Bulletin 70, p. 124-131 (just a mention that rare arthropods occur in the Ordovician of the state).
- Hannibal, Joseph T., 1996b, Ichnofossils, in Feldmann, Rodney M., and Hackathorn, Merrienne, eds., Fossils of Ohio: Ohio Division of Geological Survey Bulletin 70, p. 506-529.
- Hannibal, Joseph T., and Davis, Richard Arnold, 1992, Guide to the building stones of downtown Cincinnati: a walking tour: Ohio Division of Geological Survey Guidebook 7, 44 p.
- Hansen, Michael C., 1985, *Isotelus*—Ohio's state fossil: Ohio Division of Geological Survey, Ohio Geology Newsletter, Summer, p. 1-4.
- Hansen, Michael C., 1997, The geology of Ohio—the Ordovician: Ohio Division of Geological Survey, Ohio Geology, Fall, p. 1, 3-6.
- Hansen, Michael C., and Collins, Horace R., 1979, A brief history of the Ohio Geological Survey: Ohio Journal of Science, v. 79, no. 1, p. 3-14.
- Hansman, Robert H., Shaw, Frederick C., and Pettyjohn, Wayne A., 1962, Supplement to the catalog of type specimens of fossils in the University of Cincinnati Museum: University of Cincinnati, 131 p.
- Hariharan, Ganesan, 1970, Lithostratigraphy of subsurface Cincinnati strata in western and southern Ohio and adjacent areas: M.S. thesis (unpub.), The Ohio State University, 46 p.
- Harper, Arthur R., 1948, Ohio in the making: a brief geological history of Ohio: The Ohio State University, College of Education, 80 p.
- Harper, G. W., and Bassler, R. S., 1896, Catalogue of the fossils of the Trenton and Cincinnati Periods, occurring in the vicinity of Cincinnati, O.: Cincinnati, W. B. Carpenter Co., 34 p.
- Harris, Frank William, 1976, Textural and compositional variability in limestone beds from the Waynesville Formation (Upper Ordovician), Brookville, Indiana: M.S. thesis (unpub.), Miami University, 149 p.
- Harris, Frank W., and Martin, Wayne D., 1979, Benthic community development in limestone beds of the Waynesville (upper Dillsboro) Formation (Cincinnati Series, Upper Ordovician) of southeastern Indiana: Journal of Sedimentary Petrology, v. 49, no. 4, p. 1295-1306.
- Harrison, William B., III, 1978, The occurrence of larval and young post-larval juvenile Mollusca in the Upper Ordovician Cincinnati Series (abs.): Geological Society of America Abstracts with Programs, v. 10, no. 6, p. 256.
- Harrison, William B., III, 1979a, Paleogeologic interpretation of environmental stability, a different approach (abs.): American Association of Petroleum Geologists Bulletin, v. 63, no. 3, p. 463.
- Harrison, William B., III, 1979b, The use of larval and juvenile specimens for determining species distribution in ancient environments (abs.): Geological Society of America Abstracts with Programs, v. 11, no. 5, p. 231.
- Harrison, William B., III, 1984, Influence of physical disturbance and substrate condition on benthic molluscan assemblages preserved in the Kope Fm. (Upper Ordovician), Cincinnati, Ohio region (abs.): Geological Society of America Abstracts with Programs, v. 16, no. 3, p. 144.
- Hartwick, Wayde M., 1987, Taxonomy, morphology, and paleoecology of *Strophomena planumbona* (Hall), an articulate brachiopod from the Upper Ordovician of the Cincinnati, Ohio, area: M.S. thesis (unpub.), Miami University, 120 p.
- Hatfield, Craig Bond, 1968, Stratigraphy and paleoecology of the Saluda Formation (Cincinnati) in Indiana, Ohio, and Kentucky: Geological Society of America Special Paper 95, 34 p., 4 pls.
- Hattin, Donald E., Nosow, Edmund, Perkins, Ronald D., Stumm, Erwin C., Mound, Michael C., and Utgaard, John, 1961, Field excursion to the Falls of the Ohio, field trip 9: Guidebook for field trips, Cincinnati meeting, 1961, Geological Society of America, p. 295-350.
- Hay, Helen B., 1975, Lithofacies classification for the Cincinnati Series (Upper Ordovician), southeastern Indiana: M.S. thesis (unpub.), Miami University, 147 p.
- Hay, Helen B., 1977, Field trip No. 1, Cincinnati stratigraphy from Richmond to Aurora, Indiana, in Pope, John K., and Martin, Wayne D., eds., Field guidebook to the biostratigraphy and paleoenvironments of the Cincinnati Series in southeastern Indiana: Society of Economic Paleontologists and Mineralogists, Great Lakes Section, 7th Annual Field Conference, Miami University, p. I1-I33 (reprinted in 1979; see Pope and Martin, 1977).
- Hay, Helen B., 1981, Lithofacies and formations of the Cincinnati Series (Upper Ordovician), southeastern Indiana and southwestern Ohio: Ph.D. dissertation (unpub.), Miami University, 236 p., 3 pls.
- Hay, Helen B., Pope, John K., and Frey, Robert C., 1981, Lithostratigraphy, cyclic sedimentation, and paleoecology of the Cincinnati Series in southwestern Ohio and southeastern Indiana, in Roberts, Thomas G., ed., GSA Cincinnati '81 Field Trip Guidebooks: American Geological Institute, v. 1, p. 73-86.
- Hessin, William A., 1988, *Leviceraurus*, a new cheirurine trilobite from the Cobourg Formation (Middle-Upper Ordovician), southern Ontario, Canada: Journal of Paleontology, v. 62, no. 1, p. 87-93.
- Hickey, David R., 1987, Skeletal structure, development and elemental composition of the Ordovician trepostome bryozoan *Peronopora*: Palaeontology, v. 30, pt. 4, p. 691-716.
- Hickey, David R., 1988, Bryozoan astogeny and evolutionary novelties: their role in the origin and systematics of the Ordovician monticuliporid trepostome genus *Peronopora*: Journal of Paleontology, v. 62, no. 2, p. 180-203.
- Hinde, George Jennings, 1879, On annelid jaws from the Cambro-Silurian, Silurian, and Devonian formations in Canada and from the Lower Carboniferous in Scotland: Quarterly Journal of the Geological Society of London, v. 35, p. 370-389, pls. 18-20.
- Hinterlong, Gregory Dale, 1981, Paleogeologic succession in the Corryville Member (McMillan Formation, Upper Ordovician) Stonelick Creek, Clermont County, Ohio: M.S. thesis (unpub.), Miami University, 119 p.
- Hoare, Richard D., 1996, Phylum Mollusca, Class Rostroconchia, in Feldmann, Rodney M., and Hackathorn, Merrienne, eds., Fossils of Ohio: Ohio Division of Geological Survey Bulletin 70, p. 132-135 (just a mention that rostroconchs occur in the Ordovician of the state).
- Hoare, Richard D., and Miller, Barry B., 1996a, Phylum Mollusca, Class Pelecypoda, in Feldmann, Rodney M., and Hackathorn, Merrienne, eds., Fossils of Ohio: Ohio Division of Geological Survey Bulletin 70, p. 136-150.
- Hoare, Richard D., and Miller, Barry B., 1996b, Phylum Mollusca, Class Gastropoda, in Feldmann, Rodney M., and Hackathorn, Merrienne, eds., Fossils of Ohio: Ohio Division of Geological Survey Bulletin 70, p. 151-165.
- Hofmann, H. J., 1964, Deformational structures near Cincinnati, Ohio: Geological Society of America Bulletin, v. 77, no. 5, p. 533-548, 3 pls.
- Hofmann, H. J., 1966, Ordovician paleocurrents near Cincinnati, Ohio: Journal of Geology, v. 74, no. 6, p. 868-890.
- Hoge, Richard John, 1985, Conodont and brachiopod biostratigraphy

- in Upper Ordovician rocks near Brent, Kentucky: M.S. thesis (unpub.), The Ohio State University, 148 p.
- Holland, Steven M., 1985, Biogenic and physical sedimentation inside the Maysvillian brachiopod *Platystrophia ponderosa*: biostratigraphic implications (abs.): Ohio Journal of Science, v. 85, no. 2, p. 20.
- Holland, Steven M., 1987, Effects of sea-floor exposure on the preservation of an Ordovician brachiopod assemblage (abs.): Geological Society of America Abstracts with Programs, v. 19, no. 7, p. 705.
- Holland, Steven M., 1989a, Shoaling cycles, depth-dependent fossil assemblages, and epiboles in the Upper Ordovician of the Cincinnati Arch (abs.): Geological Society of America Abstracts with Programs, v. 21, no. 2, p. 23.
- Holland, Steven M., 1989b, Sequence-based correlation; sorting through the complicated stratigraphy of the eastern U.S. Ordovician (abs.): Geological Society of America Abstracts with Programs, v. 21, no. 6, p. 80.
- Holland, Steven M., 1989c, Taphonomic effects of sea-floor exposure on an Ordovician brachiopod assemblage: *Palaios*, v. 3, no. 6, p. 588-597.
- Holland, Steven M., 1990, Distinguishing eustasy and tectonics in foreland basin sequences: the Upper Ordovician of the Cincinnati Arch and Appalachian Basin: Ph.D. dissertation (unpub.), University of Chicago, 390 p.
- Holland, Steven M., 1993a, Sequence stratigraphy of a carbonate-clastic ramp: the Cincinnati Series (Upper Ordovician) in its type area: Geological Society of America Bulletin, v. 105, no. 3, p. 306-322.
- Holland, Steven M., 1993b, Sequence stratigraphy of a carbonate-clastic ramp: the Cincinnati Series (Upper Ordovician) in its type area: Reply: Geological Society of America Bulletin, v. 105, no. 12, p. 1639-1640.
- Holland, Steven M., 1997, Using time/environment analysis to recognize faunal events in the Upper Ordovician of the Cincinnati Arch, in Brett, C. E., ed., Paleontological event horizons: ecological and evolutionary implications: New York, Columbia University Press, p. 309-334.
- Holland, Steven M., and Patzkowsky, Mark, 1992, Sequence stratigraphy of the Middle and Upper Ordovician of the eastern United States (abs.): Geological Society of America Abstracts with Programs, v. 24, no. 4, p. 21.
- Holland, Steven M., and Patzkowsky, Mark E., 1996, Sequence stratigraphy and long-term lithologic change in the Middle and Upper Ordovician of the eastern United States, in Witzke, Brian J., Ludvigsen, Gregory A., and Day, Jed E., Paleozoic sequence stratigraphy: views from the North American craton: Geological Society of America Special Paper 306, p. 117-130.
- Holterhoff, Frank Koch, 1977, Upper Ordovician lithostratigraphy and conodont biostratigraphy at Carrollton, Kentucky: M.S. thesis (unpub.), The Ohio State University, 68 p.
- Hotchkiss, F. H. C., 1970, North American Ordovician Ophiuroidea—the genus *Taeniaster* Billings, 1858 (Protasteridae): Proceedings of the Biological Society of Washington, v. 83, no. 5, p. 59-76.
- Howe, Herbert J., 1965, Morphology of the brachiopod genera *Rhynchotrema*, *Hypsiptycha*, and *Lepidocyclus*: Journal of Paleontology, v. 39, no. 6, p. 1125-1128, pl. 134.
- Howe, Herbert J., 1979, Middle and Late Ordovician plectambonitacean, rhynchonellacean, syntrophiacean, trimerellacean, and atrypcean brachiopods: U.S. Geological Survey Professional Paper 1066-C, 18 p., 7 pls.
- Howe, Herbert J., 1982, Variation in the ventral muscle scars of the orthacean brachiopod *Plaesiomys subquadrata* (Hall) from Richmond (Upper Ordovician) strata in the Ohio Valley (abs.): Geological Society of America Abstracts with Programs, v. 14, no. 5, p. 263.
- Howe, Herbert J., 1985, Evolution of the Ordovician brachiopod *Hesperorthis tricenaria* (Conrad) and its stratigraphic distribution (abs.): Geological Society of America Abstracts with Programs, v. 17, no. 5, p. 293.
- Howe, Herbert J., 1986, A comparison of the brachiopod fauna in the Cape Limestone of Missouri with the standard Richmond fauna in the Ohio Valley (abs.): Geological Society of America Abstracts with Programs, v. 18, no. 4, p. 309.
- Howe, Herbert J., 1987, Origin of the brachiopod genus *Thaerodonta* and its biogeographic dispersal (abs.): Geological Society of America Abstracts with Programs, v. 19, no. 4, p. 205.
- Howe, Herbert J., 1988, Articulate brachiopods from the Richmond of Tennessee: Journal of Paleontology, v. 62, no. 2, p. 204-218.
- Huff, Warren D., 1983, Correlation of Middle Ordovician K-bentonites based on chemical fingerprinting: Journal of Geology, v. 91, no. 6, p. 657-669.
- Huff, Warren D., and Türkmenoglu, Asuman G., 1982, Chemical characteristics and origin of Ordovician K-bentonites along the Cincinnati Arch: A reply: Clays and Clay Minerals, v. 30, no. 6, p. 447-448.
- Hyde, David E., 1959, A structural and stratigraphic study of the Fairview-McMillan formational contact in the Cincinnati area: Compass, v. 36, no. 3, p. 161-171.
- International Commission on Zoological Nomenclature, 1957, Opinion 443, Designation under the Plenary Powers of a type species in harmony with accustomed usage for the genus *Monticulipora* D'Orbigny, October, 1849 (Class Bryozoa) (Ordovician): Opinions and Declarations rendered by the International Commission on Zoological Nomenclature, v. 5, pt. 3, p. 163-174.
- International Commission on Zoological Nomenclature, 1968, Opinion 838, Designation under the Plenary Powers of the type species in harmony with accustomed usage for genera *Heterotrypa* Nicholson, 1879, and *Peronopora* Nicholson, 1881 (Class Bryozoa) (Ordovician): Opinions and Declarations rendered by the International Commission on Zoological Nomenclature, v. 24, pt. 6, p. 835-841.
- International Commission on Zoological Nomenclature, 1990, *Strophomena* Blainville, 1825 (Brachiopoda): proposed adoption of authorship and designation of *Leptaena planumbona* Hall, 1847 as the type species: Bulletin of Zoological Nomenclature, v. 47, no. 1, p. 1.
- Jackson, Robert Tracy, 1929, Shaler on the fossil brachiopods of the Ohio Valley: Science, new ser., v. 70, p. 214-216.
- Jacobson, Stephen Richard, 1977, Acritarchs from Middle and Upper Ordovician rocks in New York state and the Cincinnati region in Ohio and Kentucky: Ph.D. dissertation (unpub.), The Ohio State University, 264 p.
- James, Carolyn, 1940, Species of *Tetradium* in the Cincinnati Series: American Midland Naturalist, v. 24, no. 3, p. 645-646.
- James, Jos. F., 1881, Catalogue of the fossils of the Cincinnati Group: Committee on Palaeontology, Cincinnati Society of Natural History, 27 p.
- James, Joseph F., 1884, The fucoids of the Cincinnati Group: Journal of the Cincinnati Society of Natural History, v. 7, no. 2, p. 124-132; no. 3, p. 151-166 (the date on part 2 of this paper may not be correct; an offprint of this paper, apparently inscribed by the author, has "December, 1884" marked out and "Jan 1885" written above it; U.S. Geological Survey Bulletin 746, *Geologic literature on North America, 1785-1918*, has 1884-85 noted in parentheses for this paper).
- James, Joseph F., 1885a, Evidences of beaches in the Cincinnati Group: Science, v. 5, p. 231-233.
- James, Joseph F., 1885b, Remarks on some markings on the rocks of the Cincinnati Group, described under the names of *Ormathichnus* and *Walcottia*: Journal of the Cincinnati Society of Natural History, v. 8, no. 3, p. 160-162.
- James, Joseph F., 1885c, Remarks on the genera *Lepidolites*, *Anomaloides*, *Ischadites*, and *Receptaculites* from the Cincinnati Group: Journal of the Cincinnati Society of Natural History, v. 8, no. 3, p. 163-166 (entire title is in italics in the original).
- James, Joseph F., 1886a, Cephalopoda of the Cincinnati Group: Journal of the Cincinnati Society of Natural History, v. 8, no. 4, p. 235-253, pl. 4.
- James, Joseph F., 1886b, The geology of Cincinnati: Journal of the Cincinnati Society of Natural History, v. 9, no. 2, p. 20-31.

- James, Joseph F., 1886c, Note on a recent synonym in the palaeontology of the Cincinnati Group: *Journal of the Cincinnati Society of Natural History*, v. 9, no. 2, p. 39.
- James, Joseph F., 1887, Protozoa of the Cincinnati Group: *Journal of the Cincinnati Society of Natural History*, v. 9, no. 4, p. 244-252.
- James, Joseph F., 1888a, Nomenclature of some Cincinnati Group fossils: *American Geologist*, v. 1, no. 5, p. 333.
- James, Joseph F., 1888b, *Monticulipora*, a coral and not a polyzoon: *American Geologist*, v. 1, no. 6, p. 386-392.
- James, Joseph F., 1889a, Remarks upon sedimentation in the Cincinnati Group: *Journal of the Cincinnati Society of Natural History*, v. 12, no. 1, p. 34-36.
- James, Joseph F., 1889b, On variation; with special reference to certain Palaeozoic genera: *American Naturalist*, v. 23, p. 1071-1087.
- James, Joseph F., 1891a, Manual of the paleontology of the Cincinnati Group. Part I: *Journal of the Cincinnati Society of Natural History*, v. 14, no. 1, p. 45-72.
- James, Joseph F., 1891b, Manual of the paleontology of the Cincinnati Group. Part II: *Journal of the Cincinnati Society of Natural History*, v. 14, no. 3-4, p. 149-163.
- James, Joseph F., 1891c, On the age of the Pt. Pleasant, Ohio, Beds: *Journal of the Cincinnati Society of Natural History*, v. 14, no. 2, p. 93-104, pls. 3, 4.
- James, Joseph F., 1892a, Manual of the paleontology of the Cincinnati Group. Part III: *Journal of the Cincinnati Society of Natural History*, v. 15, no. 2, p. 88-100.
- James, Joseph F., 1892b, Studies in problematic organisms—the genus *Scolithus*: *Geological Society of America Bulletin*, v. 3, no. 1, p. 32-44.
- James, Joseph F., 1893, Manual of the paleontology of the Cincinnati Group. Part IV: *Journal of the Cincinnati Society of Natural History*, v. 15, no. 3-4, p. 144-159.
- James, Joseph F., 1894, Manual of the paleontology of the Cincinnati Group. Part V: *Journal of the Cincinnati Society of Natural History*, v. 16, no. 4, p. 179-208.
- James, Joseph F., 1895, Manual of the paleontology of the Cincinnati Group. Part VI: *Journal of the Cincinnati Society of Natural History*, v. 18, no. 1-2, p. 67-88.
- James, Joseph F., 1896, Manual of the paleontology of the Cincinnati Group. Part VII: *Journal of the Cincinnati Society of Natural History*, v. 18, no. 3-4, p. 115-140.
- James, Joseph F., 1897, Manual of the paleontology of the Cincinnati Group. Part VIII: *Journal of the Cincinnati Society of Natural History*, v. 19, no. 3, p. 99-118.
- James, Uriah Pierson, 1871, Paleontology. Catalogue of the Lower Silurian fossils, Cincinnati Group, found at Cincinnati and vicinity—within a range of forty or fifty miles: Cincinnati, Ohio, U. P. James, 17 p.
- James, U. P., 1872, On a new species of fossil from the Lower Silurian: *American Journal of Science*, ser. 3, v. 3 (whole number 103), no. 13, p. 26.
- James, U. P., 1874a, Descriptions of new species of Brachiopoda, from the Lower Silurian rocks—Cincinnati Group: *Cincinnati Quarterly Journal of Science*, v. 1, no. 1, p. 19-22.
- James, U. P., 1874b, Descriptions of one new species of *Leptaena*, and two species of *Cyclonema* from the Lower Silurian rocks, Cincinnati Group: *Cincinnati Quarterly Journal of Science*, v. 1, no. 2, p. 151-153 (entire title is in italics in the original).
- James, U. P., 1874c, *Nullipores*: *Cincinnati Quarterly Journal of Science*, v. 1, no. 2, p. 153-154.
- James, U. P., 1874d, Descriptions of new species of fossils from the Lower Silurian formation, Cincinnati Group: *Cincinnati Quarterly Journal of Science*, v. 1, no. 3, p. 239-242.
- James, U. P., 1874e, Descriptions of a new species of Brachiopoda from the Lower Silurian rocks, Cincinnati Group: *Cincinnati Quarterly Journal of Science*, v. 1, no. 4, p. 333-335.
- James, U. P., 1875, Paleontology. Catalogue of Lower Silurian fossils of the Cincinnati Group found at Cincinnati and vicinity—within a circuit of 40 or 50 miles, new ed., much enl. with descriptions of some new species of corals and *Polyzoa*: Cincinnati, U. P. James, 8 p.
- James, U. P., 1878a, Descriptions of newly discovered species of fossils from the Lower Silurian formations—Cincinnati Group: *The Paleontologist*, no. 1, p. 1-8.
- James, U. P., 1878b, Descriptions of newly discovered species of fossils and remarks on others, from the Lower and Upper Silurian rocks of Ohio: *The Paleontologist*, no. 2, p. 9-15.
- James, U. P., 1879a, Description of new species of fossils and remarks on some others, from the Lower and Upper Silurian rocks of Ohio: *The Paleontologist*, no. 3, p. 17-23.
- James, U. P., 1879b, Geological nomenclature (The Cincinnati Group—The “Hudson River Group”): *The Paleontologist*, no. 4, p. 27-28.
- James, U. P., 1879c, Supplement to catalogue of the Lower Silurian fossils of the Cincinnati Group, found at Cincinnati and within a radius of forty or fifty miles: *The Paleontologist*, no. 4, p. 29-32.
- James, U. P., 1881, Contributions to paleontology, fossils of the Lower Silurian formation, Ohio, Indiana, and Kentucky: *The Paleontologist*, no. 5, p. 33-44.
- James, U. P., 1882, Descriptions of ten new species of *Monticulipora* from the Cincinnati Group, Ohio, index, etc.: *The Paleontologist*, no. 6, p. 45-56, pl. 1.
- James, U. P., 1883a, Descriptions of fossils from the Cincinnati group: *Journal of the Cincinnati Society of Natural History*, v. 6, no. 4, p. 235-236, pl. 10.
- James, U. P., 1883b, Descriptions of new species of fossils from the Cincinnati Group, Ohio and Kentucky: *The Paleontologist*, no. 7, p. 57-59, pl. 2.
- James, U. P., 1884a, Description of three species of fossils: *Journal of the Cincinnati Society of Natural History*, v. 7, no. 1, p. 20-24.
- James, U. P., 1884b, Descriptions of four new species of fossils from the Cincinnati Group: *Journal of the Cincinnati Society of Natural History*, v. 7, no. 3, p. 137-140, pl. 7.
- James, U. P., 1884c, On conodonts and fossil annelid jaws: *Journal of the Cincinnati Society of Natural History*, v. 7, no. 3, p. 143-149, pl. 7.
- James, U. P., 1885, *Glyptocrinus baeri*, Meek: *Journal of the Cincinnati Society of Natural History*, v. 8, no. 1, p. 71 (entire title is in italics in the original).
- James, U. P., 1887, Genus *Agelacinus*, Vanuxem: *Journal of the Cincinnati Society of Natural History*, v. 10, no. 1, p. 25-26 (generic name not in italics in the original and spelled as given here).
- James, U. P., and James, Joseph F., 1887, On the monticuliporoid corals of the Cincinnati Group, with critical revision of the species: *Journal of the Cincinnati Society of Natural History*, v. 10, no. 3, p. 118-141.
- James, U. P., and James, Joseph F., 1888a, On the monticuliporoid corals of the Cincinnati Group, with critical revision of the species (continued): *Journal of the Cincinnati Society of Natural History*, v. 10, no. 4, p. 158-184, pl. 2; v. 11, no. 1, p. 15-47, pl. 1.
- James, U. P., and James, Joseph F., 1888b, Monograph of the monticuliporoid corals of the Cincinnati Group: Cincinnati, Ohio, U. P. James, 47 p., 1 pl. (reprint of the series of papers published in the *Journal of the Cincinnati Society of Natural History* in 1887 and 1888).
- Janssens, Adriaan, 1967, Analysis of the Paleozoic movements of the Cincinnati Arch: Ph.D. dissertation (unpub.), The Ohio State University, 142 p.
- Jennette, David C., 1986, Storm-dominated cyclic ramp deposits of the Kope-Fairview transition (Upper Ordovician), southwestern Ohio and northern Kentucky: M.S. thesis (unpub.), University of Cincinnati, 210 p.
- Jennette, David C., and Pryor, Wayne A., 1986, Storm generated cyclic shelf deposits in Upper Ordovician of the Cincinnati Arch region (abs.): *American Association of Petroleum Geologists Bulletin*, v. 70, no. 5, p. 605.
- Jennette, David C., and Pryor, Wayne A., 1993, Cyclic alternation of proximal and distal storm facies: Kope and Fairview Formations (Upper Ordovician), Ohio and Kentucky: *Journal of Sedimentary Petrology*, v. 63, no. 2, p. 183-203.
- Jillson, Willard Rouse, 1945, The Cull Conglomerate: notes on the occurrence of an intraformational pebblestone in the Eden Shale of Owen County, Kentucky: Privately published by

- author, 8 p.
- Johnson, Thomas T., 1985, Trilobites of the Thomas T. Johnson collection: how to find, prepare, and photograph trilobites: published by author, printed by Litho-Print, Inc., Dayton, Ohio, 178 p.
- Jones, T. Rupert, 1890, On some Paleozoic Ostracoda from North America, Wales, and Ireland: *Quarterly Journal of the Geological Society of London*, v. 46, no. 1, p. 1-31, pls. 1-4.
- Kanizay, Stephen P., 1950, The Ordovician-Silurian problem in the Lanes Mill area, Oxford, Ohio: M.S. thesis (unpub.), Miami University, 31 p.
- Karklins, Olgerts L., 1981, Bryozoan biostratigraphy in Middle and Upper Ordovician rocks of Kentucky (abs.): *Geological Society of America Abstracts with Programs*, v. 13, no. 7, p. 482-483.
- Karklins, O. L., 1983a, Bryozoan fauna of the upper Clays Ferry, Kope, and lower Fairview Formations (Edenian, Upper Ordovician) at Moffett Road, northern Kentucky: U.S. Geological Survey Open-File Report 83-21, 53 p.
- Karklins, Olgerts L., 1983b, Ptilodictyoid Cryptostomata Bryozoa from the Middle and Upper Ordovician rocks of central Kentucky: *Journal of Paleontology* v. 57, no. 1, suppl. (Paleontological Society Memoir 14), 31 p.
- Karklins, O. L., 1984, Trepastome and cystoporate bryozoans from the Lexington Limestone and the Clays Ferry Formation (Middle and Upper Ordovician) of Kentucky: U.S. Geological Survey Professional Paper 1066-I, 105 p., 41 pls.
- Karklins, O. L., 1985, Bryozoan biostratigraphy of the upper Middle Ordovician and Upper Ordovician of Kentucky, in Nielsen, Claus, and Larwood, Gilbert P., eds., *Bryozoa, Ordovician to Recent: Papers presented at the 6th International Conference on Bryozoa* (Vienna, 1983), Fredensborg, Denmark, Olsen and Olsen, p. 346-347.
- Karlstrom, Adabell, 1950, A facies study of the Upper Ordovician Series: M.S. thesis (unpub.), Northwestern University.
- Kay, G. Marshall, 1929, *Rafinesquina incurvata* (Shepard), a Cincinnati brachiopod: *Geological Society of America Bulletin*, v. 40, no. 1, p. 211.
- Keith, Brian D., 1987, Tectonic influence on sedimentation patterns, Upper Ordovician of eastern North America (abs.): *American Association of Petroleum Geologists Bulletin*, v. 71, no. 9, p. 1107.
- Keith, Brian D., ed., 1989, The Trenton Group (Upper Ordovician Series) of eastern north America; deposition, diagenesis, and petroleum: *American Association of Petroleum Geologists Studies in Geology* 29, 317 p.
- Kelly, S. M., Frest, T. J., and Strimple, H. L., 1978, Additional information on *Simplococrinus persculptus*: *Journal of Paleontology*, v. 52, no. 6, p. 1227-1232, 1 pl.
- Kelly, S. M., and Pope, J. K., 1979, A new camerate crinoid from the Upper Ordovician of Indiana: *Journal of Paleontology*, v. 53, no. 2, p. 416-420, 1 pl.
- Kepferle, Roy C., Noger, Martin C., Meyer, David L., and Schumacher, Gregory A., 1987, Stratigraphy, sedimentology, and paleontology (Upper Ordovician), and glacial and engineering geology of northern Kentucky and southern Ohio: Guidebook for Annual Field Conference of the Geological Society of Kentucky, September 25-26, 1987, Kentucky Geological Survey, 18 p.
- Keroher, Grace C., and others, 1966, Lexicon of geologic names of the United States for 1936-1960: U.S. Geological Survey Bulletin 1200, 3 v., 4,341 p.
- Kerr, Duff, 1951, Stratigraphy of the Cincinnati area: *Compass*, v. 28, no. 2, p. 94-103.
- Kesling, Robert V., and Mintz, Leigh W., 1960, Internal structures in two edrioasteroid species, *Isorophus cincinnatiensis* (Roemer) and *Carneyella pilea* (Hall): *University of Michigan Museum of Paleontology Contributions*, v. 15, no. 14, p. 315-348, 9 pls.
- Kesling, Robert V., and Mintz, Leigh W., 1961, Notes on *Lepadocystis moorei* (Meek), an Upper Ordovician callocystitid cystoid: *University of Michigan Museum of Paleontology Contributions*, v. 17, no. 4, p. 123-148, 7 pls.
- Key, Marcus M., Jr., 1985, Environmental stability and morphogenic variation in the bryozoan *Homotrypa obliqua* (abs.): *Geological Society of America Abstracts with Programs*, v. 17, no. 7, p. 628.
- Key, Marcus M., Jr., 1987, Partitioning of morphologic variation across stability gradients in Upper Ordovician trepostomes, in Ross, J. R. P., ed., *Bryozoa: present and past*: Western Washington University, p. 145-152.
- Kirchner, Brian, 1991, Description, correlation, and interpretation of Cincinnati Series lithofacies in the Indiana State Route 1 roadcut near Brookville, Indiana: Independent study in geology, Earlham College, 42 p.
- Kissling, Don L., and Turonis, Linda J., 1977, Bryozoan minibioherms in the Upper Ordovician of north-central Kentucky (abs.): *Geological Society of America Abstracts with Programs*, v. 9, no. 3, p. 284-285.
- Knabe, Keith Alan, 1980, Kirkfieldian (Middle Ordovician) through Edenian (Upper Ordovician) chitinozoans from the Cincinnati region of Kentucky: M.S. thesis (unpub.), The Ohio State University, 154 p.
- Kobluk, David R., and Risk, Michael J., 1977, Algal borings and framboidal pyrite in Upper Ordovician brachiopods: *Lethaia*, v. 10, no. 2, p. 135-143.
- Kohut, Joseph James, 1967, Quantitative analysis, taxonomy, and distribution of Middle and Upper Ordovician conodonts from the Cincinnati region of Ohio, Kentucky, and Indiana: Ph.D. dissertation (unpub.), The Ohio State University, 152 p., 2 pls. (abstract: *Dissertation Abstracts* v. 28, no. 9, p. 3752B).
- Kohut, Joseph J., 1969, Determination, statistical analysis, and interpretation of recurrent conodont groups in Middle and Upper Ordovician strata of the Cincinnati region (Ohio, Kentucky, and Indiana): *Journal of Paleontology*, v. 43, no. 2, p. 392-412.
- Kohut, Joseph J., and Sweet, Walter C., 1968, The American Upper Ordovician Standard. X. Upper Maysville and Richmond conodonts from the Cincinnati region of Ohio, Indiana, and Kentucky: *Journal of Paleontology*, v. 42, no. 6, p. 1456-1477, pls. 185-186.
- Kohut, Joseph J., and Weiss, Malcolm P., 1981, Bedrock geology of the Withamsville quadrangle, Hamilton and Clermont Counties, Ohio: Ohio Division of Geological Survey Report of Investigations 120, 1 sheet with text.
- Kohut, Joseph J., Weiss, Malcolm P., and Luft, Stanley J., 1973, Geologic map of the Laurel quadrangle, Ohio-Kentucky: U.S. Geological Survey Geological Quadrangle Map GQ-1075, scale 1:24,000.
- Kolata, D. R., Brower, J. C., and Frest, T. J., 1987, Upper Mississippi Valley Champlainian and Cincinnati echinoderms: Minnesota Geological Survey Report of Investigations 35, p. 179-181.
- Kolata, Dennis R., and Guensburg, Thomas E., 1979, *Diamphidocystis*, a new mitrate "carpoid" from the Cincinnati (Upper Ordovician) Maquoketa Group in southern Illinois: *Journal of Paleontology*, v. 53, no. 5, p. 1121-1135, 1 pl.
- Kolata, D. R., Strimple, H. L., and Leverson, C. O., 1977, A new species of *Bothriocidaris* (Echinoidea) from the Cincinnati Maquoketa Group of Iowa: *Proceedings of the Iowa Academy of Science*, v. 84, no. 4, p. 146-149.
- Kolodny, Carol Renee, 1979, Structural characteristics of the bryozoan *Hallopora ramosa* (d'Orbigny) in the Dillsboro Formation (southeastern Indiana) and paleoenvironmental implications: M.S. thesis (unpub.), Boston University, 191 p.
- Kotila, David A., 1964, The bedrock geology of the Rielly area, Butler County, Ohio: M.S. thesis (unpub.), Miami University, 119 p.
- Kowalczyk, Gary R., 1976, Stratigraphy and paleoenvironment of the Cincinnati Series (Upper Ordovician) of southeast Indiana: M.S. thesis (unpub.), University of Toledo, 65 p.
- Kramer, Terry M., 1972, The morphology and taxonomy of the Upper Ordovician, Cincinnati Series, Lingulida (Brachiopoda) of Ohio, Indiana, and Kentucky: M.S. thesis (unpub.), Miami University, 116 p., 2 pls.

- Kreisa, Ronald D., 1981a, Origin of stratification in a Paleozoic epicontinental sea: the Cincinnati Series (abs.): Geological Society of America Abstracts with Programs, v. 13, no. 7, p. 491.
- Kreisa, Ronald D., 1981b, Storm-generated sedimentary structures in subtidal marine facies with examples from the Middle and Upper Ordovician of southwestern Virginia: Journal of Sedimentary Petrology, v. 51, no. 3, p. 823-843.
- Kreisa, Ronald D., Dorobek, Steven L., Accorti, Peter J., and Ginger, Elliot P., 1981, Recognition of storm-generated deposits in the Cincinnati Series, Ohio (abs.): Geological Society of America Abstracts with Programs, v. 13, no. 6, p. 285.
- Krumpolz, Bradley J., 1980, Paleocology, biostratigraphy, and sedimentology of the Mt. Auburn Member (McMillan Formation, uppermost Maysvillian) in the area of Cincinnati, Ohio: M.S. thesis (unpub.), University of Cincinnati, 196 p., 4 pls.
- Kuder, Harry B., 1964, The petrology of the limestone of the Liberty and lower Whitewater Formations (Cincinnati Series) northeastern Warren County, Ohio: M.S. thesis (unpub.), Miami University, 82 p., 13 pls.
- Laferrriere, Alan P., Hattin, Donald E., Foell, Christopher J., and Abdulkareem, Talal Faisal, 1986, The Ordovician-Silurian boundary in southeastern Indiana: Indiana Geological Survey Occasional Paper 53, 12 p.
- Lafferty, April, and Springer, Dale A., 1989, Epibionts on the rugose coral, *Grewingkia*, from the Tanners Creek Formation, (Upper Ordovician), of southeastern Indiana and southwestern Ohio (abs.): Proceedings of the Pennsylvania Academy of Science, v. 63, no. 1, p. 51 (65th annual meeting of the Pennsylvania Academy of Science and Piedmont Aquatic Scientists Association, Pocono, PA, April 7-9, 1989).
- Lamborn, Raymond E., Austin, Chester R., and Schaaf, Downs, 1938, Shales and surface clays of Ohio: Ohio Division of Geological Survey Bulletin 39, 281 p.
- Larabee, J. H., 1994, Stratigraphy and paleocology of coralline biostromes and patch reefs, Upper Ordovician of southeastern Indiana (abs.): Geological Society of America Abstracts with Programs, v. 26, no. 5, p. 49.
- La Rocque, Aurèle and Marple, Mildred Fisher, 1955, Ohio fossils: Ohio Division of Geological Survey Bulletin 54, 152 p.
- Lasemi, Zakaria, 1987, Origin of Cincinnati Series microcrystalline limestones (abs.): Ohio Journal of Science, v. 87, no. 2, p. 10.
- Lasemi, Z., and Boardman, M. R., 1989, Early cementation origin of nodular limestones, Elk Creek and Oregonia beds (Cincinnati Series), southwestern Ohio (abs.): Ohio Journal of Science, v. 89, no. 2, p. 9.
- Lattman, Laurence H., 1954, The sub-Eden beds of the Ohio Valley around Cincinnati: American Journal of Science, v. 252, no. 5, p. 257-276.
- Lee, Gerald Bernard, 1974, Lithostratigraphy of the Cincinnati Series (Upper Ordovician) from Maysville, Kentucky to Dayton, Ohio: M.S. thesis (unpub.), Miami University, 127 p.
- Lee, Gerald B., Martin, Wayne D., and Reinhart, Roy, 1974, Lithostratigraphy of the Cincinnati Series (Upper Ordovician) from Maysville, Kentucky to Dayton, Ohio (abs.): Geological Society of America Abstracts with Programs, v. 6, no. 6, p. 526.
- Leon, R. R., 1984, Provenance of Upper Ordovician (Richmondian) Bald Eagle and Juniata formations, central Pennsylvania; implications for the nature of the Taconian Orogeny in central Appalachians (abs.): American Association of Petroleum Geologists Bulletin, v. 68, no. 12, p. 1922-1923.
- Lerch, Norbert K., 1978, An inventory of Ohio soils, Butler County: Ohio Department of Natural Resources, Division of Lands and Soils, Report of Progress No. 53, 65 p. + map.
- Lerch, N. K., 1992, Soil survey of Brown County, Ohio: U.S. Department of Agriculture, Soil Conservation Service, 151 p., 65 maps + index map.
- Lerch, N. K., Hale, W. F., and Lemaster, D. D., 1982, Soil survey of Hamilton County, Ohio: U.S. Department of Agriculture, Soil Conservation Service, 219 p., 56 maps + index map.
- Lerch, Norbert K., Hale, William F., and Milliron, E. Larry, 1975, Soil survey of Clermont County, Ohio: U.S. Department of Agriculture, Soil Conservation Service, 97 p., 39 maps + index map.
- Lesquereux, Leo, 1874, On the remains of land plants in the Lower Silurian: American Journal of Science, ser. 3, v. 7, no. 37, p. 31-34 (reprinted in the Cincinnati Quarterly Journal of Science, v. 1, p. 43-45).
- Lesquereux, Leo, 1878, Land plants, recently discovered in the Silurian rocks of the United States: American Philosophical Society Proceedings, v. 17, p. 163-173.
- Linder, Robert Frederick, 1951, Detailed lithology of the carbonate members of the Elkhorn Formation in the crestal portion of the Cincinnati Arch: M.S. thesis (unpub.), Miami University, 24 p.
- Lobdell, Frederick K., 1988, Macropaleontology of the Gunn Member, Stoney Mountain Formation (Upper Ordovician), Manitoba and North Dakota: Ph.D. dissertation (unpub.), University of North Dakota, 382 p.
- Lobo, Cyril F., and Osborne, Robert H., 1973, The American Upper Ordovician Standard. XVIII. Investigation of micrite in typical Cincinnati limestones by means of scanning electron microscopy: Journal of Sedimentary Petrology, v. 43, no. 2, p. 478-483.
- Locke, John, 1838, Prof. Locke's geological report [on southwestern Ohio]: Geological Survey of Ohio Second Annual Report, p. 201-286, 15 pls.
- Locke, John, 1841, On a new species of trilobite found at Cincinnati, Ohio (abs.): American Journal of Science, v. 41, no. 1, p. 161; also published in American Geologists and Naturalists Report, 1843, p. 14.
- Locke, John, 1842a, On a new species of trilobite of very large size: American Journal of Science, v. 42, no. 2, p. 366-368, pl. 3.
- Locke, John, 1842b, On the fossil *Cryptolithus tessellatus*: Proceedings of the Academy of Natural Sciences of Philadelphia, v. 1, no. 15-16, p. 186, 196-197.
- Locke, John, 1843a, Notice of a new trilobite *Ceraurus crosotus*: American Journal of Science, v. 44, no. 2, p. 346.
- Locke, John, 1843b, Supplementary notice of the *Ceraurus crosotus*: American Journal of Science, v. 45, no. 2, p. 222-224.
- Locke, John, 1843c, On the fossil *Cryptolithus tessellatus* (continued): Proceedings of the Academy of Natural Sciences of Philadelphia, v. 1, nos. 22 & 23, p. 236.
- Loeblich, Alfred R., Jr., and Tappan, Helen, 1978, Some Middle and Late Ordovician microphytoplankton from central North America: Journal of Paleontology, v. 52, no. 6, p. 1233-1287, 16 pls.
- Lorenz, D. M., 1973, Edenian (Upper Ordovician) benthic community ecology in north-central Kentucky: M.S. thesis (unpub.), Northwestern University, 318 p.
- Ludvigsen, Rolf, and Chatterton, Brian D. E., 1982, Ordovician Pterygomtopidae (Trilobita) of North America: Canadian Journal of Earth Sciences, v. 19, no. 11, p. 2179-2206.
- Ludvigsen, R., and Tuffnell, P. A., 1983, A revision of the Ordovician olenid trilobite *Triarthrus* Green: Geological Magazine, v. 120, no. 6, p. 567-577, pls. 1-3.
- Luft, Stanley J., 1970, Geologic map of the De Mossville quadrangle, north-central Kentucky: U.S. Geological Survey Geologic Quadrangle Map GQ-862, scale 1:24,000.
- Luft, Stanley J., 1971, Geologic map of part of the Covington quadrangle, northern Kentucky: U.S. Geological Survey Geologic Quadrangle Map GQ-955, scale 1:24,000.
- Luft, Stanley J., 1972a, Geologic map of the Butler quadrangle, Pendleton and Campbell Counties, Kentucky: U.S. Geological Survey Geologic Quadrangle Map GQ-982, scale 1:24,000.
- Luft, Stanley J., 1972b, Geologic map of the Falmouth quadrangle, Pendleton County, Kentucky: U.S. Geological Survey Geologic Quadrangle Map GQ-1037, scale 1:24,000.
- Luft, Stanley J., 1973, Geologic map of the Williamstown quadrangle, Grant and Pendleton Counties, Kentucky: U.S. Geological Survey Geologic Quadrangle Map GQ-1104, scale 1:24,000.
- Luft, Stanley J., 1975, Geologic map of the Berlin quadrangle, Bracken and Pendleton Counties, Kentucky: U.S. Geological Survey Geologic Quadrangle Map GQ-1256, scale 1:24,000.
- Luft, Stanley J., Osborne, Robert H., and Weiss, Malcolm P., 1973, Geologic map of the Moscow quadrangle, Ohio-Kentucky. U.S. Geological Survey Geologic Quadrangle Map GQ-1069,

- scale 1:24,000.
- Lyke, Frederick P., 1982, External morphology of some trepostomatous ectoprocts of the Bellevue Member, McMillan Formation, (middle Upper Ordovician, southwest Ohio): M.S. thesis (unpub.), Miami University, 134 p.
- Lyell, Charles, 1845, Travels in North America; with geological observations on the United States, Canada, and Nova Scotia: London, John Murray, v. I, 316 p., v. II, 272 p. (the geology of the Cincinnati area is discussed on p. 48-51 of v. II, but see also the cross section of the eastern United States on p. 92 of v. I).
- MacDaniel, R. Patrick, 1976, Upper Ordovician sedimentary and benthic community patterns of the Cincinnati Arch area: Ph.D. dissertation (unpub.), University of Chicago, 181 p.
- MacDaniel, R. Patrick, 1976, Fossil benthic communities of the type Cincinnati Series (abs.): Geological Society of America Abstracts with Programs, v. 8, no. 4, p. 491.
- MacKenzie, P., and Bergström, S. M., 1993, Discovery of the zonal index conodont *Amorphognathus ordovicicus* in the Richmondian of Indiana: implications for the regional correlation of the North American Upper Ordovician Standard (abs.): Geological Society of America Abstracts with Programs, v. 25, no. 8, p. A-472.
- MacKenzie, P., and Bergström, S. M., 1994, Stratigraphic significance of the occurrence of *Amorphognathus ordovicicus* in the Richmondian Stage of the Cincinnati (Upper Ordovician) of Indiana (abs.): Geological Society of America Abstracts with Programs, v. 26, no. 5, p. 51.
- Mahan, Thomas Kent, 1981, Variations in the depositional environment of the lower Cincinnati Kope Formation: M.S. thesis (unpub.), Western Michigan University, 158 p.
- Mahan, T. K., and Harrison, W. B., III, 1980, Depositional environment of lower Cincinnati Kope Formation and some paleoecologic implications (abs.): American Association of Petroleum Geologists Bulletin, v. 64, no. 5, p. 744.
- Marak, Joe H., ed., 1992a, The Richmond Group of the Cincinnati Province, an unpublished manuscript by Dr. William Henry Shideler (circa 1927): Department of Geology, Miami University, 94 p. (available as Ohio Division of Geological Survey Open-File Report 95-1).
- Marak, Joe H., 1992b, Fossil collecting localities in the Oxford area: The Karl E. Limper Geology Museum, Department of Geology, Miami University, Oxford, Ohio, two sheets, folded.
- Marsh, O. C., 1867, Notice of a new genus of fossil sponges from the Lower Silurian: American Journal of Science and Arts, series 2, v. 44, no. 130, p. 88.
- Martin, A. J., 1986a, Burial preservation of trace fossils as indicator of storm deposition (abs.): American Association of Petroleum Geologists Bulletin, v. 70, no. 5, p. 616.
- Martin, Anthony J., 1986b, A paleoenvironmental interpretation of the "Arnheim" micromorph fossil assemblage from the Cincinnati Series (Upper Ordovician), southeastern Indiana and southwestern Ohio: M.S. thesis (unpub.), Miami University, 221 p.
- Martin, Anthony J., and Pope, John K., 1985, A paleoenvironmental interpretation of a micromorph fossil assemblage from the Cincinnati Series (Upper Ordovician), Indiana and Ohio (abs.): Geological Society of America Abstracts with Programs, v. 17, no. 7, p. 653.
- Martin, Anthony J., Martin, Wayne D., and Pope, John K., 1985, Genesis of phosphatic sediments in Cincinnati Series (Upper Ordovician), southeastern Indiana and southwestern Ohio (abs.): American Association of Petroleum Geologists Bulletin, v. 69, no. 9, p. 1441.
- Martin, Wayne D., 1971, Petrography of composite vertical section of Cincinnati Series limestones, southwestern Ohio and adjacent areas (abs.): American Association of Petroleum Geologists Bulletin, v. 55, no. 2, p. 351.
- Martin, Wayne D., 1973, Textural characteristics and faunal components of the Cincinnati Series limestones of southwestern Ohio and adjacent areas indicating the origin of these rocks (abs.): Geological Society of America Abstracts with Programs, v. 5, no. 5, p. 416-417.
- Martin, Wayne D., 1975, The petrology of a composite vertical section of Cincinnati Series limestones (Upper Ordovician) of southwestern Ohio, southeastern Indiana, and northern Kentucky: Journal of Sedimentology Petrology, v. 45, no. 4, p. 907-925.
- Martin, Wayne D., 1977, Petrology of the Cincinnati Series limestones (Upper Ordovician) of Indiana and Ohio, in Pope, John K., and Martin, Wayne D., eds., Field guidebook to the biostratigraphy and paleoenvironments of the Cincinnati Series in southeastern Indiana: Society of Economic Paleontologists and Mineralogists, Great Lakes Section, 7th Annual Field Conference, Miami University, p. III-II25 (reprinted in 1979 with four additional pages; see Pope and Martin, 1977).
- Martin, Wayne, and Shideler, W. H., 1958, Dry Dredgers excursion to Oxford, Ohio area, April 13, 1958: Department of Geology, Miami University, 11 unnumbered pages, mimeographed, no "publication" data (copy in Geology/Physics Library, University of Cincinnati). See also [National] Association of Geology Teachers [East-Central Section], 1958.
- Maurer, Robert James, 1974, The classification of bedding contacts in the Richmondian Stage (Upper Ordovician) of the Cincinnati Series: M.S. thesis (unpub.), Miami University, 149 p.
- McAuley, Robert J., Elias, Robert J., and Mattison, Blair W., 1986, Succession of solitary rugose corals across the Ordovician-Silurian boundary, east-central United States (abs.): Proceedings of the 4th North American Paleontological Convention, p. A31.
- McClish, Richard Fred, 1965, Lithostratigraphy and conodont biostratigraphy of the Richmond Group of southwestern Ohio and southeastern Indiana: M.S. thesis (unpub.), The Ohio State University, 84 p., 9 pls.
- McCormick, Charles Larry, 1960, Petrology of the Waynesville Formation (Cincinnati Series) in the Fort Ancient-Oregonia region, Warren County, Ohio: M.S. thesis (unpub.), Miami University, 89 p., 9 pls.
- McCracken, Alexander D., and Nowlan, Godfrey S., 1988, The Gamachian Stage and Fauna 13, in Landing, Ed, ed., The Canadian Paleontology and Biostratigraphy Seminar, Albany, NY, Sept. 29 - Oct. 1, 1986: New York State Museum Bulletin 462, p. 71-79.
- McDowell, Robert C., 1981, Correlation chart for units on the geologic map of Kentucky: U.S. Geological Survey Miscellaneous Field Studies Map MF-1291, 1 sheet.
- McDowell, R. C., 1983, Stratigraphy of the Silurian outcrop belt on the east side of the Cincinnati Arch in Kentucky, with revisions in the nomenclature: U.S. Geological Survey Professional Paper 1151-F, 27 p., 2 pls.
- McEwan, Eula Davis, 1919, A study of the brachiopod genus *Platystrophia*: Proceedings of the U.S. National Museum, v. 56, p. 383-448, 11 pls. Also see review by Bather, 1920.
- McEwan, Eula Davis, 1920, The Ordovician of Madison, Indiana: American Journal of Science, ser. 4, v. 50, p. 154-158.
- McFarlan, A. C., 1913, Geology of the Georgetown quadrangle: Kentucky Geological Survey Report, ser. IV, v. 1, pt. 1, p. 317-351, pls. 1-10.
- McFarlan, Arthur C., 1919, The geology of Kentucky: Ordovician System: Kentucky Department of Geology and Forestry Bulletin, ser. 5, no. 2, p. 16-53, pls. 20-23.
- McFarlan, Arthur C., 1931, The Ordovician fauna of Kentucky, in Jilison, Willard Rouse, ed., Paleontology of Kentucky: Kentucky Geological Survey Report, ser. VI, v. 36, p. 47-166, 16 pls.
- McFarlan, A. C., 1943, Geology of Kentucky: Lexington, University of Kentucky Press, 531 p.
- McKinney, Frank K., 1969, Bibliography and list of the families Constellariidae and Dianulitidae (Ectoprocta, Order Cystopora): Southeastern Geology, v. 10, no. 3, p. 175-184.
- McKinney, Frank K., 1974, Bibliography and catalogue (1900-1969) of the Trepostomata (phylum Ectoprocta): Southeastern Geology, Special Publication 4, 147 p.
- McKinney, F. K., 1975, Autozoecial budding patterns in dendroid stenolaemate bryozoans, in David, L., ed., Bryozoan 1974: Documents Laboratoires Geologie Faculté des Sciences Lyon, Hors Ser. 3, fasc. 1, p. 65-76, 4 pls.

- McKinney, Frank K., 1977, Autozoecial budding patterns in dendroid Paleozoic bryozoans: *Journal of Paleontology*, v. 51, no. 2, p. 303-329, 9 pls.
- Meek, F. B., 1871, On some new Silurian crinoids and shells: *American Journal of Science*, ser. 3, v. 2 (whole number 102), no. 10, p. 295-302.
- Meek, F. B., 1872a, Supplementary note on the genus *Lichenocrinus*: *American Journal of Science*, ser. 3, v. 3 (whole number 103), no. 13, p. 15-17 (entire title is in italics in the original).
- Meek, F. B., 1872b, Descriptions of two new star-fishes, and a crinoid, from the Cincinnati group of Ohio and Indiana: *American Journal of Science*, ser. 3, v. 3 (whole number 103), no. 16, p. 257-262.
- Meek, F. B., 1872c, Descriptions of new species of fossils from the Cincinnati Group of Ohio: *American Journal of Science*, ser. 3, v. 3 (whole number 103), no. 18, p. 423-428.
- Meek, F. B., 1872d, Descriptions of a few new species, and one new genus of Silurian fossils, from Ohio: *American Journal of Science*, ser. 3, v. 4 (whole number 104), no. 22, p. 274-281.
- Meek, F. B., 1872e, Descriptions of new western Paleozoic fossils, mainly from the Cincinnati Group of the Lower Silurian Series of Ohio: *Proceedings of the Academy of Natural Sciences of Philadelphia*, v. 21, pt. 3, p. 308-336.
- Meek, F. B., 1873, Descriptions of invertebrate fossils of the Silurian and Devonian systems: *Geological Survey of Ohio*, v. 1, pt. 2, p. 1-246, pls. 1-23.
- Meek, F. B., and Worthen, A. H., 1865a, Descriptions of new species of Crinoidea, etc., from the Paleozoic rocks of Illinois and some of the adjoining states: *Proceedings of the Academy of Natural Sciences of Philadelphia*, v. 17, p. 143-155.
- Meek, F. B., and Worthen, A. H., 1865b, Contributions to the paleontology of Illinois and other western states: *Proceedings of the Academy of Natural Sciences of Philadelphia*, v. 17, p. 245-273.
- Melchin, M., and Legault, J. A., 1985, Evolutionary lineages in some Ordovician Chitinozoa: *Palynology*, v. 9, p. 199-210.
- Meyer, David L., 1988, Population paleoecology of edrioasteroid echinoderms from the Upper Ordovician of the Cincinnati Arch region (abs.): *Geological Society of America Abstracts with Programs*, v. 20, no. 1, p. 56.
- Meyer, David L., 1990, Population paleoecology and comparative taphonomy of two edrioasteroid (Echinodermata) pavements: Upper Ordovician of Kentucky and Ohio: *Historical Biology*, v. 4, no. 3, p. 155-178.
- Meyer, D. L., Diekmeyer, S. D., and Holterhoff, P. F., 1990, Crinoid distribution in shoaling-upward carbonate-clastic sequences: Upper Ordovician (Maysvillian) of Ohio and Kentucky, U.S.A. (abs.): Atami, Japan, Seventh International Echinoderm Conference, Abstracts with Programs, p. 29.
- Meyer, David L., Schumacher, Gregory A., Swinford, E. Mac, Jennette, David C., and Brockman, C. Scott, 1985, Upper Ordovician stratigraphy, sedimentology, and paleontology along Backbone Creek, Clermont County, Ohio: Field trip guidebook for Annual Meeting of Ohio Academy of Science, Geology Section, 14 p.
- Meyer, David L., Tobin, Rick C., Pryor, Wayne A., Harrison, William B., Osgood, Richard G., Hinterlong, Gregory D., Krumpolz, Bradley J., and Mahan, Thomas K., 1981, Stratigraphy, sedimentology, and paleoecology of the Cincinnati Series (Upper Ordovician) in the vicinity of Cincinnati, Ohio, in Roberts, Thomas G., ed., *GSA Cincinnati '81 Field Trip Guidebooks*: *American Geological Institute*, v. 1, p. 31-72.
- Meyer, David L., and Weaver, Thomas R., 1980, Biostratigraphy of crinoid-dominated communities in the lower Bull Fork Formation (Upper Ordovician) of southwestern Ohio (abs.): *Geological Society of America Abstracts with Programs*, v. 12, no. 5, p. 251.
- Mickleborough, John, 1883, Locomotory appendages of trilobites: *Journal of the Cincinnati Society of Natural History*, v. 6, no. 3, p. 200-206.
- Mickleborough, John, and Wetherby, A. G., 1878a, A classified list of Lower Silurian fossils, Cincinnati Group: *Journal of the Cincinnati Society of Natural History*, v. 1, no. 2, p. 61-86.
- Mickleborough, J., and Wetherby, A. G., 1878b, A classified list of Lower Silurian fossils, Cincinnati Group: Cincinnati, Ohio, James Barclay, 26 p.
- Miller, Arnold I., 1989, Spatio-temporal transitions in Paleozoic Bivalvia: a field comparison of Upper Ordovician and Upper Paleozoic bivalve-dominated fossil assemblages: *Historical Biology*, v. 2, no. 3, p. 227-260.
- Miller, Merrell Allen, 1976, Maysvillian (Upper Ordovician) chitinozoans from the Cincinnati region in Ohio, Kentucky, and Indiana: M.S. thesis (unpub.), The Ohio State University, 251 p.
- Miller, M. A., 1980, Paleoecological significance of chitinozoan recurrent species associations (Upper Ordovician), Cincinnati region, U.S.A. (abs.): Abstracts of the Fifth International Palynological Conference, Cambridge, 1980, p. 262.
- Miller, M. A., and Wicander, R., 1982, Upper Ordovician (lower Richmondian) acritarchs and chitinozoans from southeastern Indiana, U.S.A. (abs.): *Palynology*, v. 6, p. 286-287.
- Miller, S. A., 1874a, The column of *Heterocrinus heterodactylus*: *Cincinnati Quarterly Journal of Science*, v. 1, no. 1, p. 2-3.
- Miller, S. A., 1874b, *Pasceolus Darwini*: *Cincinnati Quarterly Journal of Science*, v. 1, no. 1, p. 5-6.
- Miller, S. A., 1874c, *Pasceolus Claudi*: *Cincinnati Quarterly Journal of Science*, v. 1, no. 1, p. 6-7.
- Miller, S. A., 1874d, Remarks on the genus *Conchicolites* of Nicholson: *Cincinnati Quarterly Journal of Science*, v. 1, no. 1, p. 7-11 (all words of title except "genus" in italics in the original).
- Miller, S. A., 1874e, Genus *Ambonychia* (Hall): *Cincinnati Quarterly Journal of Science*, v. 1, no. 1, p. 14-16.
- Miller, S. A., 1874f, Genus *Anomalodonta*: *Cincinnati Quarterly Journal of Science*, v. 1, no. 1, p. 16-18.
- Miller, S. A., 1874g, The Cincinnati Group: *Cincinnati Quarterly Journal of Science*, v. 1, no. 1, p. 63-64.
- Miller, S. A., 1874h, The position of the Cincinnati Group in the geological column of fossiliferous rocks of North America: *Cincinnati Quarterly Journal of Science*, v. 1, no. 2, p. 97-115.
- Miller, S. A., 1874i, Monograph of the Crustacea of the Cincinnati Group: *Cincinnati Quarterly Journal of Science*, v. 1, no. 2, p. 115-147.
- Miller, S. A., 1874j, *Cypricardites Hainesi*: *Cincinnati Quarterly Journal of Science*, v. 1, no. 2, p. 147-148.
- Miller, S. A., 1874k, *Streptorhynchus* (?) *Hallie*: *Cincinnati Quarterly Journal of Science*, v. 1, no. 2, p. 148-149.
- Miller, S. A., 1874l, *Modiolopsis modiolaris*—(Conrad, 1838): *Cincinnati Quarterly Journal of Science*, v. 1, no. 2, p. 149-150.
- Miller, S. A., 1874m, *Modiolopsis Versaillesensis*: *Cincinnati Quarterly Journal of Science*, v. 1, no. 2, p. 150.
- Miller, S. A., 1874n, Monograph of the Lamellibranchiata of the Cincinnati Group: *Cincinnati Quarterly Journal of Science*, v. 1, no. 3, p. 211-231 (entire title in italics in the original).
- Miller, S. A., 1874o, *Cyrtoceras Vallandighamii*: *Cincinnati Quarterly Journal of Science*, v. 1, no. 3, p. 232.
- Miller, S. A., 1874p, *Beyrichia Duryi*: *Cincinnati Quarterly Journal of Science*, v. 1, no. 3, p. 232-233.
- Miller, S. A., 1874q, *Beyrichia striato-marginatus*: *Cincinnati Quarterly Journal of Science*, v. 1, no. 3, p. 233.
- Miller, S. A., 1874r, *Beyrichia Chambersi*: *Cincinnati Quarterly Journal of Science*, v. 1, no. 3, p. 234.
- Miller, S. A., 1874s, *Tentaculites Richmondensis*: *Cincinnati Quarterly Journal of Science*, v. 1, no. 3, p. 234-235.
- Miller, S. A., 1874t, *Buthotrephis ramulosus*: *Cincinnati Quarterly Journal of Science*, v. 1, no. 3, p. 235-236.
- Miller, S. A., 1874u, Letter from Professor Nicholson on the genera *Conchicolites* and *Ortonia*: *Cincinnati Quarterly Journal of Science*, v. 1, no. 3, p. 236-239 (the entire title is italicized in the original).
- Miller, S. A., 1874v, *Modiolopsis pholadiformis* (Foster and Whitney): *Cincinnati Quarterly Journal of Science*, v. 1, no. 3, p. 282.
- Miller, S. A., 1874w, Monograph of the Gasteropoda of the Cincinnati Group: *Cincinnati Quarterly Journal of Science*, v. 1, no. 4, p. 302-321.
- Miller, S. A., 1874x, Remarks upon the genus *Anomalodonta* and the

- words *Megaptera* and *Opisthoptera* and the species *gigantea* and *alata*: Cincinnati Quarterly Journal of Science, v. 1, no. 4, p. 326-333.
- Miller, S. A., 1874y, Genus *Megalograptus*: Cincinnati Quarterly Journal of Science, v. 1, no. 4, p. 343-346.
- Miller, S. A., 1874z, *Lichenocrinus tuberculatus*: Cincinnati Quarterly Journal of Science, v. 1, no. 4, p. 346-347.
- Miller, S. A., 1874aa, *Trematis Dyeri*: Cincinnati Quarterly Journal of Science, v. 1, no. 4, p. 347.
- Miller, S. A., 1874bb, *Beyrichia Richardsoni*: Cincinnati Quarterly Journal of Science, v. 1, no. 4, p. 347-348.
- Miller, S. A., 1874cc, *Glyptocrinus Fornshelli*: Cincinnati Quarterly Journal of Science, v. 1, no. 4, p. 348-351.
- Miller, S. A., 1874dd, Observations upon *Stenopora fibrosa* and the genus *Chetetes*: Cincinnati Quarterly Journal of Science, v. 1, no. 4, p. 368-375.
- Miller, S. A., 1875a, Monograph of the class Brachiopoda of the Cincinnati Group: Cincinnati Quarterly Journal of Science, v. 2, no. 1, p. 6-62.
- Miller, S. A., 1875b, *Acidaspis O'nealli*: Cincinnati Quarterly Journal of Science, v. 2, no. 1, p. 86-87.
- Miller, S. A., 1875c, Class *Cephalopoda* (Cuvier), as represented in the Cincinnati Group: Cincinnati Quarterly Journal of Science, v. 2, no. 2, p. 121-134 (italics as in the original).
- Miller, S. A., 1875d, Some remarks on *Plumulites Jamesi* of Hall and Whitfield: Cincinnati Quarterly Journal of Science, v. 2, no. 3, p. 274-277.
- Miller, S. A., 1875e, *Glyptocrinus Shafferi*: Cincinnati Quarterly Journal of Science, v. 2, no. 3, p. 277-279.
- Miller, S. A., 1875f, *Heterocrinus isodactylus*: Cincinnati Quarterly Journal of Science, v. 2, no. 3, p. 279.
- Miller, S. A., 1875g, *Crania reticularis*: Cincinnati Quarterly Journal of Science, v. 2, no. 3, p. 280.
- Miller, S. A., 1875h, Some further remarks upon the genus *Anomalodonta*: Cincinnati Quarterly Journal of Science, v. 2, no. 3, p. 280-284.
- Miller, S. A., 1875i, *Cyrtoceras magister*: Cincinnati Quarterly Journal of Science, v. 2, no. 3, p. 284.
- Miller, S. A., 1875j, Some new species of fossils from the Cincinnati Group and remarks upon some described forms: Cincinnati Quarterly Journal of Science, v. 2, no. 4, p. 349-355.
- Miller, S. A., 1875k, The square crinoid column: Cincinnati Quarterly Journal of Science, v. 2, no. 4, p. 378-379.
- Miller, S. A., 1878, Description of a new genus and eleven new species of fossils, with remarks upon others well known, from the Cincinnati Group: Journal of the Cincinnati Society of Natural History, v. 1, no. 2, p. 100-108, pl. 3.
- Miller, S. A., 1879a, Catalogue of fossils found in the Hudson River, Utica Slate, and Trenton Groups, as exposed in the southeast part of Indiana, southwest part of Ohio, and northern part of Kentucky: Indiana Geological Survey Annual Report for 1876-77-78, v. 8-9-10, p. 22-56 (reprint edition paginated p. 1-35).
- Miller, S. A., 1879b, Description of twelve new fossil species (Niagaran and Cincinnatian) and remarks upon others: Journal of the Cincinnati Society of Natural History, v. 2, no. 2, p. 104-118, pls. 9-10.
- Miller, S. A., 1880a, Silurian [Ordovician] ichnolites, with definitions of new genera and species: Journal of the Cincinnati Society of Natural History, v. 2, no. 4, p. 217-222, pls. 13-14.
- Miller, S. A., 1880b, Note upon the habits of some fossil annelids: Journal of the Cincinnati Society of Natural History, v. 2, no. 4, p. 260.
- Miller, S. A., 1880c, Description of four new species of Silurian fossils: Journal of the Cincinnati Society of Natural History, v. 3, no. 2, p. 140-144, pl. 4 (reprint paginated p. 1-5).
- Miller, S. A., 1880d, Description of four new species and a new variety of Silurian fossils, and remarks upon others: Journal of the Cincinnati Society of Natural History, v. 3, no. 3, p. 232-236, pl. 7.
- Miller, S. A., 1881a, Description of five new species of Silurian fossils, and remarks upon an undetermined form: Journal of the Cincinnati Society of Natural History, v. 3, no. 4, p. 314-317, pl. 8.
- Miller, S. A., 1881b, Description of some new and remarkable crinoids and other fossils of the Hudson River Group, and notice of *Strotocrinus bloomfieldensis*: Journal of the Cincinnati Society of Natural History, v. 4, no. 1, p. 69-77, pl. 1.
- Miller, S. A., 1881c, Description of new species of fossils from the Hudson River group, with remarks upon others: Journal of the Cincinnati Society of Natural History, v. 4, no. 4, p. 316-319, pl. 8.
- Miller, S. A., 1882a, Notice of a work by Prof. Nicholson on the genus *Monticulipora*: Journal of the Cincinnati Society of Natural History, v. 5, no. 1, p. 25-33.
- Miller, S. A., 1882b, Description of two new genera and eight new species of fossils from the Hudson River group, with remarks upon others: Journal of the Cincinnati Society of Natural History, v. 5, no. 1, p. 34-44, pls. 1-2.
- Miller, S. A., 1882c, Description of ten new species of fossils: Journal of the Cincinnati Society of Natural History, v. 5, no. 2, p. 79-88, pls. 3-4.
- Miller, S. A., 1882d, Description of three new species and remarks upon others: Journal of the Cincinnati Society of Natural History, v. 5, no. 3, p. 116-117, pl. 5.
- Miller, S. A., 1882e, Description of three new orders and four new families in the class Echinodermata and eight new species from the Silurian and Devonian formations: Journal of the Cincinnati Society of Natural History, v. 5, no. 4, p. 221-231, pl. 9.
- Miller, S. A., 1883, *Glyptocrinus* redefined and restricted, *Gaurocrinus*, *Pycnocrinus* and *Compsocrinus* established, and two new species described: Journal of the Cincinnati Society of Natural History, v. 6, no. 4, p. 217-234, pl. 11.
- Miller, S. A., 1884, Description of a beautiful starfish and other fossils: Journal of the Cincinnati Society of Natural History, v. 7, no. 1, p. 16-20, pl. 4.
- Miller, S. A., 1889, North American geology and palaeontology for the use of amateurs, students, and scientists: Cincinnati, Ohio, [Western Methodist book concern], 664 p.
- Miller, S. A., 1892, Palaeontology: Indiana Department of Geology and Natural Resources, 17th Annual Report, p. 611-705, 20 pls.
- Miller, S. A., 1897, Second appendix to North American geology and palaeontology: Cincinnati, Ohio, published by the author, p. 719-793.
- Miller, S. A., Braun, Fred., Mickleborough, Jno., Hall, John W., Jr., Ulrich, E. O., Wetherby, A. G., Harper, Geo. W., Mohr, Paul, Dyer, C. B., and Byrnes, R. M., 1879, Report of committee on geological nomenclature: Journal of the Cincinnati Society of Natural History, v. 1, no. 4, p. 193-194.
- Miller, S. A., and Dyer, C. B., 1878a, Contributions to palaeontology: Journal of the Cincinnati Society of Natural History, v. 1, no. 1, p. 24-39, pls. 1-2 (reprint paginated p. 1-16).
- Miller, S. A., and Dyer, C. B., 1878b, Contributions to paleontology. No. 2: Cincinnati, Ohio, published by the authors, p. 1-11, pls. 3-4.
- Miller, S. A., and Faber, Charles, 1892, Some new species and new structural parts of fossils: Journal of the Cincinnati Society of Natural History, v. 15, no. 2, p. 79-87.
- Miller, S. A., and Faber, C. L., 1894a, New species of fossils from the Hudson River Group, and remarks upon others: Journal of the Cincinnati Society of Natural History, v. 17, no. 1, p. 22-33, pl. 1.
- Miller, S. A., and Faber, C. L., 1894b, Description of some Cincinnati fossils: Journal of the Cincinnati Society of Natural History, v. 17, no. 3, p. 137-158, pls. 7-8.
- Mitchell, Charles E., 1986, Morphometric studies of *Climacograptus* (Hall) and the phylogenetic significance of astogeny, in Hughes, C. P., and Rickards, R. B., eds., Palaeoecology and biostratigraphy of graptolites: Geological Society Special Publication 20, p. 119-129.
- Mitchell, C. E., and Bergström, S. M., 1977, Three-dimensionally preserved Richmondian graptolites from southwestern Ohio and the graptolite correlation of the North American Upper Ordovician Standard: Bollettino della Società Paleontologica Italiana, v. 16, no. 2, p. 257-270.
- Mitchell, Charles E., and Bergström, Stig M., 1987, A new and stratigraphically significant graptolite succession across the

- base of the Cincinnati (Upper Ordovician) in eastern Indiana (abs.): Geological Society of America Abstracts with Programs, v. 19, no. 4, p. 235.
- Mitchell, Charles E., and Bergström, Stig M., 1988, Definition and correlation of the base of the Cincinnati, the North American Upper Ordovician Standard Series: graptolite, conodont, and lithostratigraphic evidence from the reference standard in Ohio, Indiana, and Kentucky (abs.): Fifth International Symposium on the Ordovician System, St. John's, Newfoundland, 1988, Program and Abstracts, p. 61.
- Mitchell, Charles E., and Bergström, Stig M., 1991, New graptolite and lithostratigraphic evidence from the Cincinnati region, U.S.A., for the definition and correlation of the base of the Cincinnati Series (Upper Ordovician), in Barnes, Christopher R., and Williams, S. Henry, eds., Advances in Ordovician Geology: Geological Survey of Canada Paper 90-9, p. 59-77, 1 pl.
- Mitchell, Charles E., Bergström, Stig M., and Schumacher, Gregory A., 1991, The Sebree Trough Project. 5. Development and possible tectonic origin of a Middle to Late Ordovician intracratonic basin (abs.): Geological Society of America Abstracts with Programs, v. 23, no. 3, p. 49.
- Mitchell, Charles E., Wilson, Mark A., and St. John, James M., 1993, In situ crustoid graptolite colonies from an Upper Ordovician hardground, southwestern Ohio: Journal of Paleontology v. 67, no. 6, p. 1011-1016.
- Morris, Robert W., and Felton, Stephen H., 1993, Symbiotic association of crinoids, platyceratid gastropods, and *Cornulites* in the Upper Ordovician (Cincinnati) of the Cincinnati, Ohio region: Palaios, v. 8, no. 5, p. 465-476.
- Morris, Robert W., and Rollins, Harold B., 1971, The distribution and paleoecological interpretation of *Cornulites* in the Waynesville Formation (Upper Ordovician) of southwestern Ohio: Ohio Journal of Science, v. 71, no. 3, p. 159-170.
- Morrison, Stanley J., and Anstey, Robert L., 1979, Ultrastructure and composition of brown bodies in some Ordovician trepostome bryozoans: Journal of Paleontology, v. 53, no. 4, p. 943-949.
- [National] Association of Geology Teachers [East-Central Section], 1958, Ordovician stratigraphy of the Oxford, Ohio area: Field Trip, 22 March 1958, Department of Geology, Miami University, 22 p. (mimeographed, using some of the same pages as Martin, Wayne, and Shideler, W. H., 1958; copy in Geology/Physics Library, University of Cincinnati).
- Neuman, Robert B., 1967, Some silicified Middle Ordovician brachiopods from Kentucky: U.S. Geological Survey Professional Paper 583-A, 14 p., 2 pls.
- Newberry, J. S., 1873, The general geological relations and structure of Ohio: Geological Survey of Ohio, v. 1, pt. 1, p. 1-167.
- Newberry, J. S., 1874, On the so-called land plants from the Lower Silurian of Ohio: American Journal of Science and the Arts, 3rd series, v. 8 (whole number 108), no. 44 (but given as 64), p. 110-113 (republished without the figures in Cincinnati Quarterly Journal of Science, 1874, v. 1, no. 4, p. 335-338).
- Nicholson, H. Alleyne, 1872, On *Ortonia*, a new genus of fossil tubicolar Annelides, with notes on the genus *Tentaculites*: Geological Magazine, v. 9, no. 100, p. 446-449.
- Nicholson, H. Alleyne, 1873, Descriptions of two new species of fossil tubicolar Annelides: Geological Magazine, v. 10, no. 104, p. 54-57, pl. 4.
- Nicholson, H. Alleyne, 1874a, Description of species of *Chaetetes* from the Lower Silurian rocks of North America: Quarterly Journal of the Geological Society of London, v. 30, p. 499-515, pls. 29-30.
- Nicholson, H. A., 1874b, On *Columnopora*, a new genus of tabulate corals: Geological Magazine, 2nd ser., v. 1, p. 253-254.
- Nicholson, H. Alleyne, 1875a, Description of the corals of the Silurian and Devonian systems: Geological Survey of Ohio, v. 2, pt. 2, p. 181-242, pls. 21-23.
- Nicholson, H. Alleyne, 1875b, Descriptions of Polyzoa from the Silurian formation: Geological Survey of Ohio, v. 2, pt. 2, Geology, p. 257-268, pl. 25.
- Nicholson, H. A., 1875c, Descriptions of species of *Hippothoa* and *Alecto* from the Lower Silurian rocks of Ohio, with a description of *Aulopora arachnoidea* Hall: Annals and Magazine of Natural History, ser. 4, v. 15, no. 86, p. 123-127.
- Nicholson, H. A., 1875d, Descriptions of new species of Polyzoa from the Lower and Upper Silurian rocks of North America: Annals and Magazine of Natural History, ser. 4, v. 15, no. 87, p. 177-184.
- Nicholson, H. A., 1876, Notes on the Palaeozoic corals of the state of Ohio: Annals and Magazine of Natural History, ser. 4, v. 18, p. 85-95, pl. 5.
- Nicholson, H. Alleyne, 1879, On the structure and affinities of the "tabulate corals" of the Paleozoic Period, with critical descriptions of illustrative species: Edinburgh, William Blackwood and Sons, 342 p., 15 pls.
- Nicholson, H. A., 1880, On the minute structure of the recent *Heteropora neozelanica* Busk and on the relations of the genus *Heteropora* to *Monticulipora*: Annals and Magazine of Natural History, ser. 5, v. 6, no. 35, p. 329-339; no. 36, p. 414-423.
- Nicholson, H. Alleyne, 1881, On the structure and affinities of the genus *Monticulipora* and its sub-genera, with critical descriptions of illustrative species: Edinburgh, William Blackwood and Sons, 240 p., 6 pls.
- Nickles, J. M., 1902a, The geology of Cincinnati: Journal of the Cincinnati Society of Natural History, v. 20, no. 2, p. 49-100, 1 pl. [2-page map]; reprint edition, p. 1-53, 1 pl.
- Nickles, John M., 1902b, Description of a new bryozoan, "*Homotrypa bassleri*" n. sp., from the Warren beds of the Lorraine Group: Journal of the Cincinnati Society of Natural History, v. 20, no. 2, p. 103-105 (no italics in the original).
- Nickles, John M., 1903, The Richmond Group in Ohio and Indiana and its subdivisions, with a note on the genus *Strophomena* and its type: American Geologist, v. 32, no. 4, p. 202-218.
- Nickles, John M., 1905, The Upper Ordovician rocks of Kentucky and their Bryozoa: Kentucky Geological Survey, Series III, Bulletin 5, 64 p., 3 pls.
- Nickles, John M., and Bassler, Ray S., 1900, A synopsis of American fossil Bryozoa, including bibliography and synonymy: U.S. Geological Survey Bulletin 173, 663 p.
- Norman, Carl Edgar, 1959, Stratigraphy and petrology of the Upper Eden strata in the Ohio Valley: M.S. thesis (unpub.), The Ohio State University, 133 p., 4 pls.
- Norrish, Winston, 1982, Petrographic analysis of *Diplocraterion* ichnofossils from the Cincinnati Group (Ordovician) (abs.): Ohio Journal of Science, v. 82, no. 2, p. 26.
- Norton, L. D., 1979, Soils developed from the Kope geologic formation: Ohio Journal of Science, v. 79, no. 4, p. 160-163.
- Nuttall, Thomas, 1821, A journal of travels into Arkansas Territory, during the year 1819, with occasional observations on the manners of the aborigines: Philadelphia, Thos. M. Palmer, xii + 296 p., including two-page map; reprinted 1966, Readex Microprint Corporation (section that includes Cincinnati begins on p. 30; fossils mentioned, including *Trilobites paradoxus*, on p. 31).
- O'Brien, John M., 1970, The stratigraphy of the upper part of the Richmond Group (Ordovician) of Indiana, Kentucky, and Ohio: M.S. thesis (unpub.), Miami University, 155 p.
- Oldroyd, John David, 1978, Paleocommunities and environments of the Dillsboro and Saluda formations (Upper Ordovician), Madison, Indiana: Ph.D. dissertation (unpub.), University of Michigan, 340 p.
- Ordovician Subcommittee of the Committee on Stratigraphy of the National Research Council, W. H. Twenhofel, Chairman, 1954, Correlation of the Ordovician formations of North America: Geological Society of America Bulletin, v. 65, no. 3, p. 247-298, 1 pl.
- Orton, Edward, 1873, Report on the Third Geological District: geology of the Cincinnati Group; Hamilton, Clermont, Warren and Butler Counties: Geological Survey of Ohio, v. 1, pt. 1, Geology, p. 365-480.
- Orton, Edward, 1878, Report on the geology of Warren County; Butler County; Preble County; Madison County: Geological Survey of Ohio, v. 3, pt. 1 [no other parts were published], p. 381-428.

- Orton, Edward, 1890, Geological scale and geological structure of Ohio: Geological Survey of Ohio, Third Organization, First Annual Report, p. 9-54.
- Orton, Edward, Jr., and Peppel, Samuel Vernon, 1906, The limestone resources and lime industry in Ohio: Ohio Division of Geological Survey Bulletin 4, 365 p.
- Osborne, Robert H., 1966, Bedrock geology and limestone petrology of eastern Hamilton County, Ohio: Ph.D. dissertation (unpub.), The Ohio State University, 143 p.
- Osborne, Robert H., 1967a, The American Upper Ordovician Standard. VIII. R-mode factor analysis of Cincinnati limestones: *Journal of Sedimentary Petrology*, v. 37, no. 2, p. 649-657.
- Osborne, Robert H., 1967b, R-mode factor analysis of Cincinnati (Ordovician) limestones (abs.): *American Association of Petroleum Geologists Bulletin*, v. 51, no. 3, pt. 2, p. 477.
- Osborne, Robert H., 1968, The American Upper Ordovician Standard. IX. Bedrock geology of eastern Hamilton County, Ohio: *American Association of Petroleum Geologists Bulletin*, v. 52, no. 11, pt. 1, p. 2137-2152.
- Osborne, Robert H., 1969, The American Upper Ordovician Standard. XI. Multivariate classification of typical Cincinnati calcarenites: *Journal of Sedimentary Petrology*, v. 39, no. 2, p. 769-776.
- Osborne, Robert H., 1970a, The American Upper Ordovician Standard. XII. Time-trend analysis of Cincinnati calcarenites, Hamilton County, Ohio: *Journal of Geology*, v. 78, no. 3, p. 363-371.
- Osborne, Robert H., 1970b, Bedrock geology of the Madeira quadrangle, Hamilton and Clermont Counties, Ohio: Ohio Division of Geological Survey Report of Investigations 77, 1 sheet, scale 1:24,000.
- Osborne, Robert H., 1971a, The American Upper Ordovician Standard. XIV. Markov analysis of typical Cincinnati sedimentation: *Journal of Sedimentary Petrology*, v. 41, no. 2, p. 444-449.
- Osborne, Robert H., 1971b, The American Upper Ordovician Standard. XVI. Utility of elastic-ratio values to distinguish Kope and Fairview Formations, Hamilton County, Ohio: *Ohio Journal of Science*, v. 71, no. 5, p. 317-319.
- Osborne, Robert H., 1972a, Depositional models and trend-surface analysis of Cincinnati (Upper Ordovician) limestones, Hamilton County, Ohio (abs.): *Geological Society of America Abstracts with Programs*, v. 4, no. 5, p. 343.
- Osborne, Robert H., 1972b, A distribution-free discrimination technique applied to typical Cincinnati (Upper Ordovician) limestones: *Ohio Journal of Science*, v. 72, no. 4, p. 224-232.
- Osborne, Robert H., 1973, The American Upper Ordovician Standard. XVII. Areal variation of limestone frequencies in the Kope and Fairview Formations, Hamilton County, Ohio: *Journal of Sedimentary Petrology*, v. 43, no. 1, p. 137-146.
- Osborne, Robert H., 1974, Bedrock geology of the Cincinnati East quadrangle, Hamilton County, Ohio: Ohio Division of Geological Survey Report of Investigations 94, 1 sheet, scale 1:24,000.
- Osborne, Robert H., Buck, S. P., Jalajas, E. W., and Wolfe, W. I., 1983, Limestone thickness distributions and paleoenvironmental implications for Kope and Fairview formations (Cincinnati), southwestern Ohio (abs.): *Geological Society of America Abstracts with Programs*, v. 15, no. 4, p. 253.
- Osborne, Robert H., Weiss, Malcolm P., and Outerbridge, William F., 1973, Geologic map of the Felicity quadrangle, Ohio-Kentucky: U.S. Geological Survey Geologic Quadrangle Map GQ-1063, scale 1:24,000.
- Osgood, Richard G., Jr., 1965, Trace fossils of the Cincinnati area: Ph.D. dissertation (unpub.), University of Cincinnati, 2 v., 536 p., 27 pls.
- Osgood, Richard G., Jr., 1970, Trace fossils of the Cincinnati area: *Palaeontographica Americana*, v. 6, no. 41, p. 277-444, pls. 57-83.
- Osgood, Richard G., Jr., 1977a, Selected Cincinnati trace fossils, in Pope, John K., and Martin, Wayne D., eds., Field guidebook to the biostratigraphy and paleoenvironments of the Cincinnati Series in southeastern Indiana: *Society of Economic Paleontologists and Mineralogists, Great Lakes Section, 7th Annual Field Conference*, Miami University, p. III1-III24 (reprinted in 1979; see Pope and Martin, 1977).
- Osgood, R. G., Jr., 1977b, "Trace fossils of the Cincinnati area" revisited, a decade later (abs.): *Journal of Paleontology*, v. 51, no. 2, suppl., part III, p. 19.
- Osgood, Richard G., Jr., and Szmuc, Eugene J., 1972, A trace fossil *Zoophycos* as an indicator of water depth: *Bulletins of American Paleontology*, v. 62, no. 271, p. 1-22, pls. 1-2.
- Outerbridge, William F., 1971a, Geologic map of the Brooksville quadrangle, Bracken County, Kentucky: U.S. Geological Survey Geologic Quadrangle Map GQ-905, scale 1:24,000.
- Outerbridge, William F., 1971b, Geologic map of the Germantown quadrangle, Mason and Bracken Counties, Kentucky: U.S. Geological Survey Geologic Quadrangle Map GQ-971, scale 1:24,000.
- Outerbridge, William F., Weiss, Malcolm P., and Osborne, Robert H., 1973, Geologic map of the Higginsport quadrangle, Ohio-Kentucky, and part of the Russellville quadrangle, Mason County, Kentucky: U.S. Geological Survey Geologic Quadrangle Map GQ-1065, scale 1:24,000.
- Pachut, Joseph F., 1987, Population genetics of four species of Ordovician bryozoans: stereology and jackknifed analysis of variance: *Journal of Paleontology*, v. 61, no. 5, p. 927-941.
- Pachut, Joseph F., 1989, Heritability and intraspecific heterochrony in Ordovician bryozoans from environments differing in diversity: *Journal of Paleontology*, v. 63, no. 2, p. 182-194.
- Pachut, Joseph F., 1992, Morphological integration and covariance during astogeny of an Ordovician trepostome bryozoan from communities of different diversities: *Journal of Paleontology*, v. 66, no. 5, p. 751-757.
- Pachut, Joseph F., and Anstey, Robert L., 1977, Diversity, stability and stratigraphic gradients in bryozoan dominated communities of Eden Shale (Cincinnati, Ohio Valley) (abs.): *Geological Society of America Abstracts with Programs*, v. 9, no. 5, p. 639-640.
- Pachut, Joseph F., and Anstey, Robert L., 1979, A developmental explanation of stability-diversity-variation hypotheses: morphogenetic regulation in Ordovician bryozoan colonies: *Paleobiology*, v. 5, no. 2, p. 168-187.
- Palmer, T. J., Hudson, J. D., and Wilson, M. A., 1988, Palaeoecological evidence for early aragonite dissolution in ancient calcite seas: *Nature*, v. 335, no. 6193, p. 809-810.
- Palmer, T. J., and Wilson, M. A., 1987, Parasitism of Ordovician bryozoans and the origin of pseudoborings (abs.): *Palaeontological Association Annual Conference*, December 17-20, 1987, Abstracts, p. 12-13.
- Palmquist, W. N., Jr., and Hall, F. R., 1960, Availability of ground water in Bracken, Harrison, Mason, Nicholas, and Robertson Counties, Kentucky: U.S. Geological Survey Hydrologic Investigations Atlas HA-16, scale 1:125,000.
- Parsley, Ronald L., 1981, Echinoderms from Middle and Upper Ordovician rocks of Kentucky: U.S. Geological Survey Professional Paper 1066-K, 9 p., 1 pl.
- Patton, John B., Perry, Thomas G., and Wayne, William J., 1953, Ordovician stratigraphy and the physiography of part of southeastern Indiana: *Indiana Geological Survey Field Conference Guidebook* 6, 29 p.
- Patzkowsky, Mark E., and Holland, Steven M., 1992, Faunal extinction, migration, and stratigraphic sequences: Middle and Late Ordovician of North America (abs.): *Geological Society of America Abstracts with Programs*, v. 24, no. 4, p. 59.
- Patzkowsky, Mark E., and Holland, Steven M., 1993, Biotic response to a Middle Ordovician paleoceanographic event in eastern North America: *Geology*, v. 21, no. 7, p. 619-622.
- Peck, J. H., 1966, Upper Ordovician formations in the Maysville area, Kentucky: U.S. Geological Survey Bulletin 1244-B, 30 p.
- Pemberton, S. George, and Frey, Robert W., 1982, Trace fossil nomenclature and the *Planolites-Palaeophycus* dilemma: *Journal of Paleontology*, v. 56, no. 4, p. 843-881, 5 pls.
- Perry, Nelson W., 1889, The Cincinnati rocks: What has been their physical history?: *American Geologist*, v. 4, no. 6, p. 326-336.
- Perry, T. G., Horowitz, A. S., and Anstey, R. L., 1973, Restudy of some of Nicholson's types of Palaeozoic bryozoans, in Larwood,

- G. P., ed., *Living and fossil Bryozoa*: London, Academic Press, p. 429-436, 1 pl.
- Peterson, W. L., 1981, Lithostratigraphy of Silurian rocks exposed on the west side of the Cincinnati Arch in Kentucky: U.S. Geological Survey Professional Paper 1151-C, 29 p., 2 pls.
- Phillips, June R. P., 1960, Restudy of types of seven Ordovician bifoliate Bryozoa: *Palaeontology*, v. 3, pt. 1, p. 1-25, pls. 1-10.
- Phillips, June R. P. Also see Ross, June Phillips.
- Podell, Mark E., and Anstey, Robert L., 1979, The interrelationship of early colony development, monticules and branches in Palaeozoic bryozoans: *Palaeontology*, v. 22, pt. 4, p. 965-982, pls. 126-130.
- Pohana, Richard E., 1992, Landslides along Columbia Parkway (stops #1, 3), in Haneberg, William C., Riestenberg, Mary M., Pohana, Richard E., and Diekmeyer, Sharon C., Cincinnati's geologic environment: a trip for secondary-school science teachers: Ohio Division of Geological Survey Guidebook 9, p. 2-6.
- Pohowsky, Robert A., 1978, The boring ctenostomate Bryozoa: taxonomy and paleobiology based on cavities in calcareous substrata: *Bulletins of American Paleontology*, v. 73, no. 301, 192 p., 24 pls.
- Pojeta, John, Jr., 1961, The pelecypod genus *Byssonychia* as it occurs in the Cincinnati at Cincinnati, Ohio: M.S. thesis (unpub.), University of Cincinnati, 157 p., 14 pls.
- Pojeta, John, Jr., 1962, The pelecypod genus *Byssonychia* as it occurs in the Cincinnati at Cincinnati, Ohio: *Palaeontographica Americana*, v. 4, no. 30, p. 166-216, pls. 22-31.
- Pojeta, John, Jr., 1963, North American Ambonychiidae (Pelecypoda): Ph.D. dissertation (unpub.), University of Cincinnati, 396 p., 29 pls.
- Pojeta, John, Jr., 1966, North American Ambonychiidae (Pelecypoda): *Palaeontographica Americana*, v. 5, no. 36, p. 129-241, pls. 19-47.
- Pojeta, John, Jr., 1971, Review of Ordovician pelecypods: U.S. Geological Survey Professional Paper 695, 46 p., 20 pls.
- Pojeta, John, Jr., 1979, The Ordovician paleontology of Kentucky and nearby states—Introduction: U.S. Geological Survey Professional Paper 1066-A, 48 p.
- Pojeta, John, Jr., 1981, Proposed stratotypes for Ordovician stages in Kentucky (abs.): *Geological Society of America Abstracts with Programs*, v. 13, no. 7, p. 530.
- Pojeta, John, Jr., 1982, Geologic field excursion to the Ordovician rocks of southwestern Ohio and central Kentucky. Part 2 of the first Richard G. Osgood Memorial Lecture: Department of Geology, College of Wooster, 18 p.
- Pojeta, John, Jr., and Palmer, T. J., 1976, The origin of rock boring in mytilacean pelecypods: *Alcheringa*, v. 1, no. 2, p. 167-179.
- Pope, John Keyler, 1966, Comparative morphology and shell histology of the Ordovician Strophomenacea (Brachiopoda): Ph.D. dissertation (unpub.), University of Cincinnati, 270 p. (abstract: *Dissertation Abstracts International*, sec. B, *Science and Engineering*, v. 27, no. 9, p. 3152B-3153B, 1967).
- Pope, John Keyler, 1976, Comparative morphology and shell histology of the Ordovician Strophomenacea (Brachiopoda): *Palaeontographica Americana*, v. 8, no. 49, p. 129-214.
- Pope, John Keyler, 1982, Some silicified strophomenacean brachiopods from the Ordovician of Kentucky, with comments on the genus *Pionomena*: U.S. Geological Survey Professional Paper 1066-L, 30 p., 8 pls.
- Pope, John K., and Martin, Wayne D., eds., 1977, Field guidebook to the biostratigraphy and paleoenvironments of the Cincinnati Series of southeastern Indiana: Society of Economic Paleontologists and Mineralogists, Great Lakes Section, 7th Annual Field Conference, Miami University, 88 p. (not consecutively paginated) (revised edition, labeled "second printing, enlarged," issued in 1979).
- Potter, A. W., and Boucot, A. J., 1992, Middle and late Ordovician brachiopod benthic assemblages of North America, in Webby, B. D., and Laurie, J. R., eds., *Global perspectives on Ordovician geology: Proceedings of the Sixth International Symposium on the Ordovician System*, University of Sydney, Australia, 15-19 July 1991, Rotterdam, The Netherlands, A. Balkema, p. 307-323.
- Potter, Paul Edwin, 1993, Structure on top of the Middle Ordovician High Bridge/Black River Groups in the tristate area of northern Kentucky, southwestern Ohio, and southeastern Indiana: Kentucky Geological Survey Map and Chart Series 5, Series XI, 1 sheet.
- Potter, Paul Edwin, 1996, Exploring the geology of the Cincinnati/northern Kentucky region: Kentucky Geological Survey Special Publication 22, Series XI, 115 p.
- Potter, Paul E., Ausich, William I., Klee, John, Krissek, Lawrence A., Mason, Charles E., Schumacher, Gregory A., Wilson, Richard T., and Wright, Earl M., 1991, Geology of the Alexandria-Ashland Highway (Kentucky Highway 546), Maysville to Garrison: Joint Field Conference of the Geological Society of Kentucky and the Ohio Geological Society, Kentucky Geological Survey, 64 p. (the version distributed at the time of the field trip lacks the photograph of the version for public distribution).
- Potter, Paul Edwin, and Spohn, Richard, 1992, The Geoscience Information Society explores the geology of the Tri-state: Field trip guide in conjunction with 1992 Annual Meeting of the Geological Society of America, University of Cincinnati (no regular pagination).
- Prosser, Charles S., 1905, Revised nomenclature of the Ohio geological formations: Ohio Division of Geological Survey Bulletin 7, 36 p.
- Pulse, Richard Reid, 1959, Conodonts from the Upper Ordovician Fairview Formation, in Cincinnati region, Ohio, Kentucky, and Indiana: M.S. thesis (unpub.), The Ohio State University, 125 p., 3 pls.
- Pulse, Richard R., and Sweet, Walter C., 1960, The American Upper Ordovician Standard. III. Conodonts from the Fairview and McMillan Formations of Ohio, Kentucky, and Indiana: *Journal of Paleontology*, v. 34, no. 2, p. 237-264, pls. 35-37.
- Purdy, Ann C., 1991, Taphonomic classification of Kope Formation limestones (Ordovician, Cincinnati area) (abs.): *Geological Society of America Abstracts with Programs*, v. 23, no. 3, p. 54.
- Pursell, Benjamin F., and Cuffey, Roger J., 1995, Bryozoan species surrounding and below the Maysville bryozoan reefs (Upper Ordovician, north-central Kentucky) (abs.): *Geological Society of America Abstracts with Programs*, v. 27, no. 3, p. 81.
- Rabbio, Salvatore Frank, 1988, Ecological and taphonomic gradients in storm disturbed bryozoan communities of the Kope Formation (Cincinnati Series, Upper Ordovician), Cincinnati Arch region: M.S. thesis (unpub.), Michigan State University, 219 p.
- Rassman, Barbara, 1981, A sedimentologic comparison of two dissimilar Ordovician edriasteroid pavements (hardgrounds): M.S. thesis (unpub.), University of Cincinnati, 224 p.
- Raymond, William H., 1960, A study of the fauna of the Whitewater Formation at Indian Creek, Butler County, Ohio: M.S. thesis (unpub.), Miami University, 111 p., 17 pls.
- Reierson, Carolyn J., Wilson, Mark A., and Palmer, Timothy J., 1988, Diagenetic origin, depositional environments and stratigraphy of encrusted limestone cobbles in the Cincinnati Series (Upper Ordovician) of northern Kentucky (abs.): *Geological Society of America Abstracts with Programs*, v. 20, no. 5, p. 385.
- Reinhart, Roy H., 1977a, Field Trip No. 2. Road log to Armco Steel Company, Middletown, Ohio (Stop 5) and NYC RR cut—I-75 near Maud, Ohio (Stop 6), and section descriptions, in Pope, John K., and Martin, Wayne D., eds., *Field guidebook to the biostratigraphy and paleoenvironments of the Cincinnati Series in southeastern Indiana: Society of Economic Paleontologists and Mineralogists, Great Lakes Section, 7th Annual Field Conference*, Miami University, p. 134-138 (reprinted in 1979; see Pope and Martin, 1977).
- Reinhart, Roy H., 1977b, Field Trip No. 3. Road log to Harrison, Ohio (Stop 7) and outcrops along I-74 Indiana (Stop 8), in Pope, John K., and Martin, Wayne D., eds., *Field guidebook to the biostratigraphy and paleoenvironments of the Cincinnati Series in southeastern Indiana: Society of Economic Paleontologists and Mineralogists, Great Lakes Section, 7th Annual*

- Field Conference, Miami University, p. I39-I42 (reprinted in 1979; see Pope and Martin, 1977).
- Reinhart, Roy H., and Bever, James E., 1969, Introductory geology, 3rd ed.: Dubuque, Iowa, Kendall Hunt, 182 p. (laboratory manual used at Miami University).
- Rexroad, Carl B., 1967, Stratigraphy and conodont paleontology of the Brassfield (Silurian) in the Cincinnati Arch area: Indiana Geological Survey Bulletin 36, 64 p., 4 pls.
- Richards, R. Peter, 1972, Autecology of Richmondian brachiopods (Late Ordovician of Indiana and Ohio): *Journal of Paleontology*, v. 46, no. 3, p. 386-405, pls. 1-5.
- Richards, R. Peter, 1974, Ecology of the Cornulitidae: *Journal of Paleontology*, v. 48, no. 3, p. 514-523, 1 pl.
- Riddell, J. L., 1837, Remarks on the geological features of Ohio, and some of the desiderata which might be supplied by a geological survey of this state: Report no. 60 to the 35th General Assembly of Ohio, Columbus, 34 p.
- Riesterberg, Mary M., and Haneberg, William C., 1992, Hillside evolution and landslides at Delhi Pike (stop #8), in Haneberg, William C., Riesterberg, Mary M., Pohana, Richard E., and Diekmeyer, Sharon C., Cincinnati's geologic environment: a trip for secondary-school science teachers: Ohio Division of Geological Survey Guidebook 9, p. 18-22.
- Rigby, J. Keith, 1966, Microstructure and classification of an Ordovician sponge, *Dystactospongia madisonensis* Foerste: *Journal of Paleontology*, v. 40, no. 5, p. 1127-1130, pl. 146.
- Rigby, J. Keith, 1970, *Brachiospongia tuberculata* James from the Ordovician of central Ontario: *Journal of Paleontology*, v. 44, no. 6, p. 1139-1142.
- Rigby, J. Keith, 1996, Phylum Porifera, sponges, in Feldmann, Rodney M., and Hackathorn, Merriane, eds., *Fossils of Ohio: Ohio Division of Geological Survey Bulletin 70*, p. 51-63.
- Riva, John, 1974, A revision of some Ordovician graptolites of eastern North America: *Palaeontology*, v. 17, pt. 1, p. 1-40, pls. 1-2.
- Riva, John, 1987, The graptolite *Amplexograptus praetypicalis* n. sp. and the origin of the *typicalis* group: *Canadian Journal of Earth Sciences*, v. 24, no. 5, p. 924-933.
- Rockwell, Lynn A., and Cuffey, Roger J., 1996, Great bryozoan diversity and sampling in the uppermost Ordovician (Whitewater Formation) at Caesar Creek (southwestern Ohio) (abs.): *Geological Society of America Abstracts with Programs*, v. 28, no. 6, p. 62.
- Roeminger, Carl, 1866, Observations on *Chaetetes* and some related genera, in regard to their systematic position, with an appended description of some new species: *Proceedings of the Academy of Natural Sciences of Philadelphia*, 1866, p. 113-123.
- Roeminger, C., 1890, Studies on *Monticulipora*: *American Geologist*, v. 6, no. 2, p. 102-121.
- Rosenberg, Gary D., 1982, Growth rhythms in the brachiopod *Rafinesquina alternata* from the Late Ordovician of southeastern Indiana: *Paleobiology*, v. 8, no. 4, p. 389-401.
- Ross, June Phillips, 1960, Reevaluation of the type species of *Arthropora* Ulrich: *Journal of Paleontology*, v. 34, no. 5, p. 859-861, pl. 108.
- Ross, June Phillips, 1967, Fossil problematica from Upper Ordovician, Ohio: *Journal of Paleontology*, v. 41, no. 1, p. 37-42, pls. 4-8.
- Ross, June R. P., 1970, Discussion of "On the validity of the name *Hallopora* for a genus of trepostome bryozoan" by R. J. Singh: *Journal of Paleontology*, v. 44, no. 4, p. 785.
- Ross, J. R. P., 1985, Biogeography of Ordovician ectoproct (bryozoan) faunas, in Nielsen, Claus, and Larwood, Gilbert P., eds., *Bryozoa, Ordovician to Recent: Papers presented at 6th International Conference on Bryozoa*, Vienna, 1983, Fredensborg, Denmark, Olsen and Olsen, p. 265-271.
- Ross, J. R. P. Also see Phillips, June R. P.
- Ross, Reuben James, Jr., 1967, Calymenid and other Ordovician trilobites from Kentucky and Ohio: U.S. Geological Survey Professional Paper 583-B, 19 p., 5 pls.
- Ross, Reuben James, Jr., 1979, Additional trilobites from the Ordovician of Kentucky: U.S. Geological Survey Professional Paper 1066-D, 13 p., 6 pls.
- Ross, Reuben J., Jr., and others, 1982, The Ordovician System in the United States: International Union of Geological Sciences Publication 12, 73 p., 3 correlation charts.
- Rudkin, David M., and Tripp, Ronald P., 1989, The type species of the Ordovician trilobite genus *Isotelus*: *I. gigas* DeKay, 1824: *Royal Ontario Museum Life Sciences Contributions* 152, 19 p.
- Ruedemann, Rudolf, 1901, Hudson River beds near Albany and their taxonomic equivalents: *Bulletin of the New York State Museum*, v. 8, no. 42, p. 485-596, 2 pls.
- Ruedemann, Rudolf, 1925a, The Utica and Lorraine Formations of New York. Part I. Stratigraphy: *New York State Museum Bulletin*, no. 258, p. 1-175.
- Ruedemann, Rudolf, 1925b, The Utica and Lorraine Formations of New York. Part 2. Systematic paleontology. No. 1. Plants, sponges, corals, graptolites, crinoids, worms, bryozoans, brachiopods: *New York State Museum Bulletin*, no. 262, p. 1-171.
- Ruedemann, Rudolf, 1926, The Utica and Lorraine Formations of New York. Part 2. Systematic paleontology. No. 2. Mollusks, crustaceans and eurypterids: *New York State Museum Bulletin*, no. 272, p. 1-227.
- Ruedemann, Rudolf, 1947, Graptolites of North America: *Geological Society of America Memoir* 19, 652 p., 92 pls.
- Rytel, Jenny Lyn, 1982, Conodont and brachiopod biostratigraphy of Upper Ordovician rocks near Ripley, Ohio: M.S. thesis (unpub.), The Ohio State University, 108 p.
- Rytel, Jenny, and Bergström, Stig M., 1985, Lower and Middle Paleozoic geology of southern Ohio: Sixth Gondwana Symposium, The Ohio State University, 1985, Field Excursion 10, Guidebook, 32 p.
- Sandy, Michael R., 1996, Oldest record of peduncular attachment of brachiopods to crinoid stems, Upper Ordovician, Ohio, U.S.A. (Brachiopoda; Atrypida; Echinodermata; Crinoidea): *Journal of Paleontology* v. 70, no. 3, p. 532-534.
- Sardeson, Frederick W., 1901, Problems of the Monticuliporidae, I: *Journal of Geology*, v. 9, no. 1, p. 1-27, pl. A.
- Sardeson, Frederick W., 1936, Early bryozoans: *Monotrypa* to *Eridotrypa*: *Pan-American Geologist*, v. 66, no. 3, p. 179-190, pl. 15.
- Schallreuter, Roger, 1981, Ordovizische Problematika III. *Ancientia* Ross, 1967 aus Europa: *Paläontologische Zeitschrift*, v. 55, no. 3/4, p. 209-218, 3 pls.
- Scheufler, Lowell W., 1956, The geology of the Darrrtown-McGonigle area, Butler County, Ohio: M.S. thesis (unpub.), Miami University, 58 p.
- Schmidt, Ronald G., McFarlan, Arthur C., Nosow, Edmund, Bowman, Richard S., and Alberts, Robert, 1961, Examination of the Ordovician through Devonian stratigraphy and the Serpent Mound chaotic structure area, field trip 8: Guidebook for Field Trips, Cincinnati meeting, 1961, Geological Society of America, p. 259-293.
- Schopf, J. M., Ehlers, E. G., Stiles, D. V., and Birle, J. D., 1965, Fossil iron bacteria preserved in pyrite: *American Philosophical Society Proceedings*, v. 109, no. 5, p. 288-308.
- Schopf, Thomas J. M., 1969, Paleocology of ectoprocts (bryozoans): *Journal of Paleontology*, v. 43, no. 2, p. 234-244.
- Schumacher, Gregory A., 1984, Sedimentology, biostratigraphy, and paleocology of Cincinnati (Upper Ordovician) *Iocrinus subcrassus* crinoid (Echinodermata) assemblages: M.S. thesis (unpub.), University of Cincinnati, 210 p., 8 pls.
- Schumacher, Gregory A., 1985, Fossil lag deposits: a key to Upper Ordovician crinoid biostratigraphy (abs.): *Geological Society of America Abstracts with Programs*, v. 17, no. 5, p. 325.
- Schumacher, Gregory A., 1986a, Bedrock geology of Clermont County, Ohio (abs.): *Ohio Journal of Science*, v. 86, no. 2, p. 9.
- Schumacher, Gregory A., 1986b, Ordovician echinoderm fossils: Ohio Division of Geological Survey, Ohio Geology Newsletter, Summer, p. 5-7.
- Schumacher, Gregory A., 1986c, Storm processes and crinoid preservation (abs.): *Proceedings of the 4th North American Paleontological Convention*, p. A41.
- Schumacher, Gregory A., 1988, Middle and Upper Ordovician stratigraphy of south-western Ohio (abs.): *Geological Society*

- of America Abstracts with Programs, v. 20, no. 5, p. 387.
- Schumacher, Gregory A., 1990, Bedrock geology of the Goshen, Ohio, quadrangle: Ohio Division of Geological Survey Open-File Map BG-B5B2.
- Schumacher, Gregory A., 1992, Lithostratigraphy, cyclic sedimentation, and event stratigraphy of the Maysville, Kentucky area, Stop 8, *in* Ettensohn, Frank R., ed., Changing interpretations of Kentucky geology: layer cake, facies, flexure, and eustasy (field trip for 1992 Annual Meeting of the Geological Society of America): Ohio Division of Geological Survey Miscellaneous Report 5, p. 165-172.
- Schumacher, Gregory A., and Ausich, William I., 1983, New Upper Ordovician echinoderm site: Bull Fork Formation, Caesar Creek Reservoir (Warren County, Ohio): Ohio Journal of Science, v. 83, no. 1, p. 60-64.
- Schumacher, Gregory A., and Ausich, William I., 1985, Catastrophic sedimentation: impact on a Late Ordovician crinoid assemblage (abs.): Geological Society of America Abstracts with Programs, v. 17, no. 5, p. 325.
- Schumacher, Gregory A., Bergström, Stig M., and Mitchell, Charles E., 1991, The Sebree Trough project. 4: preliminary tracing of surficial Upper Ordovician lithostratigraphic units into the subsurface of western and northern Ohio (abs.): Geological Society of America Abstracts with Programs, v. 23, no. 3, p. 59.
- Schumacher, Gregory A., and Carlton, Richard W., 1991, Impure K-bentonite beds from the Lexington Limestone and the Point Pleasant Formation (Middle Ordovician) of northern Kentucky and southwestern Ohio: Southeastern Geology, v. 32, no. 2, p. 83-105.
- Schumacher, Gregory A., and Caudill, Michael R., 1992, Combined flow storm-generated crinoid taphonomy: an example from the Upper Ordovician, Cincinnati Series in northeastern Kentucky (abs.): Geological Society of America Abstracts with Programs, v. 24, no. 4, p. 63.
- Schumacher, Gregory A., and Meyer, David L., 1986, Tempestites and variable crinoid preservation: examples from the Upper Ordovician of Ohio and Indiana (abs.): Geological Society of America Abstracts with Programs, v. 18, no. 4, p. 323.
- Schumacher, Gregory A., Potter, Paul E., and Noger, Martin C., 1993, The new Alexandria to Ashland, Kentucky, highway: an extraordinary geologic "classroom" (abs.): Ohio Journal of Science, v. 93, no. 2, p. 39.
- Schumacher, Gregory A., and Shrake, Douglas L., 1989, *The Isotelus gigas* (trilobite) bed: revisited (abs.): Geological Society of America Abstracts with Programs, v. 21, no. 2, p. 65.
- Schumacher, Gregory A., Shrake, Douglas L., Swinford, E. Mac, Brockman, C. Scott, and Wickstrom, Lawrence H., 1987, Stratigraphy and depositional environments of the Cincinnati Group of southwestern Ohio: Field trip guidebook for Eastern Section meeting, American Association of Petroleum Geologists, Ohio Geological Society, 73 p.
- Schumacher, Gregory A., Swinford, E. Mac, and Bergström, Stig M., 1990, The Sebree Trough Project. 1. Lithostratigraphy and geologic setting of the Oxford, Ohio, core (abs.): Geological Society of America Abstracts with Programs, v. 22, no. 3, p. 43.
- Schumacher, Gregory A., Swinford, E. Mac, and Shrake, Douglas L., 1986, Southwestern Ohio stratigraphy, reevaluated (abs.): Geological Society of America Abstracts with Programs, v. 18, no. 4, p. 323.
- Schumacher, Gregory A., Swinford, E. Mac, and Shrake, Douglas L., 1987, Straight Creek Member of the Grant Lake Limestone: a new map unit in southwestern Ohio (abs.): Geological Society of America Abstracts with Programs, v. 19, no. 4, p. 243.
- Schumacher, Gregory A., Swinford, E. Mac, and Shrake, Douglas L., 1989, A new lithostratigraphic unit in southwestern Ohio: the Straight Creek Member of the Grant Lake Limestone (abs.): Ohio Journal of Science, v. 89, no. 2, p. 11.
- Schumacher, Gregory A., Swinford, E. Mac, and Shrake, Douglas L., 1991, Lithostratigraphy of the Grant Lake Limestone and Grant Lake Formation (Upper Ordovician) in southwestern Ohio: Ohio Journal of Science, v. 91, no. 1, p. 56-68.
- Schwimmer, Barbara A., and Sandy, Michael R., 1996, Phylum Brachiopoda, *in* Feldmann, Rodney M., and Hackathorn, Merrienne, eds., Fossils of Ohio: Ohio Division of Geological Survey, Bulletin 70, p. 210-241.
- Scotford, David M., 1964, The Cincinnati Arch: mineralogical-statistical evidence of a post-Ordovician origin: American Association of Petroleum Geologists Bulletin, v. 48, no. 4, p. 427-436.
- Scotford, David M., 1965, Petrology of the Cincinnati Series shales and environmental implications: Geological Society of America Bulletin, v. 76, no. 2, p. 193-222.
- Scotford, David M., and Smoot, Thomas W., 1962, Petrography of Cincinnati Series mudstones of southwestern Ohio and adjacent areas (abs.): Geological Society of America Special Paper 68, p. 265-266.
- Seddon, George, and Sweet, Walter C., 1971, An ecologic model for conodonts: Journal of Paleontology, v. 45, no. 5, p. 869-880.
- Shade, John W., 1963, Mineral equilibria investigations in the Cincinnati Series shales of Ohio, Indiana, and Kentucky: M.S. thesis (unpub.), Miami University, 61 p.
- Shafor, Kenneth W., 1959, The geology of the Trenton quadrangle Butler County, Ohio: M.S. thesis (unpub.), Miami University, 101 p., 3 pls.
- Shaler, N. S., 1876, On the fossil brachiopods of the Ohio Valley: Memoirs of the Geological Survey of Kentucky, v. 1, pt. 3, 44 p., 7 pls.
- Sharp, Everett Ray, 1957, Petrography and petrology of the Southgate Member, Eden Group, southwestern Ohio: M.S. thesis (unpub.), The Ohio State University, 151 p., 4 pls.
- Shaver, Robert H., Ault, Curtis H., Burger, Ann M., Carr, Donald D., Droste, John B., Eggert, Donald L., Gray, Henry H., Harper, Denver, Hasenmueller, Nancy R., Hasenmueller, Walter A., Horowitz, Alan S., Hutchison, Harold C., Keith, Brian D., Keller, Stanley J., Patton, John B., Rexroad, Carl B., and Wier, Charles E., 1986, Compendium of Paleozoic rock-unit stratigraphy in Indiana—a revision: Indiana Geological Survey Bulletin 59, 203 p., 2 pls.
- Shaver, Robert H., Burger, Ann M., Gates, Gary R., Gray, Henry H., Hutchison, Harold C., Keller, Stanley J., Patton, John B., Rexroad, Carl B., Smith, Ned M., Wayne, William J., and Wier, Charles E., 1970, Compendium of rock-unit stratigraphy in Indiana: Indiana Geological Survey Bulletin 43, 229 p., 1 pl.
- Shaw, Frederick Carleton, 1960, Richmondian (Upper Ordovician) Rafinesquinae, Cincinnati, Ohio: M.S. thesis (unpub.), University of Cincinnati, 156 p., 7 pls.
- Shaw, Frederick C., 1962, Cincinnati rafinesquinoid brachiopods (abs.): Geological Society of America Special Paper 68, p. 268-269.
- Shaw, Frederick C., and Lespérance, Pierre J., 1994, North American biogeography and taxonomy of *Cryptolithus* (Trilobita, Ordovician): Journal of Paleontology, v. 68, no. 4, p. 808-823.
- Sheehan, Peter M., 1988, Constraints of brachiopod zoogeography on correlation in the Late Ordovician and Early Silurian of North America: New York State Museum Bulletin 462, p. 80-83.
- Shideler, W. H., 1914, The upper Richmond beds of the Cincinnati Group: Ohio Naturalist, v. 14, no. 3, p. 229-235.
- Shideler, W. H., 1916, The Ordovician-Silurian boundary: Ohio Journal of Science, v. 16, no. 8, p. 329-335.
- Shideler, W. H., 1918, A primitive type of *Agelacrinites* from the Richmond: Ohio Journal of Science, v. 19, no. 1, p. 58.
- Shideler, W. H., 1923, The basal Richmond of the Cincinnati province (abs.): Geological Society of America Bulletin, v. 34, p. 132.
- Shideler, W. H., 1928, A re-study of the Hamburg, Indiana, section (abs.): Ohio Academy of Science Proceedings, v. 8, pt. 4, p. 173.
- Shideler, William Henry, 1934, New Cincinnati sponges (abs.): Geological Society of America Proceedings for 1933, p. 341-342.
- Shideler, William Henry, 1937, New species of Cincinnati fossils (abs.): Geological Society of America Proceedings for 1936, p. 367.
- Shideler, W. H., 1939, Paleogeography of the Cincinnati Province: Compass, v. 19, no. 3, p. 201-211.
- Shideler, William Henry. See also Marak, Joe H., 1992.
- Shimer, Hervey W., and Shrock, Robert R., 1944, Index fossils of North America; a new work based on the complete revision and reillustration of Grabau and Shimer's "North American index fossils": New York, J. Wiley and Sons, Inc.; London, Chapman

- & Hall, Ltd., 837 p., 303 pls.
- Shrake, Douglas L., 1987, Paleocology of a Cincinnati (Upper Ordovician) bryozoan-trilobite association (abs.): *Ohio Journal of Science*, v. 87, no. 2, p. 10.
- Shrake, Douglas Lee, 1989, Paleocology of a Cincinnati (Upper Ordovician) bryozoan-trilobite association: M.S. thesis (unpub.), Wright State University, 90 p.
- Shrake, Douglas L., 1990, Common trilobites of Ohio: *Ohio Division of Geological Survey, Ohio Geology Newsletter*, Summer, p. 1, 3-5.
- Shrake, Douglas L., 1992, Excursion to Caesar Creek State Park in Warren County, Ohio: a classic Upper Ordovician fossil-collecting locality: *Ohio Division of Geological Survey Guidebook* 12, 18 p., 9 pls.
- Shrake, Douglas L., 1994, Ohio trilobites: *Ohio Division of Geological Survey GeoFacts* 5, one sheet.
- Shrake, Douglas L., 1995, *Isotelus*: Ohio's state fossil: *Ohio Division of Geological Survey GeoFacts* 6, one sheet.
- Shrake, Douglas L., Carlton, Richard W., Wickstrom, Lawrence H., Potter, Paul Edwin, Richard, Benjamin H., Wolfe, Paul J., and Sittler, Gary M., 1991, Pre-Mount Simon basin under Cincinnati Arch: *Geology*, v. 19, no. 2, p. 139-142.
- Shrake, Douglas L., and Schumacher, Gregory A., 1989, Excavation of an *Isotelus* shale layer at Caesar Creek State Park (abs.): *Ohio Journal of Science*, v. 89, no. 2, p. 9.
- Shrake, Douglas L., Schumacher, Gregory A., and Swinford, E. Mac, 1988, The stratigraphy, sedimentology and paleontology of the Upper Ordovician Cincinnati Group of southwest Ohio: Field trip guidebook for Fifth Midyear Meeting of Society of Economic Paleontologists and Mineralogists, Ohio Division of Geological Survey, 81 p. (title as given on title page is: Field guidebook to the stratigraphy, sedimentology, and paleontology of the Upper Ordovician rocks of southwestern Ohio).
- Shrake, Douglas L., Wolfe, Paul J., Richard, Benjamin H., Swinford, E. Mac, Wickstrom, Lawrence H., Potter, Paul E., and Sittler, Gary W., 1990, Lithologic and geophysical description of a continuously cored hole in Warren County, Ohio, including description of the Middle Run Formation (Precambrian?) and a seismic profile across the core site: *Ohio Division of Geological Survey Information Circular* 56, 11 p., 2 pls.
- Shuff, Sheldon G., 1974a, A mineralogical study of the Cincinnati Series limestones (Upper Ordovician), southwestern Ohio and southeastern Indiana: M.S. thesis (unpub.), Miami University, 67 p.
- Shuff, Sheldon G., 1974b, Use of the Ca/Mg ratios of carbonate sediments as a stratigraphic tool in correlating portions of Cincinnati Series (Upper Ordovician) limestones: *Compass*, v. 52, no. 1, p. 1-8.
- Sigma Gamma Epsilon, 1968, 1968 National Convention, Cincinnati, Ohio—handbook and guidebook: Cincinnati, Sigma Gamma Epsilon, Alpha Delta Chapter, 79 p.
- Simmons, George C., and Oliver, William A., Jr., 1967, Otter Creek coral bed and its fauna, east-central Kentucky: *U.S. Geological Survey Bulletin* 1244-F, 13 p.
- Simpson, C. B., 1897, A handbook of the genera of the North American Paleozoic Bryozoa: with an introduction upon the structure of living species: *State Geologist of New York, 14th Annual Report for 1894*, p. 407-669, pls. A-E.
- Singh, Raman J., 1966, Conodonts from the Bellevue Limestone Member of the McMillan Formation (Upper Ordovician) of Cincinnati, Ohio: M.S. thesis (unpub.), University of Cincinnati, 98 p., 3 pls.
- Singh, Raman J., 1970, On the validity of the name *Hallopora* for a genus of trepostome bryozoan: *Journal of Paleontology*, v. 44, no. 4, p. 784-785.
- Singh, Raman J., 1971, Trepostomatous bryozoan fauna and stratigraphy of the Bellevue Limestone, Upper Ordovician, in the tri-state area of Ohio, Indiana, and Kentucky: Ph.D. dissertation (unpub.), University of Cincinnati, 188 p., 33 pls.
- Singh, Raman J., 1979, Trepostomatous bryozoan fauna from the Bellevue Limestone, Upper Ordovician, in the tri-state area of Ohio, Indiana, and Kentucky: *Bulletins of American Paleontology*, v. 76, no. 307, p. 157-288, pls. 16-48.
- Slaughter, Arthur E., compiler, 1953, The Ordovician stratigraphy of the Cincinnati, Ohio and Richmond, Indiana areas: Guidebook for Annual Geological Excursion of the Michigan Geological Society 14 p., 8 pls. (not consistently paginated).
- Sloan, Robert E., 1986, Tectonics, biostratigraphy and lithostratigraphy of the Mohawkian and Cincinnati of the Upper Mississippi Valley (abs.): *Geological Society of America Abstracts with Programs*, v. 18, no. 4, p. 324.
- Sloan, Robert E., 1987, Tectonics, biostratigraphy and lithostratigraphy of the Middle and Late Ordovician of the Upper Mississippi Valley: *Minnesota Geological Survey Report of Investigations* 35, p. 7-20.
- Sloan, Robert E., 1991, A chronology of North American trilobite genera, in Barnes, Christopher R., and Williams, S. Henry, eds., *Advances in Ordovician geology: Geological Survey of Canada Paper* 90-9, p. 165-177.
- Smith, Alison J., and Tillman, John R., 1996, Phylum Arthropoda, Class Crustacea, Subclass Ostracoda, in Feldmann, Rodney M., and Hackathorn, Merrienne, eds., *Fossils of Ohio: Ohio Division of Geological Survey, Bulletin* 70, p. 114-123.
- Smith, Andrew B., and Wilson, Mark A., 1995, A new cyclostoid (Echinodermata) from the Late Ordovician of Kentucky, U.S.A.: *Journal of Paleontology*, v. 69, no. 6, p. 1186-1187.
- Smith, Leigh, 1988, Cyclic sealevel variation as the cause of siliciclastic influx and burrow dolomitization, Upper Ordovician, Manitoba (abs.): *Society of Economic Mineralogists and Paleontologists Annual Midyear Meeting Abstracts*, v. 5, p. 50.
- Spjeldnaes, Nils, 1963, A redescription of the type species of the bryozoan genus *Crepipora*: *Journal of Paleontology*, v. 37, no. 1, p. 64-68, pls. 9-10.
- Spoley, Robert J., 1967, The bedrock geology of the Oxford quadrangle, Butler-Preble Counties, Ohio: M.A. thesis (unpub.), Miami University, 140 p.
- Staursky, Geoffrey N., 1981, The petrology of selected limestone beds of the Bull Fork Formation (Cincinnati Series), Caesar Creek Lake spillway, Warren County, Ohio: M.S. thesis (unpub.), Miami University, 154 p.
- Stith, David A., 1979, Chemical composition, stratigraphy, and depositional environments of the Black River Group (Middle Ordovician), southwestern Ohio: *Ohio Division of Geological Survey Report of Investigations* 113, 36 p., 3 pls.
- Stith, David A., 1986, Supplemental core investigations for high-calcium limestones in western Ohio and discussion of natural gas and stratigraphic relationships in the Middle to Upper Ordovician rocks of southwestern Ohio: *Ohio Division of Geological Survey Report of Investigations* 132, 17 p., 1 pl.
- Stout, Wilber E., 1941, Dolomites and limestones of western Ohio: *Ohio Division of Geological Survey Bulletin* 42, 468 p.
- Stout, Wilber, 1943, Generalized section of rocks in Ohio: *Ohio Division of Geological Survey Information Circular* 4 (reprinted from *Ohio Division of Geological Survey Bulletin* 44, p. 78-117).
- Stout, Wilber E., and Schaaf, Downs, 1931, Minford silts of southern Ohio: *Geological Society of America Bulletin*, v. 42, no. 3, p. 663-672.
- Stout, Wilber, Ver Steeg, Karl, and Lamb, G. F., 1943, *Geology of water in Ohio (a basic report): Ohio Division of Geological Survey Bulletin* 44, 694 p.
- Straw, William Thomas, 1960, Stratigraphy of the Marble Hill facies and contiguous strata of the Waynesville Formation (Richmondian) in southeastern Indiana and north-central Kentucky: M.S. thesis (unpub.), Indiana University, 50 p.
- Strete, R. F., 1939, The Saluda Division of the Whitewater Formation of Ohio: *Compass*, v. 19, no. 3, p. 212-217.
- Swadley, W C, 1972a, Geologic map of parts of the Lawrenceburg, Aurora, and Hooven quadrangles, Boone County, Kentucky: U.S. Geological Survey Geologic Quadrangle Map GQ-989, scale 1:24,000.
- Swadley, W C, 1972b, Geologic map of the Elliston quadrangle, Grant County, Kentucky: U.S. Geological Survey Geologic Quadrangle Map GQ-994, scale 1:24,000.
- Swadley, W C, 1973a, Geologic map of the Sanders quadrangle,

- north-central Kentucky: U.S. Geological Survey Geologic Quadrangle Map GQ-1095, scale 1:24,000.
- Swadley, W C, 1973b, Geologic map of parts of the Vevay South and Vevay North quadrangles, north-central Kentucky: U.S. Geological Survey Geologic Quadrangle Map GQ-1123, scale 1:24,000.
- Swadley, W C, 1974, Geologic map of the Glencoe quadrangle, north-central Kentucky: U.S. Geological Survey Geologic Quadrangle Map GQ-1154, scale 1:24,000.
- Swadley, W C, 1976, Geologic map of part of the Carrollton quadrangle, Carroll and Trimble Counties, Kentucky: U.S. Geological Survey Geologic Quadrangle Map GQ-1281, scale 1:24,000.
- Swadley, W C, 1979, The Marble Hill Bed: an offshore bar-tidal channel complex in the Upper Ordovician Drakes Formation of Kentucky: U.S. Geological Survey Professional Paper 1126-D, 8 p.
- Swadley, W C, and Gibbons, A. B., 1976, Geologic map of the Campbellsburg quadrangle, north-central Kentucky: U.S. Geological Survey Geologic Quadrangle Map GQ-1354, scale 1:24,000.
- Swadley, W C, Luft, S. J., and Gibbons, A. B., 1975, The Point Pleasant Tongue of the Clays Ferry Formation, northern Kentucky: U.S. Geological Survey Bulletin 1405-A, p. 30-31.
- Sweet, Walter C., 1959, Distribution and stratigraphic significance of conodonts in the type section of the Cincinnati Series (abs.): Geological Society of America Bulletin, v. 70, no. 12, pt. 2, p. 1684.
- Sweet, Walter C., 1979a, Conodonts and conodont biostratigraphy of post-Tyrone Ordovician rocks of the Cincinnati region: U.S. Geological Survey Professional Paper 1066-G, 26 p.
- Sweet, Walter C., 1979b, Late Ordovician conodonts and biostratigraphy of the western Midcontinent Province: Brigham Young University, Geology Studies, v. 26, no. 3, p. 45-86, figs. (= pls.) 5-10.
- Sweet, Walter C., 1983, Graphic correlation of Middle and Late Ordovician rocks, North America: Appalachian Basin Industrial Associates Program for Meeting, v. 5, p. 4-22.
- Sweet, Walter C., 1984a, A conodont-based standard reference section in Ordovician rocks of the Cincinnati region (abs.): Geological Society of America Abstracts with Programs, v. 16, no. 3, p. 201.
- Sweet, Walter C., 1984b, Conodont provinces and biofacies of the Late Ordovician: Geological Society of America Special Paper 196, p. 69-87.
- Sweet, Walter C., 1984c, Conodonts, conodont biostratigraphy, and correlation of the Moffett Road section (Middle and Upper Ordovician), Kenton County, Kentucky: U.S. Geological Survey Open-File Report 84-270, 20 p.
- Sweet, Walter C., 1984d, Graphic correlation of upper Middle and Upper Ordovician rocks, North American Midcontinent Province, U.S.A., in Bruton, D. L., ed., Aspects of the Ordovician System: Palaeontological Contributions from the University of Oslo 295, p. 23-25.
- Sweet, W. C., 1986, Graphic rejuvenation of Ordovician chronostratigraphic scale (abs.): American Association of Petroleum Geologists Bulletin, v. 70, no. 5, p. 654.
- Sweet, Walter C., 1988, Mohawkian and Cincinnati chronostratigraphy: New York State Museum Bulletin 462, p. 84-90.
- Sweet, Walter C., 1996, Phylum Conodonta, in Feldmann, Rodney M., and Hackathorn, Merrienne, eds., Fossils of Ohio: Ohio Division of Geological Survey Bulletin 70, p. 262-269.
- Sweet, Walter C., and Bergström, Stig M., 1970, Stratigraphic significance of conodonts from the Lexington Limestone and Kope Formation in the Cincinnati region (abs.): Geological Society of America Abstracts with Programs, v. 2, no. 3, p. 242-243.
- Sweet, Walter C., and Bergström Stig M., 1971a, A revised time-stratigraphic classification of the North American upper Middle and Upper Ordovician (abs.): Geological Society of America Abstracts with Programs, v. 2, no. 7, p. 698.
- Sweet, W. C., and Bergström, S. M., 1971b, The American Upper Ordovician Standard: XIII. A revised time-stratigraphic classification of North American Upper Middle and Upper Ordovician rocks: Geological Society of America Bulletin, v. 82, no. 3, p. 613-628.
- Sweet, W. C., and Bergström, S. M., 1976, Conodont biostratigraphy of the Middle and Upper Ordovician of the U.S. Midcontinent, in Bassett, M. G., ed., The Ordovician System: Proceedings of a Palaeontological Association Symposium, Birmingham, September, 1974: Cardiff, University of Wales Press and National Museum of Wales, p. 121-151.
- Sweet, Walter C., and Bergström, Stig M., 1981, Conodonts, conodont biostratigraphy and correlation of Middle and Late Ordovician rocks of the Cincinnati region (Kentucky, Ohio, Indiana) (abs.): Geological Society of America Abstracts with Programs, v. 13, no. 7, p. 563.
- Sweet, Walter C., and Bergström, Stig M., 1984, Conodont provinces and biofacies of the Late Ordovician: Geological Society of America Special Paper 196, p. 69-87.
- Sweet, Walter C., and Bergström, Stig M., 1986, Conodonts and biostratigraphic correlation: Annual Review of Earth and Planetary Science, v. 14, p. 85-112.
- Sweet, Walter C., Bergström, Stig M., and Rust, Claude C., 1965, Relative abundance analysis: an aid to Ordovician conodont biostratigraphy in the Cincinnati region of Ohio (abs.): Geological Society of America Special Paper 82, p. 203.
- Sweet, Walter C., Ethington, Raymond L., and Barnes, Christopher R., 1971, North American Middle and Upper Ordovician conodont faunas, in Sweet, Walter C., and Bergström, Stig M., eds., Symposium on conodont biostratigraphy: Geological Society of America Memoir 127, p. 163-193, 2 pls.
- Sweet, W. C., Harper, Howard, Jr., and Zlatkin, Dennis, 1974, The American Upper Ordovician Standard. XIX. A Middle and Upper Ordovician reference standard for the eastern Cincinnati region: Ohio Journal of Science, v. 74, no. 1, p. 47-54.
- Sweet, Walter C., and Miller, A. K., 1958, Ordovician cephalopods from Cornwallis and Little Cornwallis Islands, District of Franklin, Northwest Territories: Geological Survey of Canada Bulletin, v. 38, 86 p., 7 pls.
- Sweet, Walter C., Turco, Caroline A., Warner, Earl, Jr., and Wilkie, Lorna C., 1959, The American Upper Ordovician Standard. I. Eden conodonts from the Cincinnati region, Ohio and Kentucky: Journal of Paleontology, v. 33, no. 6, p. 1029-1068, pls. 130-133.
- Swinford, E. Mac, 1983, Geology of the Peebles quadrangle, Adams County, Ohio: M.S. thesis (unpub.), Eastern Kentucky University, 104 p.
- Swinford, E. Mac, 1985, Geology of the Peebles quadrangle, Adams County, Ohio: Ohio Journal of Science, v. 85, no. 5, p. 218-230.
- Swinford, E. Mac, 1986, Bedrock geologic map of Hamilton County, Ohio (abs.): Ohio Journal of Science, v. 86, no. 2, p. 8-9.
- Swinford, E. Mac, 1991, Lithostratigraphy of five bedrock cores of some Upper, Middle, and Lower Ordovician rocks in Cincinnati, Ohio (abs.): Ohio Journal of Science, v. 91, no. 2, p. 34.
- Swinford, E. Mac, and Schumacher, Gregory A., 1985, New bedrock geologic maps of Hamilton and Clermont Counties, Ohio (abs.): Ohio Journal of Science, v. 85, no. 2, p. 24-25.
- Tang, Su, and Cuffey, Roger J., 1989, Constellariid bryozoan species in the type Cincinnati (Upper Ordovician: Ohio, Indiana, Kentucky) (abs.): Geological Society of America Abstracts with Programs, v. 21, no. 4, p. 49.
- Tankersley, Ken, and Selms, Dennis C., 1978, A handbook to the Upper Ordovician of Cincinnati: Privately published by Selms, 1220 Fairbanks Ave., Cincinnati, Ohio, 31 p.
- Taylor, J. S., 1850, Notice of trilobites in the cabinet of Dr. Julius S. Taylor: American Journal of Science, ser. 2, v. 10, no. 28, p. 113-114.
- Taylor, Paul D., and Wilson, Mark A., 1994, *Corynotrypa* from the Ordovician of North America: colony growth in a primitive stenolaemate bryozoan: Journal of Paleontology, v. 68, no. 2, p. 241-257.
- Taylor, Paul D., and Wilson, Mark A., 1996, *Cuffeyella*, a new bryozoan genus from the Late Ordovician of North America, and its bearing on the origin of the post-Paleozoic cyclostomates, in Gordon, Dennis P., Smith, Abigail M., and Grant-Mackie, Jack A., eds., Bryozoans in space and time: Wellington, New

- Zealand, Proceedings of the 10th International Bryozoology Conference (1995), National Institute of Water & Atmospheric Research, Ltd., p. 351-360.
- Thompson, D. R., 1963, Astogenetic study of some trepostomatous Bryozoa in the Waynesville Formation of the Richmond Group: M.S. thesis (unpub.), Miami University, 138 p.
- Thompson, Esther H., 1969, Morphology and taxonomy of *Cyclonema* Hall (Gastropoda), Upper Ordovician, Cincinnati Province: M.S. thesis (unpub.), Miami University, 148 p.
- Thompson, Esther H., 1970, Morphology and taxonomy of *Cyclonema* Hall (Gastropoda), Upper Ordovician, Cincinnati Province: *Bulletins of American Paleontology*, v. 58, no. 261, p. 215-284, pls. 31-35.
- Tobin, Rick C., 1980, Sedimentology of the Fairview Formation, Miami-town Shale, and Bellevue Limestone (Upper Ordovician, Cincinnati Series) of southwestern Ohio and northern Kentucky: M.S. thesis (unpub.), University of Cincinnati, 144 p. 8 pls.
- Tobin, Rick C., 1982, A model for cyclic deposition in the Cincinnati Series of southwestern Ohio, northern Kentucky, and southeastern Indiana: Ph.D. dissertation (unpub.), University of Cincinnati, 483 p.
- Tobin, Rick C., 1986, An assessment of the lithostratigraphic and interpretive value of the traditional "biostratigraphy" of the type Upper Ordovician of North America: *American Journal of Science*, v. 286, no. 9, p. 673-701.
- Tobin, Rick C., and Pryor, Wayne A., 1985, Cincinnati Series—model for cyclic and episodic deposition of carbonates and shales on a storm-dominated ramp (abs.): *American Association of Petroleum Geologists Bulletin*, v. 69, no. 2, p. 311-312.
- Tofel, Jane Evans, and Bretsky, Peter W., 1987, Middle Ordovician *Lophospira* (Archaeogastropoda) from the upper Mississippi Valley: *Journal of Paleontology*, v. 61, no. 4, p. 700-723.
- Totten, Stanley M., and Hay, Helen B., 1987, Madison, Indiana: geomorphology, and Paleozoic and Quaternary geology, in Biggs, Donald L., ed., North-Central Section of the Geological Society of America: *Geological Society of America Centennial Field Guide*, v. 3, p. 365-370.
- Tri-City Paleontology Conference, 1992, Field guidebook, first annual Tri-City Paleontology Conference, February 14-16, 1992: University of Cincinnati, Department of Geology.
- Tuckey, Michael E., and Anstey, Robert L., 1992, Late Ordovician extinctions of bryozoans: *Lethaia*, v. 25, no. 1, p. 111-117.
- Uhri, Duane Christian, 1954, The correlation of (Ordovician) Cincinnati environments by spectrographic analyses of associated fossils: M.S. thesis (unpub.), Michigan State University, 53 p.
- Ulrich, E. O., 1878a, Observations on fossil annelids, and descriptions of some new forms: *Journal of the Cincinnati Society of Natural History*, v. 1, no. 2, p. 87-91, pl. 4.
- Ulrich, E. O., 1878b, Descriptions of some new species of fossils, from the Cincinnati Group: *Journal of the Cincinnati Society of Natural History*, v. 1, no. 2, p. 92-100, pl. 4.
- Ulrich, E. O., 1879a, Descriptions of new genera and species of fossils from the Lower Silurian about Cincinnati: *Journal of the Cincinnati Society of Natural History*, v. 2, no. 1, p. 8-30, pl. 7.
- Ulrich, E. O., 1879b, Description of a new genus and some new species of bryozoans from the Cincinnati Group: *Journal of the Cincinnati Society of Natural History*, v. 2, no. 3, p. 119-131, pl. 12.
- Ulrich, E. O., 1880, Catalogue of fossils occurring in the Cincinnati Group, of Ohio, Indiana & Kentucky: Cincinnati, Ohio, James Barclay, 31 p.
- Ulrich, E. O., 1882a, American Paleozoic Bryozoa: *Journal of the Cincinnati Society of Natural History*, v. 5, no. 3, p. 121-175, pls. 6-8; no. 4, p. 232-257, pls. 10-11.
- Ulrich, E. O., 1882b, Descriptions of two new species of crinoids: *Journal of the Cincinnati Society of Natural History*, v. 5, no. 3, p. 175-177, pl. 5.
- Ulrich, E. O., 1883, American Paleozoic Bryozoa (continued): *Journal of the Cincinnati Society of Natural History*, v. 6, no. 1, p. 82-92, pl. 1; no. 2, p. 148-168, pls. 6-7; no. 4, p. 245-279 pls. 12-14.
- Ulrich, E. O., 1884, American Paleozoic Bryozoa (continued): *Journal of the Cincinnati Society of Natural History*, v. 7, no. 1, p. 24-51, pls. 1-3.
- Ulrich, E. O., 1886, Report on the Lower Silurian Bryozoa with preliminary descriptions of some new species: *Geological and Natural History Survey of Minnesota*, 14th Annual Report for 1885, p. 57-103.
- Ulrich, E. O., 1888a, A correlation of the Lower Silurian horizons of Tennessee and of the Ohio and Mississippi valleys with those of New York and Canada: *American Geologist*, v. 1, no. 2, p. 100-110.
- Ulrich, E. O., 1888b, A correlation of the Lower Silurian horizons of Tennessee and of the Ohio and Mississippi valleys with those of New York and Canada [part 2]: *American Geologist*, v. 1, no. 3, p. 179-190.
- Ulrich, E. O., 1888c, A correlation of the Lower Silurian horizons of Tennessee and of the Ohio and Mississippi valleys with those of New York and Canada [part 3]: *American Geologist*, v. 1, no. 5, p. 305-315.
- Ulrich, E. O., 1888d, A correlation of the Lower Silurian horizons of Tennessee and of the Ohio and Mississippi valleys with those of New York and Canada [part 4]: *American Geologist*, v. 2, no. 1, p. 39-44.
- Ulrich, E. O., 1888e, Nomenclature of some Cincinnati group fossils: *American Geologist*, v. 1, no. 5, p. 333-335.
- Ulrich, E. O., 1889a, Preliminary description of new Lower Silurian sponges: *American Geologist*, v. 3, no. 4, p. 233-248.
- Ulrich, E. O., 1889b, On *Lingulasma*, a new genus, and eight new species of *Lingula* and *Trematis*: *American Geologist*, v. 3, no. 6, p. 377-391; v. 4, no. 1, p. 21-25.
- Ulrich, E. O., 1890a, New Lower Silurian Bryozoa: *Journal of the Cincinnati Society of Natural History*, v. 12, no. 4, p. 173-198.
- Ulrich, E. O., 1890b, New and little known American Paleozoic Ostracoda: *Journal of the Cincinnati Society of Natural History*, v. 13, no. 3, p. 104-137, pls. 7-10.
- Ulrich, E. O., 1890c, New Lamellibranchiata: *American Geologist*, v. 5, no. 5, p. 270-284; v. 6, no. 3, p. 173-181; v. 6, no. 5, p. 382-389.
- Ulrich, E. O., 1890d, Paleozoic Bryozoa: *Illinois Geological Survey*, v. 8, pt. 2, sec. 6, p. 285-688, pls. 29-78.
- Ulrich, E. O., 1891, New and little known American Paleozoic Ostracoda (continued): *Journal of the Cincinnati Society of Natural History*, v. 13, no. 4, p. 173-211, pls. 12-18.
- Ulrich, E. O., 1892, New Silurian Lamellibranchiata chiefly from Minnesota rocks: *Geological and Natural History Survey of Minnesota*, 19th Annual Report for 1890, p. 211-248.
- Ulrich, E. O., 1893a, New and little known Lamellibranchiata from the Lower Silurian rocks of Ohio and adjacent states: *Geological Survey of Ohio*, v. 7, p. 627-693, pls. 45-56.
- Ulrich, E. O., 1893b, On Lower Silurian Bryozoa of Minnesota: *Minnesota Geological and Natural History*, Final Report, v. 3, pt. 1, p. 96-332, 28 pls.
- Ulrich, E. O., 1895, On Lower Silurian Bryozoa of Minnesota: *Minnesota Geological Survey Final Report*, v. 3, pt. 2, p. 96-332.
- Ulrich, E. O., 1896, Bryozoa, in Zittel, K. A., von (Eastman, C. R., translator and ed.), *Text-book of paleontology*: New York, MacMillan and Co., v. 1, p. 257-291, text figs. 411-488.
- Ulrich, E. O., 1897a, The Lower Silurian Lamellibranchiata of Minnesota: *Minnesota Geological Survey Final Report*, v. 3, pt. 2, p. 475-628.
- Ulrich, E. O., 1897b, The Lower Silurian Ostracoda of Minnesota: *Minnesota Geological Survey Final Report*, v. 3, pt. 2, p. 629-693.
- Ulrich, E. O., 1911, Revision of the Paleozoic Systems: *Geological Society of America Bulletin*, v. 22, p. 281-680, pls. 25-29.
- Ulrich, E. O., and Bassler, R. S., 1904a, A revision of the Paleozoic Bryozoa, pt. 1, on genera and species of Ctenostomata: *Smithsonian Miscellaneous Collections*, v. 45, p. 256-294, 4 pls.
- Ulrich, E. O., and Bassler, R. S., 1904b, A revision of the Paleozoic Bryozoa, pt. 2, on genera and species of Trepostomata: *Smithsonian Miscellaneous Collections*, v. 47, no. 1470, p. 15-55, pls. 6-14.
- Ulrich, E. O., and Bassler, R. S., 1914, Report on the stratigraphy of the Cincinnati, Ohio quadrangle: U.S. Geological Survey open-file report, Washington, D.C., 122 p. (incomplete manuscript)

- of text for the never-published Cincinnati Folio, apparently completed to its present state early in 1914; there is a copy made in 1951 in the Geology/Physics Library at the University of Cincinnati).
- U.S. Army Corps of Engineers, no date, Caesar Creek Lake: U.S. Army Corps of Engineers, Louisville District, pamphlet.
- U.S. Geological Survey, 1969, Cincinnati Arch Ordovician pelecypods, in Geological Survey research 1969: U.S. Geological Survey Professional Paper 650-A, p. 132.
- University of Dayton, Department of Geology, 1989, The 1989 Ohio Intercollegiate Geology Field Trip, October 14, 1989: 14 p.
- Utgaard, John Edward, 1963, Trepostomatous bryozoan fauna of the upper part of the Whitewater Formation (Cincinnatian) of eastern Indiana and western Ohio: Ph.D. dissertation (unpub.), Indiana University, 264 p.
- Utgaard, John, 1968a, A revision of North American genera of ceramoporoid bryozoans (Ectoprocta)—Part I, Anolotichiidae: *Journal of Paleontology*, v. 42, no. 4, p. 1033-1041, pls. 129-132.
- Utgaard, John, 1968b, A revision of North American genera of ceramoporoid bryozoans (Ectoprocta)—Part II, *Crepipora*, *Ceramoporella*, *Acanthoceramoporella*, and *Ceramophylla*: *Journal of Paleontology*, v. 42, no. 6, p. 1444-1455, pls. 181-184.
- Utgaard, John, 1969, A revision of North American genera of ceramoporoid bryozoans (Ectoprocta)—Part III, the ceramoporoid genera *Ceramopora*, *Papillalunaria*, *Favositelia*, and *Haplotrypa*: *Journal of Paleontology*, v. 43, no. 2, p. 289-297, pls. 51-54.
- Utgaard, John, 1973, Mode of colony growth, autozooids, and polymorphism in the bryozoan order Cystoporata, in Boardman, Richard S., Cheetham, Alan H., and Oliver, William A., Jr., eds., *Animal colonies*: Stroudsburg, Pennsylvania, Dowden, Hutchinson and Ross, p. 317-360.
- Utgaard, John, and Boardman, R. S., 1965, *Heterotrypa* Nicholson, 1879, and *Peronopora* Nicholson, 1881, Bryozoa, Trepostomata: proposed designation of a type species in conformity with generally accepted usage: *Bulletin of Zoological Nomenclature*, v. 22, pt. 2, p. 112-118.
- Utgaard, John E., and Perry, T. G., 1964a, Trepostomatous bryozoan fauna of the upper part of the Whitewater Formation (Cincinnatian) of eastern Indiana and western Ohio (abs.): *Geological Society of America Special Paper* 76, p. 171.
- Utgaard, John E., and Perry, T. G., 1964b, Trepostomatous bryozoan fauna of the upper part of the Whitewater Formation (Cincinnatian) of eastern Indiana and western Ohio: *Indiana Geological Survey Bulletin* 33, 111 p., 23 pls.
- Van Der Voo, Robert, 1988, Paleozoic paleogeography of North America, Gondwana, and intervening displaced terranes: comparisons of paleomagnetism with paleoclimatology and biogeographical patterns: *Geological Society of America Bulletin*, v. 100, no. 3, p. 311-324.
- Van Fossen, John Doan, 1951, A study of the Rafinesquinae of the Middle Maysville (Upper Ordovician): M.S. thesis (unpub.), University of Cincinnati, 98 p.
- Van Hart, Dirk, 1966, The physical stratigraphy of the Saluda and Whitewater Formations (Cincinnatian Series), southeastern Indiana: M.S. thesis (unpub.), Miami University, 142 p.
- Van Iten, Heyo, Fitzke, Julie Ann, and Cox, Robt S., 1996, Problematical fossil cnidarians from the Upper Ordovician of the north-central USA: *Palaeontology*, v. 39, no. 4, p. 1037-1064, 3 pls.
- Vanuxem, Lardner, 1829, Remarks on the characters and classification of certain American rock formations: *American Journal of Science*, v. 16, no. 2, p. 254-256.
- Velbel, Danita Brandt. See Brandt Velbel, Danita.
- Versfelt, Porter LaRoy, Jr., 1953, Richmond sections in Ripley County, Indiana: M.S. thesis (unpub.), Miami University, 48 p.
- Wahlman, Gregory Paul, 1984, Middle and Upper Ordovician Monoplacophora and bellerophonitacean Gastropoda of the Cincinnati Arch region: Ph.D. dissertation (unpub.), University of Cincinnati, 678 p., 45 pls.
- Wahlman, G. P., 1992, Middle and Upper Ordovician symmetrical univalved mollusks (Monoplacophora and Bellerophonitina) of the Cincinnati Arch region: U.S. Geological Survey Professional Paper 1066-O, 213 p., 45 pls.
- Walker, Eugene H., 1953, Geology and ground-water resources of the Covington-Newport alluvial area, Kentucky: U.S. Geological Survey Circular 240, 26 p., 2 pls.
- Walker, Laurence G., 1982, The brachiopod genera *Herbertella*, *Dalmanella* and *Heterorthis* from the Ordovician of Kentucky: U.S. Geological Survey Professional Paper 1066-M, 17 p., 5 pls.
- Walter, Otto Theodore, 1925?, Trilobites of Iowa and some related Paleozoic forms: *Iowa Geological Survey*, v. 31, Annual Reports, 1923 and 1924, with accompanying papers, p. 167-400, pls. X-XXVII.
- Warn, John Michael, 1971, Variation in the Cincinnatian (Upper Ordovician) crinoid *Heterocrinus heterodactylus* Hall: ontogeny, regeneration, and pathology: M.S. thesis (unpub.), University of Cincinnati, 127 p., 9 pls.
- Warn, John M., 1973, The Ordovician crinoid *Heterocrinus* with reference to brachial variability in *H. tenuis*: *Journal of Paleontology*, v. 47, no. 1, p. 10-18, 1 pl.
- Warn, John Michael, 1974a, The disparid crinoid superfamilies Homocrinacea (Ordovician-Silurian) and Cincinnaticrinacea (Ordovician): Ph.D. dissertation (unpub.), University of Cincinnati, 296 p. (abstract: *Dissertation Abstracts International*, sec. B, Science and Engineering, v. 35, no. 6, p. 2828B-2829B).
- Warn, John M., 1974b, Presumed myzostomid infestation of an Ordovician crinoid: *Journal of Paleontology*, v. 48, no. 3, p. 506-513, 1 pl.
- Warn, J. M., and Strimple, H. L., 1977, The disparid inadunate superfamilies Homocrinacea and Cincinnaticrinacea (Echinodermata, Crinoidea), Ordovician-Silurian, North America: *Bulletins of American Paleontology*, v. 72, no. 296, p. 1-138, pls. 1-18.
- Warshauer, Steven M., 1969, A preliminary investigation of Ostracoda from the Kope Formation (Upper Ordovician): M.S. thesis (unpub.), University of Cincinnati, 85 p.
- Warshauer, Steven M., 1973, The taxonomy, ontogeny, biostratigraphy and paleoecology of the Edenian (Upper Ordovician) ostracods of the Ohio Valley: Ph.D. dissertation (unpub.), University of Cincinnati, 222 p., 8 pls.
- Warshauer, S. M., and Berdan, Jean M., 1982, Palaeocopid and podocopid Ostracoda from the Lexington Limestone and Clays Ferry Formation (Middle and Upper Ordovician) of central Kentucky: U.S. Geological Survey Professional Paper 1066-H, 80 p., 19 pls.
- Weaver, Thomas R., 1975, Disconformable Maysville-Richmond stage boundary (Upper Ordovician) of the type area (abs.): *Geological Society of America Abstracts with Programs*, v. 7, no. 6, p. 878.
- Weaver, Thomas R., 1976, Adaptive strategies of disparid inadunate crinoids of the type Cincinnatian (Upper Ordovician) (abs.): *Geological Society of America Abstracts with Programs*, v. 8, no. 4, p. 516.
- Weidner, Jerry R., 1962, X-ray spectrochemical investigations of the Cincinnatian Series shale of Ohio, Indiana, and Kentucky: M.S. thesis (unpub.), Miami University, 96 p.
- Weir, Gordon W., and Green, Robert C., 1965, Clays Ferry Formation (Ordovician)—a new map unit in south-central Kentucky: U.S. Geological Survey Bulletin 1224-B, 18 p.
- Weir, Gordon W., Greene, Robert C., and Simmons, George C., 1965, Calloway Creek Limestone and Ashlock and Drakes Formations (Upper Ordovician) in south-central Kentucky: U.S. Geological Survey Bulletin 1224-D, 36 p.
- Weir, Gordon W., and Peck, John H., 1968, Lithofacies of Upper Ordovician rocks exposed between Maysville and Stanford, Kentucky, in Geological Survey research 1968: U.S. Geological Survey Professional Paper 600-D, p. 162-168.
- Weir, Gordon W., Peterson, Warren L., and Swadley, W C, 1979, Lithofacies and stratigraphic nomenclature of part of the Upper Ordovician section of Kentucky: U.S. Geological Survey Miscellaneous Investigations Series Map I-1155, scale 1:250,000.

- Weir, G. W., Peterson, W. L., Swadley, W. C., and Pojeta, John, Jr., 1984, Lithostratigraphy of Upper Ordovician strata exposed in Kentucky; with a section on biostratigraphy: U.S. Geological Survey Professional Paper 1151-E, 121 p., 5 pls.
- Weisenberger, B. C., Dowell, C. W., Leathers, T. R., Odor, H. B., and Richardson, A. J., 1973, Soil survey of Boone, Campbell, and Kenton Counties, Kentucky: U.S. Department of Agriculture, Soil Conservation Service, 67 p. + maps.
- Weiss, Malcolm P., 1961, The American Upper Ordovician Standard. V. A critical appraisal of the classification of the typical Cincinnati beds: Geological Society of America Bulletin, v. 72, no. 4, p. 645-648.
- Weiss, M. P., Edwards, W. R., Norman, C. E., and Sharp, E. R., 1965, The American Upper Ordovician Standard. VII. Stratigraphy and petrology of the Cynthiana and Eden Formations of the Ohio Valley: Geological Society of America Special Paper 81, 76 p., 8 pls.
- Weiss, Malcolm P., and Norman, Carl E., 1960a, The American Upper Ordovician Standard. II. Development of stratigraphic classification of Ordovician rocks in the Cincinnati region: Ohio Division of Geological Survey Information Circular 26, 14 p., 1 pl.
- Weiss, Malcolm P., and Norman, Carl E., 1960b, The American Upper Ordovician Standard. IV. Classification of the limestones of the type Cincinnati: Journal of Sedimentary Petrology, v. 30, no. 2, p. 283-296.
- Weiss, M. P., Schilling, F. A., Pierce, K. L. and Ali, S. A., 1972, Geologic map of the Maysville East quadrangle, Ohio-Kentucky: U.S. Geological Survey Geologic Quadrangle Map GQ-1006, scale 1:24,000.
- Weiss, M. P., and Sweet, W. C., 1962, Contributions of facies analysis to Ordovician history of Cincinnati Arch geologic province (abs.): Geological Society of America Special Paper 68, p. 294.
- Weiss, Malcolm P., and Sweet, Walter C., 1964, Kope Formation (Upper Ordovician): Ohio and Kentucky: Science, v. 145, no. 3638, p. 1296-1302.
- Welch, James R., 1976, *Phosphannulus* on Paleozoic crinoid stems: Journal of Paleontology, v. 50, no. 2, p. 218-225, 2 pls.
- Wells, John W., 1942, Supposed color-markings in Ordovician trilobites from Ohio: American Journal of Science, v. 240, no. 10, p. 710-713, 1 pl.
- Wetherby, A. G., 1879, Description of a new family and genus of Lower Silurian Crustacea: Journal of the Cincinnati Society of Natural History, v. 1, no. 4, p. 162-166.
- Wetherby, A. G., 1880, Descriptions of new crinoids from the Cincinnati Group of the Lower Silurian and the Subcarboniferous of Kentucky: Journal of the Cincinnati Society of Natural History, v. 2, no. 4, p. 245-253, pl. 16.
- Wetherby, A. G., 1881, Description of new fossils from the Lower Silurian and Subcarboniferous rocks of Ohio and Kentucky: Journal of the Cincinnati Society of Natural History, v. 4, no. 1, p. 77-85, pl. 2.
- Wetzel, John M., 1968, Petrography of selected limestones of the Fairview Formation and Bellevue Member, McMillan Formation (Cincinnati Series) from southwestern Hamilton County, Ohio: M.S. thesis (unpub.), Miami University, 93 p., 1 pl., 38 text-figs.
- White, C. A., 1882, Van Cleve's fossil corals: Indiana Department of Geology and Natural Resources Eleventh Annual Report, p. 376-401, pls. 44-45.
- Whitfield, R. P., 1878, Remarks on some lamellibranchiate shells of the Hudson River Group, with descriptions of four new species: Journal of the Cincinnati Society of Natural History, v. 1, no. 3, p. 137-141, pl. 6.
- Whittington, Harry B., 1956, Type and other species of *Odontopleuridae* (Trilobita): Journal of Paleontology, v. 30, no. 3, p. 504-520, pls. 57-60.
- Whittington, H. B., 1965, *Platycoryphe*, an Ordovician homalonotid trilobite: Journal of Paleontology, v. 39, no. 3, p. 487-491, pl. 64.
- Whittington, H. B., 1968, *Cryptolithus* (Trilobita): specific characters and occurrence in Ordovician of eastern North America: Journal of Paleontology, v. 42, no. 3, p. 702-714, pls. 87-89.
- Wilkie, Lorna Christine, 1957, The conodont fauna of the Upper Ordovician McMicken Member, southwestern Ohio: M.S. thesis (unpub.), The Ohio State University, 148 p., 5 pls.
- Williams, Stephen R., 1914, A starfish found in the Whitewater Division of the Richmond on Blue Creek, Adams County, Ohio: Ohio Naturalist, v. 14, no. 3, p. 221-224.
- Williams, Stephen R., 1918, Concerning the structure of *Agelacrinites* and *Streptaster*, Edrioasteroidea of the Richmond and Maysville divisions of the Ordovician: Ohio Journal of Science, v. 19, no. 1, p. 59-82.
- Wilmarth, M. Grace, compiler, 1938, Lexicon of geologic names of the United States (including Alaska): U.S. Geological Survey Bulletin 896, 2 v., 2,396 p.
- Wilson, Mark A., 1985, Disturbance and ecologic succession in an Upper Ordovician cobble-dwelling hardground fauna: Science, v. 228, no. 4699, p. 575-577.
- Wilson, Mark A., and Norrish, Winston A., 1984, Paleoecology of the Upper Ordovician ichnofossil *Diplocraterion* in northern Kentucky (abs.): Geological Society of America Abstracts with Programs, v. 16, no. 3, p. 206.
- Wilson, Mark A., and Palmer, Timothy J., 1988, Nomenclature of a bivalve boring from the Upper Ordovician of the midwestern United States: Journal of Paleontology, v. 62, no. 2, p. 306-308.
- Wilson, Mark A., Palmer, Timothy J., and Taylor, Paul D., 1994, Earliest preservation of soft-bodied fossils by epibiont bioimmuration, Upper Ordovician of Kentucky: Lethaia, v. 27, no. 3, p. 269-270.
- Winchell, N. H., and Ulrich, E. O., 1897, The Lower Silurian deposits of the upper Mississippi province: a correlation of the strata with those in the Cincinnati, Tennessee, New York and Canadian Provinces, and the stratigraphic and geographic distribution of the fossils, in Ulrich, E. O., and others, The geology of Minnesota: Geological and Natural History Survey of Minnesota, v. 3, pt. 2 of Final Report, Paleontology, p. lxxxiii-cxxviii.
- Winder, Charles Gordon, 1953, Paleoecology and sedimentation of (Ordovician) Mohawkian limestones in south-central Ontario; a revision of Mohawkian and Cincinnati stratigraphy: Ph.D. dissertation (unpub.), Cornell University, 109 p.
- Wolford, John J., 1927, The geology of the Oregonia-Fort Ancient region, Warren County, Ohio: M.A. thesis (unpub.), The Ohio State University, 56 p., 19 pls.
- Wolford, J. J., 1930, The stratigraphy of the Oregonia-Fort Ancient region, southwestern Ohio: Ohio Journal of Science, v. 30, no. 5, p. 301-308.
- Woodward, Henry, 1884, Notes on appendages of trilobites: Geological Magazine, ser. 3, v. 1, no. 4, p. 162-165.
- Yang, King-Chih, 1947, The Bryozoa of the Greendale Member, Cynthiana Formation, Kentucky: M.A. thesis (unpub.), Miami University, 41 p.
- Ziegler, Willi, ed., 1973-91, Catalogue of conodonts: Stuttgart, Schwelbarts'sche Verlagsbuchhandlung, v. 1-5.

Addresses of authors: Richard Arnold Davis, Biology Department, College of Mount St. Joseph, 5701 Delhi Road, Cincinnati, OH, 45233-1670; Richard A. Spohn, Geology/Physics Library, University of Cincinnati, Cincinnati, OH 45221-0153.

**MAP OF THE TRI-STATE AREA SHOWING
MAJOR LOCATIONS DESCRIBED IN THIS GUIDEBOOK**

